

Checklist of Yorkshire Coleoptera

Figures based on records on the YNU database in January 2011

Figures after family names are British species/Yorkshire species

| <u>Taxon</u> | <u>VC61</u> | <u>VC62</u> | <u>VC63</u> | <u>VC64</u> | <u>VC65</u> | <u>VC Total</u> |
|---|-------------|-------------|-------------|-------------|-------------|-----------------|
| <u>Gyrinidae [12/9]</u> | | | | | | |
| Gyrinus aeratus Stephens, 1835 | 10 | 0 | 2 | 0 | 0 | 12 |
| Gyrinus caspius Ménétrés, 1832 | 5 | 1 | 2 | 1 | 0 | 9 |
| Gyrinus distinctus Aubé, 1838 | 1 | 0 | 2 | 0 | 0 | 3 |
| Gyrinus marinus Gyllenhal, 1808 | 22 | 4 | 28 | 8 | 0 | 62 |
| Gyrinus minutus Fabricius, 1798 | 0 | 2 | 3 | 0 | 0 | 5 |
| Gyrinus paykulli Ochs, 1927 | 0 | 1 | 1 | 1 | 0 | 3 |
| Gyrinus substriatus Stephens, 1828 | 22 | 14 | 79 | 20 | 4 | 139 |
| Gyrinus urinator Illiger, 1807 | 0 | 3 | 1 | 0 | 0 | 4 |
| Orectochilus villosus (Müller, O.F., 1776) | 1 | 5 | 1 | 19 | 2 | 28 |
| <u>Haliplidae [19/17]</u> | | | | | | |
| Brychius elevatus (Panzer, 1793) | 2 | 2 | 4 | 7 | 1 | 16 |
| Peltodytes caesus (Duftschmid, 1805) | 0 | 0 | 2 | 0 | 0 | 2 |
| Haliplus (Haliplus) confinis Stephens, 1828 | 15 | 5 | 37 | 12 | 2 | 71 |
| Haliplus (Haliplus) obliquus (Fabricius, 1787) | 18 | 4 | 14 | 18 | 4 | 58 |
| Haliplus (Neohalipus) lineatocollis (Marsham, 1802) | 24 | 12 | 63 | 11 | 3 | 113 |
| Haliplus (Haliplinus) apicalis Thomson, C.G., 1868 | 6 | 2 | 1 | 0 | 0 | 9 |
| Haliplus (Haliplinus) fluviatilis Aubé, 1836 | 5 | 2 | 32 | 4 | 0 | 43 |
| Haliplus (Haliplinus) heydeni Wehncke, 1875 | 2 | 2 | 6 | 3 | 0 | 13 |
| Haliplus (Haliplinus) immaculatus Gerhardt, 1877 | 19 | 3 | 48 | 11 | 0 | 81 |
| Haliplus (Haliplinus) lineolatus Mannerheim, 1844 | 13 | 3 | 25 | 9 | 0 | 50 |
| Haliplus (Haliplinus) ruficollis (De Geer, 1774) | 25 | 14 | 82 | 29 | 2 | 152 |
| Haliplus (Haliplinus) sibiricus Motschulsky, 1860 | 44 | 22 | 34 | 20 | 4 | 124 |
| Haliplus (Liaphlus) flavicollis Sturm, 1834 | 11 | 1 | 18 | 17 | 5 | 52 |
| Haliplus (Liaphlus) fulvus (Fabricius, 1801) | 7 | 3 | 7 | 7 | 1 | 25 |
| Haliplus (Liaphlus) laminatus (Schaller, 1783) | 0 | 0 | 9 | 6 | 0 | 15 |
| Haliplus (Liaphlus) mucronatus Stephens, 1828 | 1 | 0 | 1 | 0 | 0 | 2 |
| Haliplus (Liaphlus) variegatus Sturm, 1834 | 1 | 1 | 2 | 0 | 0 | 4 |
| <u>Noteridae [2/2]</u> | | | | | | |
| Noterus clavicornis (De Geer, 1774) | 37 | 6 | 66 | 29 | 1 | 139 |
| Noterus crassicornis (Müller, O.F., 1776) | 5 | 2 | 10 | 2 | 0 | 19 |
| <u>Paelobiidae [1/1]</u> | | | | | | |
| Hygrobia hermanni (Fabricius, 1775) | 17 | 4 | 19 | 6 | 1 | 47 |
| <u>Dytiscidae [117/103]</u> | | | | | | |
| Agabus (Acatodes) arcticus (Paykull, 1798) | 0 | 0 | 15 | 8 | 4 | 27 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| Agabus (Acatodes) congener (Thunberg, 1794) | 4 | 2 | 7 | 10 | 4 | 27 |
| Agabus (Acatodes) sturmii (Gyllenhal, 1808) | 28 | 13 | 83 | 30 | 5 | 159 |
| Agabus (Agabus) labiatus (Brahm, 1790) | 6 | 2 | 2 | 1 | 0 | 11 |
| Agabus (Agabus) uliginosus (Linnaeus, 1761) | 9 | 9 | 7 | 12 | 1 | 38 |
| Agabus (Agabus) undulatus (Schrank, 1776) | 4 | 1 | 0 | 15 | 0 | 20 |
| Agabus (Gaurodytes) affinis (Paykull, 1798) | 4 | 2 | 6 | 3 | 0 | 15 |
| Agabus (Gaurodytes) biguttatus (Olivier, 1795) | 1 | 3 | 11 | 3 | 0 | 18 |
| Agabus (Gaurodytes) bipustulatus (Linnaeus, 1767) | 84 | 36 | 184 | 74 | 6 | 384 |
| Agabus (Gaurodytes) conspersus (Marsham, 1802) | 16 | 2 | 3 | 0 | 0 | 21 |
| Agabus (Gaurodytes) didymus (Olivier, 1795) | 21 | 8 | 35 | 13 | 3 | 80 |
| Agabus (Gaurodytes) guttatus (Paykull, 1798) | 0 | 12 | 64 | 21 | 3 | 100 |
| Agabus (Gaurodytes) melanarius Aubé, 1837 | 1 | 3 | 1 | 0 | 0 | 5 |
| Agabus (Gaurodytes) nebulosus (Forster, 1771) | 33 | 16 | 65 | 32 | 2 | 148 |
| Agabus (Gaurodytes) paludosus (Fabricius, 1801) | 10 | 11 | 27 | 14 | 0 | 62 |
| Agabus (Gaurodytes) striolatus (Gyllenhal, 1808) | 0 | 0 | 1 | 0 | 0 | 1 |
| Agabus (Gaurodytes) unguicularis (Thomson, C.G., 1867) | 0 | 3 | 6 | 8 | 0 | 17 |
| Ilybius aenescens Thomson, C.G., 1870 | 5 | 7 | 7 | 5 | 0 | 24 |
| Ilybius ater (De Geer, 1774) | 17 | 5 | 47 | 16 | 4 | 89 |
| Ilybius chalconatus (Panzer, 1796) | 10 | 17 | 12 | 10 | 1 | 50 |
| Ilybius fenestratus (Fabricius, 1781) | 7 | 2 | 18 | 4 | 0 | 31 |
| Ilybius fuliginosus (Fabricius, 1792) | 31 | 12 | 93 | 28 | 11 | 175 |
| Ilybius guttiger (Gyllenhal, 1808) | 8 | 2 | 7 | 7 | 0 | 24 |
| Ilybius montanus (Stephens, 1828) | 2 | 2 | 4 | 6 | 2 | 16 |
| Ilybius quadriguttatus (Lacordaire, 1835) | 16 | 2 | 24 | 8 | 0 | 50 |
| Ilybius subaeneus Erichson, 1837 | 1 | 2 | 8 | 1 | 0 | 12 |
| Platambus maculatus (Linnaeus, 1758) | 6 | 12 | 30 | 21 | 6 | 75 |
| Colymbetes fuscus (Linnaeus, 1758) | 43 | 9 | 55 | 23 | 2 | 132 |
| Rhantus (Nartus) grapii (Gyllenhal, 1808) | 4 | 1 | 14 | 9 | 0 | 28 |
| Rhantus (Rhantus) bistriatus (Bergsträsser, 1778) | 0 | 0 | 1 | 0 | 0 | 1 |
| Rhantus (Rhantus) exsoletus (Forster, 1771) | 5 | 2 | 17 | 5 | 0 | 29 |
| Rhantus (Rhantus) frontalis (Marsham, 1802) | 1 | 0 | 0 | 1 | 0 | 2 |
| Rhantus (Rhantus) suturalis (MacLeay, 1825) | 2 | 1 | 8 | 6 | 0 | 17 |
| Rhantus (Rhantus) suturellus (Harris, T.W., 1828) | 13 | 6 | 9 | 5 | 0 | 33 |
| Liopterus haemorrhoidalis (Fabricius, 1787) | 14 | 0 | 19 | 11 | 0 | 44 |
| Acilius canaliculatus (Nicolai, 1822) | 3 | 0 | 26 | 0 | 0 | 29 |
| Acilius sulcatus (Linnaeus, 1758) | 9 | 12 | 27 | 9 | 0 | 57 |
| Dytiscus circumcinctus Ahrens, 1811 | 3 | 1 | 12 | 9 | 0 | 25 |
| Dytiscus circumflexus Fabricius, 1801 | 4 | 0 | 4 | 1 | 0 | 9 |
| Dytiscus dimidiatus Bergsträsser, 1778 | 1 | 1 | 2 | 11 | 0 | 15 |
| Dytiscus marginalis Linnaeus, 1758 | 22 | 11 | 73 | 14 | 3 | 123 |
| Dytiscus semisulcatus Müller, O.F., 1776 | 6 | 4 | 13 | 6 | 1 | 30 |
| Hydaticus continentalis Balfour-Browne, J., 1944 | 0 | 0 | 1 | 0 | 0 | 1 |
| Hydaticus seminiger (De Geer, 1774) | 0 | 0 | 0 | 4 | 0 | 4 |
| Hydaticus transversalis (Pontoppidan, 1763) | 0 | 0 | 2 | 7 | 0 | 9 |
| Bidessus unistriatus (Goeze, 1777) | 0 | 0 | 1 | 0 | 0 | 1 |
| Hydroglyphus geminus (Fabricius, 1792) | 4 | 1 | 12 | 4 | 0 | 21 |
| Deronectes latus (Stephens, 1829) | 0 | 4 | 2 | 2 | 0 | 8 |
| Graptodytes bilineatus (Sturm, 1835) | 6 | 0 | 0 | 0 | 0 | 6 |
| Graptodytes granularis (Linnaeus, 1767) | 3 | 0 | 7 | 5 | 0 | 15 |
| Graptodytes pictus (Fabricius, 1787) | 22 | 7 | 48 | 2 | 0 | 79 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| Hydroporus angustatus Sturm, 1835 | 11 | 5 | 26 | 19 | 0 | 61 |
| Hydroporus discretus Fairmaire & Brisout, 1859 | 10 | 6 | 16 | 11 | 1 | 44 |
| Hydroporus erythrocephalus (Linnaeus, 1758) | 25 | 3 | 22 | 27 | 0 | 77 |
| Hydroporus ferrugineus Stephens, 1829 | 0 | 5 | 3 | 3 | 1 | 12 |
| Hydroporus glabriusculus Aubé, 1838 | 0 | 0 | 0 | 1 | 0 | 1 |
| Hydroporus gyllenhalii Schiödte, 1841 | 13 | 13 | 23 | 23 | 4 | 76 |
| Hydroporus incognitus Sharp, 1869 | 1 | 5 | 13 | 5 | 0 | 24 |
| Hydroporus longicornis Sharp, 1871 | 0 | 1 | 1 | 2 | 2 | 6 |
| Hydroporus longulus Mulsant & Rey, 1861 | 0 | 2 | 5 | 2 | 3 | 12 |
| Hydroporus marginatus (Duftschmid, 1805) | 0 | 0 | 1 | 0 | 0 | 1 |
| Hydroporus melanarius Sturm, 1835 | 5 | 11 | 12 | 12 | 3 | 43 |
| Hydroporus memnonius Nicolai, 1822 | 14 | 17 | 27 | 18 | 3 | 79 |
| Hydroporus morio Aubé, 1838 | 0 | 2 | 0 | 16 | 4 | 22 |
| Hydroporus neglectus Schaum, 1845 | 3 | 1 | 6 | 3 | 0 | 13 |
| Hydroporus nigrita (Fabricius, 1792) | 7 | 11 | 20 | 16 | 5 | 59 |
| Hydroporus obscurus Sturm, 1835 | 6 | 3 | 3 | 13 | 2 | 27 |
| Hydroporus obsoletus Aubé, 1838 | 0 | 2 | 0 | 0 | 1 | 3 |
| Hydroporus palustris (Linnaeus, 1761) | 58 | 16 | 158 | 54 | 7 | 293 |
| Hydroporus planus (Fabricius, 1781) | 47 | 17 | 107 | 25 | 2 | 198 |
| Hydroporus pubescens (Gyllenhal, 1808) | 35 | 22 | 81 | 34 | 4 | 176 |
| Hydroporus rufifrons (Müller, O.F., 1776) | 0 | 4 | 3 | 8 | 0 | 15 |
| Hydroporus scalesianus Stephens, 1828 | 0 | 0 | 3 | 3 | 0 | 6 |
| Hydroporus striola (Gyllenhal, 1826) | 6 | 3 | 14 | 13 | 0 | 36 |
| Hydroporus tessellatus (Drapiez, 1819) | 15 | 13 | 24 | 14 | 0 | 66 |
| Hydroporus tristis (Paykull, 1798) | 9 | 18 | 11 | 22 | 4 | 64 |
| Hydroporus umbrosus (Gyllenhal, 1808) | 9 | 10 | 13 | 13 | 1 | 46 |
| Nebrioporus (Nebrioporus) assimilis (Paykull, 1798) | 16 | 3 | 22 | 11 | 0 | 52 |
| Nebrioporus (Nebrioporus) depressus (Fabricius, 1775) | 0 | 0 | 0 | 1 | 0 | 1 |
| Nebrioporus (Nebrioporus) elegans (Panzer, 1794) | 7 | 2 | 21 | 16 | 3 | 49 |
| Oreodytes davisii (Curtis, 1831) | 0 | 6 | 6 | 9 | 7 | 28 |
| Oreodytes sanmarkii (Sahlberg, C.R., 1826) | 2 | 19 | 20 | 19 | 11 | 71 |
| Oreodytes septentrionalis (Gyllenhal, 1826) | 0 | 5 | 3 | 6 | 4 | 18 |
| Porhydrus lineatus (Fabricius, 1775) | 12 | 2 | 14 | 5 | 0 | 33 |
| Scarodytes halensis (Fabricius, 1787) | 5 | 0 | 5 | 5 | 0 | 15 |
| Stictonectes lepidus (Olivier, 1795) | 2 | 5 | 33 | 13 | 0 | 53 |
| Stictotarsus duodecimpustulatus (Fabricius, 1792) | 6 | 6 | 24 | 9 | 1 | 46 |
| Stictotarsus multilineatus (Falkenström, 1922) | 0 | 0 | 0 | 1 | 3 | 4 |
| Suphrodytes dorsalis (Fabricius, 1787) | 14 | 0 | 15 | 22 | 0 | 51 |
| Hygrotus (Coelambus) confluens (Fabricius, 1787) | 12 | 5 | 13 | 7 | 0 | 37 |
| Hygrotus (Coelambus) impressopunctatus (Schaller, 1783) | 27 | 6 | 64 | 25 | 1 | 123 |
| Hygrotus (Coelambus) nigrolineatus (Steven, 1808) | 0 | 0 | 1 | 1 | 0 | 2 |
| Hygrotus (Coelambus) novemlineatus (Stephens, 1829) | 0 | 2 | 0 | 0 | 3 | 5 |
| Hygrotus (Coelambus) parallelogrammus (Ahrens, 1812) | 8 | 0 | 0 | 0 | 0 | 8 |
| Hygrotus (Hygrotus) decoratus (Gyllenhal, 1810) | 2 | 2 | 0 | 7 | 0 | 11 |
| Hygrotus (Hygrotus) inaequalis (Fabricius, 1777) | 45 | 7 | 86 | 38 | 3 | 179 |
| Hygrotus (Hygrotus) quinquelineatus (Zetterstedt, 1828) | 0 | 0 | 3 | 1 | 0 | 4 |
| Hygrotus (Hygrotus) versicolor (Schaller, 1783) | 20 | 0 | 29 | 5 | 0 | 54 |
| Hyphydrus ovatus (Linnaeus, 1761) | 35 | 4 | 71 | 21 | 0 | 131 |
| Laccornis oblongus (Stephens, 1835) | 0 | 1 | 0 | 5 | 0 | 6 |
| Laccophilus hyalinus (De Geer, 1774) | 10 | 2 | 14 | 8 | 0 | 34 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|-----|----|-----|
| <i>Laccophilus minutus</i> (Linnaeus, 1758) | 26 | 11 | 56 | 36 | 1 | 130 |
| <i>Laccophilus poecilus</i> Klug, 1834 | 0 | 0 | 4 | 0 | 0 | 4 |
| <u>Carabidae [368/270]</u> | | | | | | |
| <i>Cicindela campestris</i> Linnaeus, 1758 | 11 | 41 | 115 | 13 | 8 | 188 |
| <i>Brachinus</i> (<i>Brachinus</i>) <i>crepitans</i> (Linnaeus, 1758) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Carabus</i> (<i>Eucarabus</i>) <i>arvensis</i> Herbst, 1784 | 2 | 32 | 2 | 7 | 3 | 46 |
| <i>Carabus</i> (<i>Carabus</i>) <i>granulatus</i> Linnaeus, 1758 | 20 | 3 | 26 | 4 | 0 | 53 |
| <i>Carabus</i> (<i>Morphocarabus</i>) <i>monilis</i> Fabricius, 1792 | 4 | 8 | 9 | 12 | 0 | 33 |
| <i>Carabus</i> (<i>Archicarabus</i>) <i>nemoralis</i> Müller, O.F., 1764 | 9 | 19 | 36 | 30 | 5 | 99 |
| <i>Carabus</i> (<i>Autocarabus</i>) <i>auratus</i> Linnaeus, 1761 | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Carabus</i> (<i>Autocarabus</i>) <i>cancellatus</i> Illiger, 1798 | 1 | 0 | 1 | 2 | 1 | 5 |
| <i>Carabus</i> (<i>Hemicarabus</i>) <i>nitens</i> Linnaeus, 1758 | 2 | 40 | 12 | 9 | 18 | 81 |
| <i>Carabus</i> (<i>Oreocarabus</i>) <i>glabratus</i> Paykull, 1790 | 0 | 10 | 1 | 0 | 3 | 14 |
| <i>Carabus</i> (<i>Mesocarabus</i>) <i>problematicus</i> Herbst, 1786 | 6 | 33 | 70 | 19 | 18 | 146 |
| <i>Carabus</i> (<i>Megodontus</i>) <i>violaceus</i> Linnaeus, 1758 | 8 | 16 | 39 | 19 | 15 | 97 |
| <i>Cychrus caraboides</i> (Linnaeus, 1758) | 2 | 15 | 82 | 31 | 11 | 141 |
| <i>Leistus</i> (<i>Pogonophorus</i>) <i>rufomarginatus</i> (Duftschmid, 1812) | 1 | 6 | 33 | 3 | 1 | 44 |
| <i>Leistus</i> (<i>Pogonophorus</i>) <i>spinibarbis</i> (Fabricius, 1775) | 7 | 5 | 23 | 4 | 2 | 41 |
| <i>Leistus</i> (<i>Leistophorus</i>) <i>fulvibarbis</i> Dejean, 1826 | 15 | 7 | 75 | 35 | 5 | 137 |
| <i>Leistus</i> (<i>Leistus</i>) <i>ferrugineus</i> (Linnaeus, 1758) | 24 | 12 | 106 | 20 | 0 | 162 |
| <i>Leistus</i> (<i>Leistus</i>) <i>terminatus</i> (Hellwig in Panzer, 1793) | 16 | 27 | 72 | 44 | 7 | 166 |
| <i>Nebria</i> (<i>Paranebria</i>) <i>livida</i> (Linnaeus, 1758) | 12 | 6 | 0 | 0 | 0 | 18 |
| <i>Nebria</i> (<i>Nebria</i>) <i>brevicollis</i> (Fabricius, 1792) | 32 | 47 | 278 | 92 | 27 | 476 |
| <i>Nebria</i> (<i>Nebria</i>) <i>salina</i> Fairmaire & Laboulbène, 1854 | 2 | 15 | 50 | 39 | 16 | 122 |
| <i>Nebria</i> (<i>Boreonebria</i>) <i>rufescens</i> (Ström, 1768) | 0 | 20 | 33 | 32 | 16 | 101 |
| <i>Notiophilus aesthuans</i> Motschulsky, 1864 | 1 | 0 | 0 | 1 | 6 | 8 |
| <i>Notiophilus aquaticus</i> (Linnaeus, 1758) | 4 | 26 | 38 | 11 | 14 | 93 |
| <i>Notiophilus biguttatus</i> (Fabricius, 1779) | 76 | 57 | 298 | 102 | 34 | 567 |
| <i>Notiophilus germinyi</i> Fauvel, 1863 | 3 | 23 | 12 | 20 | 12 | 70 |
| <i>Notiophilus palustris</i> (Duftschmid, 1812) | 17 | 12 | 50 | 12 | 2 | 93 |
| <i>Notiophilus quadripunctatus</i> Dejean, 1826 | 4 | 3 | 4 | 0 | 0 | 11 |
| <i>Notiophilus rufipes</i> Curtis, 1829 | 1 | 2 | 9 | 3 | 0 | 15 |
| <i>Notiophilus substriatus</i> Waterhouse, G.R., 1833 | 11 | 3 | 30 | 15 | 1 | 60 |
| <i>Blethisa multipunctata</i> (Linnaeus, 1758) | 11 | 0 | 6 | 6 | 0 | 23 |
| <i>Elaphrus</i> (<i>Elaphrus</i>) <i>cupreus</i> Duftschmid, 1812 | 24 | 11 | 86 | 36 | 3 | 160 |
| <i>Elaphrus</i> (<i>Elaphrus</i>) <i>lapponicus</i> Gyllenhal, 1810 | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Elaphrus</i> (<i>Elaphrus</i>) <i>uliginosus</i> Fabricius, 1792 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Elaphrus</i> (<i>Trichelaphrus</i>) <i>riparius</i> (Linnaeus, 1758) | 41 | 13 | 38 | 36 | 5 | 133 |
| <i>Loricera pilicornis</i> (Fabricius, 1775) | 41 | 28 | 189 | 84 | 20 | 362 |
| <i>Clivina collaris</i> (Herbst, 1784) | 7 | 7 | 6 | 18 | 2 | 40 |
| <i>Clivina fossor</i> (Linnaeus, 1758) | 40 | 19 | 67 | 52 | 12 | 190 |
| <i>Dyschirius</i> (<i>Dyschirius</i>) <i>angustatus</i> (Ahrens, 1830) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Dyschirius</i> (<i>Dyschirius</i>) <i>thoracicus</i> (Rossi, 1790) | 13 | 1 | 0 | 0 | 0 | 14 |
| <i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>aeneus</i> (Dejean, 1825) | 6 | 1 | 4 | 4 | 0 | 15 |
| <i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>globosus</i> (Herbst, 1784) | 27 | 24 | 66 | 10 | 6 | 133 |
| <i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>luedersi</i> Wagner, 1915 | 16 | 3 | 18 | 1 | 0 | 38 |
| <i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>nitidus</i> (Dejean, 1825) | 3 | 1 | 0 | 0 | 0 | 4 |
| <i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>politus</i> (Dejean, 1825) | 7 | 1 | 3 | 1 | 1 | 13 |
| <i>Dyschirius</i> (<i>Dyschiriodes</i>) <i>salinus</i> Schaum, 1843 | 9 | 0 | 1 | 0 | 0 | 10 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Brosicus cephalotes</i> (Linnaeus, 1758) | 16 | 1 | 10 | 0 | 0 | 27 |
| <i>Miscodera arctica</i> (Paykull, 1798) | 2 | 13 | 8 | 9 | 7 | 39 |
| <i>Aepus marinus</i> (Ström, 1783) | 0 | 7 | 0 | 0 | 0 | 7 |
| <i>Aepus robinii</i> (Laboulbène, 1849) | 0 | 7 | 0 | 0 | 0 | 7 |
| <i>Trechus</i> (<i>Epaphius</i>) <i>rivularis</i> (Gyllenhal, 1810) | 0 | 0 | 0 | 8 | 0 | 8 |
| <i>Trechus</i> (<i>Epaphius</i>) <i>secalis</i> (Paykull, 1790) | 7 | 9 | 12 | 24 | 4 | 56 |
| <i>Trechus</i> (<i>Trechus</i>) <i>fulvus</i> Dejean, 1831 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Trechus</i> (<i>Trechus</i>) <i>obtusus</i> Erichson, 1837 | 45 | 32 | 94 | 39 | 14 | 224 |
| <i>Trechus</i> (<i>Trechus</i>) <i>quadristriatus</i> (Schrank, 1781) | 58 | 13 | 149 | 29 | 13 | 262 |
| <i>Trechus</i> (<i>Trechus</i>) <i>rubens</i> (Fabricius, 1792) | 0 | 5 | 14 | 2 | 2 | 23 |
| <i>Trechus</i> (<i>Trechus</i>) <i>subnotatus</i> Dejean, 1831 | 0 | 0 | 6 | 0 | 0 | 6 |
| <i>Blemus discus</i> (Fabricius, 1792) | 8 | 5 | 8 | 11 | 0 | 32 |
| <i>Trechoblemus micros</i> (Herbst, 1784) | 2 | 7 | 6 | 8 | 5 | 28 |
| <i>Tachys</i> (<i>Paratachys</i>) <i>bistriatus</i> (Duftschmid, 1812) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Tachys</i> (<i>Tachys</i>) <i>scutellaris</i> Stephens, 1828 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Elaphropus parvulus</i> (Dejean, 1831) | 0 | 0 | 12 | 0 | 0 | 12 |
| <i>Asaphidion curtum</i> (Heyden, 1870) | 12 | 2 | 25 | 15 | 0 | 54 |
| <i>Asaphidion flavipes</i> (Linnaeus, 1761) | 2 | 1 | 1 | 5 | 0 | 9 |
| <i>Asaphidion pallipes</i> (Duftschmid, 1812) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Asaphidion stierlini</i> (Heyden, 1880) | 10 | 0 | 13 | 5 | 0 | 28 |
| <i>Ocys harpaloides</i> (Audinet-Serville, 1821) | 23 | 17 | 83 | 40 | 7 | 170 |
| <i>Ocys quinquestriatus</i> (Gyllenhal, 1810) | 0 | 1 | 11 | 8 | 0 | 20 |
| <i>Bracteon litorale</i> (Olivier, 1790) | 2 | 13 | 1 | 10 | 4 | 30 |
| <i>Bembidion</i> (<i>Neja</i>) <i>nigricorne</i> Gyllenhal, 1827 | 2 | 26 | 3 | 5 | 0 | 36 |
| <i>Bembidion</i> (<i>Metallina</i>) <i>lampros</i> (Herbst, 1784) | 65 | 49 | 247 | 83 | 12 | 456 |
| <i>Bembidion</i> (<i>Metallina</i>) <i>properans</i> (Stephens, 1828) | 12 | 25 | 51 | 14 | 2 | 104 |
| <i>Bembidion</i> (<i>Principidium</i>) <i>punctulatum</i> Drapiez, 1821 | 0 | 15 | 3 | 23 | 7 | 48 |
| <i>Bembidion</i> (<i>Testedium</i>) <i>bipunctatum</i> (Linnaeus, 1761) | 1 | 1 | 1 | 2 | 1 | 6 |
| <i>Bembidion</i> (<i>Eupetedromus</i>) <i>dentellum</i> (Thunberg, 1787) | 32 | 6 | 69 | 32 | 5 | 144 |
| <i>Bembidion</i> (<i>Notaphus</i>) <i>obliquum</i> Sturm, 1825 | 9 | 7 | 52 | 16 | 3 | 87 |
| <i>Bembidion</i> (<i>Notaphus</i>) <i>varium</i> (Olivier, 1795) | 47 | 1 | 23 | 1 | 0 | 72 |
| <i>Bembidion</i> (<i>Plataphus</i>) <i>prasinum</i> (Duftschmid, 1812) | 0 | 3 | 2 | 13 | 9 | 27 |
| <i>Bembidion</i> (<i>Bembidionetolitzkya</i>) <i>atrocaeruleum</i> (Stephens, 1828) | 3 | 18 | 1 | 28 | 20 | 70 |
| <i>Bembidion</i> (<i>Bembidionetolitzkya</i>) <i>geniculatum</i> Heer, 1837/8 | 6 | 7 | 38 | 12 | 6 | 69 |
| <i>Bembidion</i> (<i>Bembidionetolitzkya</i>) <i>tibiale</i> (Duftschmid, 1812) | 0 | 19 | 8 | 31 | 16 | 74 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>bruxellense</i> Wesmael, 1835 | 3 | 13 | 29 | 10 | 1 | 56 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>cruciatum</i> Dejean, 1831 | 13 | 8 | 2 | 0 | 2 | 25 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>decorum</i> (Zenker in Panzer, 1800) | 0 | 17 | 2 | 21 | 10 | 50 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>deletum</i> Audinet-Serville, 1821 | 15 | 6 | 50 | 22 | 4 | 97 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>femoratum</i> Sturm, 1825 | 5 | 5 | 16 | 10 | 2 | 38 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>fluviale</i> Dejean, 1831 | 0 | 0 | 0 | 3 | 0 | 3 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>lunatum</i> (Duftschmid, 1812) | 11 | 4 | 11 | 6 | 0 | 32 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>maritimum</i> (Stephens, 1835) | 7 | 2 | 4 | 0 | 0 | 13 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>monticola</i> Sturm, 1825 | 0 | 4 | 2 | 9 | 3 | 18 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>saxatile</i> Gyllenhal, 1827 | 2 | 9 | 2 | 1 | 1 | 15 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>stephensii</i> Crotch, 1866 | 10 | 8 | 7 | 5 | 0 | 30 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>testaceum</i> (Duftschmid, 1812) | 0 | 1 | 0 | 2 | 2 | 5 |
| <i>Bembidion</i> (<i>Ocydromus</i>) <i>tetracolum</i> Say, 1825 | 31 | 26 | 103 | 61 | 30 | 251 |
| <i>Bembidion</i> (<i>Nepha</i>) <i>illigeri</i> Netolitzky, 1914 | 33 | 9 | 29 | 10 | 2 | 83 |
| <i>Bembidion</i> (<i>Sinechostictus</i>) <i>stomoides</i> Dejean, 1831 | 1 | 2 | 0 | 14 | 4 | 21 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|-----|----|-----|
| Bembidion (Semicampa) gilvipes Sturm, 1825 | 46 | 19 | 19 | 16 | 0 | 100 |
| Bembidion (Semicampa) schuppelii Dejean, 1831 | 0 | 0 | 1 | 1 | 0 | 2 |
| Bembidion (Diplocampa) assimile Gyllenhal, 1810 | 34 | 2 | 60 | 1 | 1 | 98 |
| Bembidion (Diplocampa) clarkii (Dawson, 1849) | 18 | 11 | 37 | 20 | 0 | 86 |
| Bembidion (Diplocampa) fumigatum (Duftschmid, 1812) | 24 | 1 | 38 | 2 | 0 | 65 |
| Bembidion (Emphanes) minimum (Fabricius, 1792) | 43 | 0 | 0 | 1 | 0 | 44 |
| Bembidion (Emphanes) normannum Dejean, 1831 | 6 | 0 | 0 | 0 | 0 | 6 |
| Bembidion (Bembidion) humerale Sturm, 1825 | 0 | 0 | 26 | 0 | 0 | 26 |
| Bembidion (Bembidion) quadrimaculatum (Linnaeus, 1761) | 29 | 9 | 55 | 32 | 2 | 127 |
| Bembidion (Bembidion) quadripustulatum Audinet-Serville, 1821 | 0 | 0 | 1 | 0 | 0 | 1 |
| Bembidion (Trepanedoris) doris (Panzer, 1796) | 17 | 3 | 16 | 18 | 0 | 54 |
| Bembidion (Trepanes) articulatum (Panzer, 1795) | 7 | 0 | 8 | 1 | 0 | 16 |
| Bembidion (Phyla) obtusum Audinet-Serville, 1821 | 47 | 15 | 64 | 30 | 1 | 157 |
| Bembidion (Philochthus) aeneum Germar, 1824 | 52 | 24 | 94 | 45 | 5 | 220 |
| Bembidion (Philochthus) biguttatum (Fabricius, 1779) | 64 | 24 | 83 | 42 | 2 | 215 |
| Bembidion (Philochthus) guttula (Fabricius, 1792) | 60 | 32 | 194 | 75 | 8 | 369 |
| Bembidion (Philochthus) iricolor Bedel, 1879 | 20 | 0 | 4 | 0 | 0 | 24 |
| Bembidion (Philochthus) lunulatum (Geoffroy in Fourcroy, 1785) | 13 | 2 | 54 | 25 | 0 | 94 |
| Bembidion (Philochthus) mannerheimii Sahlberg, C.R., 1827 | 8 | 20 | 20 | 19 | 5 | 72 |
| Pogonus chalceus (Marsham, 1802) | 10 | 0 | 0 | 0 | 0 | 10 |
| Pogonus littoralis (Duftschmid, 1812) | 1 | 0 | 0 | 0 | 0 | 1 |
| Patrobus assimilis Chaudoir, 1844 | 1 | 14 | 2 | 16 | 12 | 45 |
| Patrobus atrorufus (Ström, 1768) | 6 | 11 | 29 | 41 | 10 | 97 |
| Patrobus septentrionis Dejean, 1828 | 2 | 0 | 2 | 0 | 0 | 4 |
| Stomis pumicatus (Panzer, 1795) | 16 | 12 | 32 | 18 | 1 | 79 |
| Poecilus cupreus (Linnaeus, 1758) | 11 | 11 | 47 | 5 | 1 | 75 |
| Poecilus lepidus (Leske, 1785) | 3 | 18 | 0 | 0 | 0 | 21 |
| Poecilus versicolor (Sturm, 1824) | 14 | 20 | 46 | 10 | 1 | 91 |
| Pterostichus (Pterostichus) cristatus (Dufour, 1820) | 0 | 0 | 2 | 6 | 7 | 15 |
| Pterostichus (Steropus) aethiops (Panzer, 1796) | 0 | 3 | 0 | 4 | 3 | 10 |
| Pterostichus (Steropus) madidus (Fabricius, 1775) | 29 | 48 | 378 | 124 | 39 | 618 |
| Pterostichus (Pediis) longicollis (Duftschmid, 1812) | 4 | 0 | 0 | 2 | 0 | 6 |
| Pterostichus (Adelosia) macer (Marsham, 1802) | 12 | 9 | 6 | 4 | 1 | 32 |
| Pterostichus (Platysma) niger (Schaller, 1783) | 40 | 21 | 216 | 55 | 16 | 348 |
| Pterostichus (Bothriopterus) adstrictus Eschscholtz, 1823 | 3 | 15 | 13 | 18 | 11 | 60 |
| Pterostichus (Bothriopterus) oblongopunctatus (Fabricius, 1787) | 9 | 23 | 21 | 4 | 0 | 57 |
| Pterostichus (Bothriopterus) quadrifoveolatus Letzner, 1852 | 7 | 2 | 36 | 11 | 0 | 56 |
| Pterostichus (Omaseus) melanarius (Illiger, 1798) | 37 | 15 | 97 | 62 | 10 | 221 |
| Pterostichus (Pseudomaseus) anthracinus (Panzer, 1795) | 15 | 0 | 5 | 0 | 0 | 20 |
| Pterostichus (Pseudomaseus) gracilis (Dejean, 1828) | 6 | 0 | 2 | 6 | 0 | 14 |
| Pterostichus (Pseudomaseus) minor (Gyllenhal, 1827) | 24 | 17 | 91 | 27 | 0 | 159 |
| Pterostichus (Pseudomaseus) nigrita (Paykull, 1790) | 25 | 34 | 152 | 50 | 5 | 266 |
| Pterostichus (Pseudomaseus) rhaeticus Heer, 1837/8 | 2 | 7 | 23 | 12 | 3 | 47 |
| Pterostichus (Lagarus) vernalis (Panzer, 1795) | 13 | 9 | 32 | 25 | 0 | 79 |
| Pterostichus (Argutor) diligens (Sturm, 1824) | 33 | 44 | 242 | 84 | 12 | 415 |
| Pterostichus (Argutor) strenuus (Panzer, 1796) | 92 | 40 | 329 | 119 | 19 | 599 |
| Abax parallelepipedus (Piller & Mitterpacher, 1783) | 10 | 11 | 130 | 53 | 9 | 213 |
| Platyderus depressus (Audinet-Serville, 1821) | 8 | 1 | 3 | 0 | 0 | 12 |
| Synuchus vivalis (Illiger, 1798) | 6 | 6 | 24 | 11 | 4 | 51 |
| Calathus (Amphigynus) rotundicollis Dejean, 1828 | 6 | 2 | 37 | 11 | 0 | 56 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| Calathus (Calathus) ambiguus (Paykull, 1790) | 4 | 3 | 6 | 0 | 0 | 13 |
| Calathus (Calathus) erratus (Sahlberg, C.R., 1827) | 4 | 9 | 7 | 2 | 0 | 22 |
| Calathus (Calathus) fuscipes (Goeze, 1777) | 20 | 9 | 62 | 22 | 8 | 121 |
| Calathus (Calathus) melanocephalus (Linnaeus, 1758) | 18 | 1 | 61 | 10 | 7 | 97 |
| Calathus (Calathus) micropterus (Duftschmid, 1812) | 0 | 13 | 5 | 14 | 14 | 46 |
| Calathus (Calathus) mollis (Marsham, 1802) | 24 | 6 | 1 | 0 | 0 | 31 |
| Sphodrus leucophthalmus (Linnaeus, 1758) | 0 | 2 | 3 | 0 | 0 | 5 |
| Laemostenus (Laemostenus) complanatus (Dejean, 1828) | 1 | 1 | 1 | 1 | 0 | 4 |
| Laemostenus (Pristonychus) terricola (Herbst, 1784) | 7 | 8 | 8 | 4 | 0 | 27 |
| Olisthopus rotundatus (Paykull, 1790) | 4 | 20 | 52 | 14 | 5 | 95 |
| Oxypselaphus obscurus (Herbst, 1784) | 23 | 7 | 152 | 47 | 4 | 233 |
| Paranchus albipes (Fabricius, 1796) | 33 | 40 | 180 | 74 | 29 | 356 |
| Batenus livens (Gyllenhal, 1810) | 6 | 0 | 14 | 0 | 0 | 20 |
| Platynus assimilis (Paykull, 1790) | 6 | 17 | 148 | 41 | 8 | 220 |
| Anchomenus dorsalis (Pontoppidan, 1763) | 42 | 10 | 81 | 27 | 1 | 161 |
| Sericoda quadripunctata (De Geer, 1774) | 0 | 1 | 0 | 0 | 0 | 1 |
| Agonum (Europhilus) fuliginosum (Panzer, 1809) | 66 | 21 | 226 | 73 | 20 | 406 |
| Agonum (Europhilus) gracile Sturm, 1824 | 14 | 9 | 79 | 22 | 3 | 127 |
| Agonum (Europhilus) micans Nicolai, 1822 | 7 | 2 | 21 | 11 | 1 | 42 |
| Agonum (Europhilus) piceum (Linnaeus, 1758) | 12 | 5 | 20 | 5 | 0 | 42 |
| Agonum (Europhilus) thoreyi Dejean, 1828 | 37 | 5 | 87 | 21 | 2 | 152 |
| Agonum (Agonum) emarginatum (Gyllenhal, 1827) | 25 | 7 | 24 | 12 | 3 | 71 |
| Agonum (Agonum) ericeti (Panzer, 1809) | 0 | 7 | 11 | 6 | 2 | 26 |
| Agonum (Agonum) gracilipes (Duftschmid, 1812) | 2 | 0 | 0 | 0 | 0 | 2 |
| Agonum (Agonum) marginatum (Linnaeus, 1758) | 27 | 8 | 89 | 16 | 4 | 144 |
| Agonum (Agonum) muelleri (Herbst, 1784) | 8 | 14 | 46 | 34 | 5 | 107 |
| Agonum (Agonum) nigrum Dejean, 1828 | 0 | 0 | 1 | 0 | 0 | 1 |
| Agonum (Agonum) versutum Sturm, 1824 | 10 | 0 | 2 | 0 | 0 | 12 |
| Agonum (Agonum) viduum (Panzer, 1796) | 6 | 0 | 7 | 7 | 0 | 20 |
| Amara (Zezea) plebeja (Gyllenhal, 1810) | 31 | 26 | 165 | 49 | 5 | 276 |
| Amara (Amara) aenea (De Geer, 1774) | 21 | 34 | 178 | 23 | 7 | 263 |
| Amara (Amara) anthobia Villa & Villa, 1833 | 0 | 0 | 2 | 0 | 0 | 2 |
| Amara (Amara) communis (Panzer, 1797) | 4 | 8 | 41 | 17 | 2 | 72 |
| Amara (Amara) convexior Stephens, 1828 | 0 | 3 | 4 | 1 | 0 | 8 |
| Amara (Amara) curta Dejean, 1828 | 0 | 3 | 1 | 0 | 0 | 4 |
| Amara (Amara) eurynota (Panzer, 1796) | 6 | 5 | 23 | 3 | 1 | 38 |
| Amara (Amara) famelica Zimmermann, 1832 | 1 | 3 | 0 | 0 | 0 | 4 |
| Amara (Amara) familiaris (Duftschmid, 1812) | 38 | 24 | 136 | 40 | 6 | 244 |
| Amara (Amara) lucida (Duftschmid, 1812) | 7 | 1 | 2 | 1 | 0 | 11 |
| Amara (Amara) lunicollis Schiödte, 1837 | 4 | 12 | 75 | 14 | 1 | 106 |
| Amara (Amara) ovata (Fabricius, 1792) | 9 | 4 | 17 | 12 | 2 | 44 |
| Amara (Amara) similata (Gyllenhal, 1810) | 22 | 11 | 75 | 21 | 3 | 132 |
| Amara (Amara) spreta Dejean, 1831 | 0 | 2 | 0 | 0 | 0 | 2 |
| Amara (Amara) tibialis (Paykull, 1798) | 15 | 2 | 25 | 1 | 0 | 43 |
| Amara (Celia) bifrons (Gyllenhal, 1810) | 8 | 0 | 12 | 8 | 1 | 29 |
| Amara (Celia) fusca Dejean, 1828 | 0 | 0 | 1 | 0 | 0 | 1 |
| Amara (Celia) praetermissa (Sahlberg, C.R., 1827) | 2 | 2 | 9 | 0 | 1 | 14 |
| Amara (Bradytus) apricaria (Paykull, 1790) | 8 | 5 | 45 | 9 | 9 | 76 |
| Amara (Bradytus) consularis (Duftschmid, 1812) | 4 | 2 | 2 | 2 | 0 | 10 |
| Amara (Bradytus) fulva (Müller, O.F., 1776) | 11 | 0 | 9 | 0 | 2 | 22 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|---|-----|
| <i>Amara (Percosia) equestris</i> (Duftschmid, 1812) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Curtonotus aulicus</i> (Panzer, 1796) | 24 | 17 | 74 | 30 | 3 | 148 |
| <i>Curtonotus convexiusculus</i> (Marshall, 1802) | 26 | 3 | 14 | 7 | 0 | 50 |
| <i>Harpalus (Pseudoophonus) calceatus</i> (Duftschmid, 1812) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Harpalus (Pseudoophonus) rufipes</i> (De Geer, 1774) | 38 | 13 | 83 | 28 | 4 | 166 |
| <i>Harpalus (Harpalus) affinis</i> (Schrank, 1781) | 30 | 12 | 76 | 21 | 4 | 143 |
| <i>Harpalus (Harpalus) anxius</i> (Duftschmid, 1812) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Harpalus (Harpalus) attenuatus</i> Stephens, 1828 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Harpalus (Harpalus) froelichii</i> Sturm, 1818 | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Harpalus (Harpalus) latus</i> (Linnaeus, 1758) | 9 | 11 | 10 | 5 | 2 | 37 |
| <i>Harpalus (Harpalus) neglectus</i> Audinet-Serville, 1821 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Harpalus (Harpalus) rubripes</i> (Duftschmid, 1812) | 3 | 2 | 12 | 4 | 0 | 21 |
| <i>Harpalus (Harpalus) rufipalpis</i> Sturm, 1818 | 0 | 0 | 9 | 1 | 0 | 10 |
| <i>Harpalus (Harpalus) servus</i> (Duftschmid, 1812) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Harpalus (Harpalus) smaragdinus</i> (Duftschmid, 1812) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Harpalus (Harpalus) tardus</i> (Panzer, 1796) | 26 | 1 | 6 | 1 | 0 | 34 |
| <i>Ophonus (Ophonus) ardosiacus</i> Lutshnik, 1922 | 5 | 1 | 0 | 0 | 0 | 6 |
| <i>Ophonus (Metophonus) puncticeps</i> Stephens, 1828 | 3 | 0 | 2 | 0 | 0 | 5 |
| <i>Ophonus (Metophonus) puncticollis</i> (Paykull, 1798) | 2 | 0 | 2 | 0 | 0 | 4 |
| <i>Ophonus (Metophonus) rufibarbis</i> (Fabricius, 1792) | 32 | 9 | 41 | 8 | 5 | 95 |
| <i>Ophonus (Metophonus) rupicola</i> (Sturm, 1818) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Ophonus (Metophonus) schaubergerianus</i> (Puel, 1937) | 3 | 3 | 1 | 1 | 0 | 8 |
| <i>Anisodactylus binotatus</i> (Fabricius, 1787) | 2 | 2 | 53 | 4 | 0 | 61 |
| <i>Dicheirotichus gustavii</i> Crotch, 1871 | 15 | 0 | 0 | 0 | 0 | 15 |
| <i>Dicheirotichus obsoletus</i> (Dejean, 1829) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Trichocellus cognatus</i> (Gyllenhal, 1827) | 6 | 22 | 29 | 15 | 6 | 78 |
| <i>Trichocellus placidus</i> (Gyllenhal, 1827) | 37 | 23 | 130 | 60 | 4 | 254 |
| <i>Bradycellus caucasicus</i> (Chaudoir, 1846) | 2 | 6 | 12 | 10 | 7 | 37 |
| <i>Bradycellus distinctus</i> (Dejean, 1829) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Bradycellus harpalinus</i> (Audinet-Serville, 1821) | 13 | 26 | 133 | 36 | 4 | 212 |
| <i>Bradycellus ruficollis</i> (Stephens, 1828) | 4 | 34 | 52 | 14 | 8 | 112 |
| <i>Bradycellus sharpi</i> Joy, 1912 | 2 | 5 | 12 | 5 | 0 | 24 |
| <i>Bradycellus verbasci</i> (Duftschmid, 1812) | 8 | 4 | 56 | 11 | 4 | 83 |
| <i>Stenolophus mixtus</i> (Herbst, 1784) | 25 | 6 | 96 | 25 | 1 | 153 |
| <i>Acupalpus brunnipes</i> (Sturm, 1825) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Acupalpus dubius</i> Schilsky, 1888 | 19 | 23 | 173 | 32 | 1 | 248 |
| <i>Acupalpus exiguus</i> Dejean, 1829 | 1 | 0 | 5 | 0 | 0 | 6 |
| <i>Acupalpus meridianus</i> (Linnaeus, 1761) | 5 | 0 | 12 | 2 | 0 | 19 |
| <i>Acupalpus parvulus</i> (Sturm, 1825) | 21 | 3 | 43 | 8 | 0 | 75 |
| <i>Anthracus consputus</i> (Duftschmid, 1812) | 7 | 1 | 4 | 4 | 0 | 16 |
| <i>Chlaenius nigricornis</i> (Fabricius, 1787) | 14 | 2 | 1 | 3 | 1 | 21 |
| <i>Chlaenius tristis</i> (Schaller, 1783) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Chlaenius vestitus</i> (Paykull, 1790) | 19 | 1 | 2 | 0 | 0 | 22 |
| <i>Oodes helopioides</i> (Fabricius, 1792) | 1 | 1 | 0 | 1 | 0 | 3 |
| <i>Licinus depressus</i> (Paykull, 1790) | 9 | 1 | 1 | 0 | 0 | 11 |
| <i>Licinus punctatulus</i> (Fabricius, 1792) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Badister (Badister) bullatus</i> (Schrank, 1798) | 43 | 10 | 85 | 25 | 1 | 164 |
| <i>Badister (Badister) unipustulatus</i> Bonelli, 1813 | 4 | 0 | 11 | 0 | 0 | 15 |
| <i>Badister (Trimorphus) sodalis</i> (Duftschmid, 1812) | 1 | 3 | 17 | 4 | 0 | 25 |
| <i>Panagaeus bipustulatus</i> (Fabricius, 1775) | 13 | 0 | 0 | 0 | 0 | 13 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|-----|----|-----|----|---|-----|
| Panagaeus cruxmajor (Linnaeus, 1758) | 1 | 0 | 2 | 0 | 0 | 3 |
| Perigona nigriceps (Dejean, 1831) | 1 | 0 | 0 | 2 | 0 | 3 |
| Lebia (Lamprias) chlorocephala (Hoffmann, J., 1803) | 3 | 7 | 2 | 0 | 0 | 12 |
| Lebia (Lamprias) cyanocephala (Linnaeus, 1758) | 0 | 0 | 0 | 1 | 0 | 1 |
| Demetrias (Demetrias) atricapillus (Linnaeus, 1758) | 77 | 19 | 113 | 23 | 1 | 233 |
| Demetrias (Demetrias) monostigma Samouelle, 1819 | 7 | 1 | 0 | 0 | 0 | 8 |
| Cymindis (Tarulus) vaporariorum (Linnaeus, 1758) | 0 | 9 | 3 | 0 | 0 | 12 |
| Paradromius linearis (Olivier, 1795) | 104 | 35 | 207 | 50 | 7 | 403 |
| Paradromius longiceps Dejean, 1826 | 8 | 0 | 7 | 0 | 0 | 15 |
| Dromius agilis (Fabricius, 1787) | 2 | 3 | 11 | 10 | 0 | 26 |
| Dromius angustus Brullé, 1834 | 1 | 7 | 2 | 2 | 0 | 12 |
| Dromius meridionalis Dejean, 1825 | 11 | 4 | 7 | 7 | 0 | 29 |
| Dromius quadrimaculatus (Linnaeus, 1758) | 9 | 20 | 47 | 23 | 1 | 100 |
| Calodromius spilotus (Illiger, 1798) | 17 | 13 | 32 | 9 | 1 | 72 |
| Philorhizus melanocephalus (Dejean, 1825) | 65 | 11 | 125 | 26 | 0 | 227 |
| Philorhizus notatus (Stephens, 1827) | 12 | 4 | 0 | 1 | 0 | 17 |
| Philorhizus quadrisignatus (Dejean, 1825) | 0 | 0 | 1 | 0 | 0 | 1 |
| Philorhizus sigma (Rossi, 1790) | 0 | 1 | 6 | 7 | 0 | 14 |
| Microlestes maurus (Sturm, 1827) | 18 | 0 | 1 | 0 | 0 | 19 |
| Lionychus quadrillum (Duftschmid, 1812) | 0 | 1 | 0 | 0 | 0 | 1 |
| Syntomus foveatus (Geoffroy in Fourcroy, 1785) | 21 | 8 | 15 | 1 | 0 | 45 |
| Syntomus obscuroguttatus (Duftschmid, 1812) | 0 | 0 | 1 | 0 | 0 | 1 |
| Syntomus truncatellus (Linnaeus, 1761) | 8 | 1 | 5 | 1 | 0 | 15 |
| Odacantha melanura (Linnaeus, 1767) | 0 | 0 | 1 | 0 | 0 | 1 |

Helophoridae [20/19]

| | | | | | | |
|--|----|----|-----|----|----|-----|
| Helophorus (Empleurus) nubilus Fabricius, 1777 | 4 | 3 | 7 | 9 | 0 | 23 |
| Helophorus (Empleurus) porculus Bedel, 1881 | 0 | 4 | 0 | 2 | 0 | 6 |
| Helophorus (Empleurus) rufipes (Bosc d'Antic, 1791) | 1 | 0 | 0 | 0 | 0 | 1 |
| Helophorus (Cyphelophorus) tuberculatus Gyllenhal, 1808 | 0 | 10 | 2 | 2 | 0 | 14 |
| Helophorus (Trichohelophorus) alternans Gené 1836 | 0 | 0 | 0 | 1 | 0 | 1 |
| Helophorus (Megahelophorus) aequalis Thomson, C.G., 1868 | 25 | 8 | 74 | 24 | 4 | 135 |
| Helophorus (Megahelophorus) grandis Illiger, 1798 | 22 | 24 | 98 | 18 | 7 | 169 |
| Helophorus (Atracthelophorus) arvernicus Mulsant, 1846 | 1 | 7 | 3 | 2 | 7 | 20 |
| Helophorus (Atracthelophorus) brevipalpis Bedel, 1881 | 44 | 44 | 187 | 54 | 26 | 355 |
| Helophorus (Helophorus) dorsalis (Marsham, 1802) | 4 | 1 | 1 | 0 | 0 | 6 |
| Helophorus (Helophorus) flavipes Fabricius, 1792 | 7 | 19 | 37 | 19 | 7 | 89 |
| Helophorus (Helophorus) fulgidicollis Motschulsky, 1860 | 6 | 0 | 0 | 0 | 0 | 6 |
| Helophorus (Helophorus) granularis (Linnaeus, 1760) | 3 | 1 | 4 | 3 | 0 | 11 |
| Helophorus (Helophorus) griseus Herbst, 1793 | 9 | 1 | 7 | 1 | 0 | 18 |
| Helophorus (Helophorus) longitarsis Wollaston, 1864 | 2 | 0 | 0 | 0 | 0 | 2 |
| Helophorus (Helophorus) minutus Fabricius, 1775 | 41 | 8 | 82 | 31 | 5 | 167 |
| Helophorus (Helophorus) nanus Sturm, 1836 | 11 | 0 | 2 | 3 | 0 | 16 |
| Helophorus (Helophorus) obscurus Mulsant, 1844 | 11 | 5 | 24 | 7 | 1 | 48 |
| Helophorus (Helophorus) strigifrons Thomson, C.G., 1868 | 1 | 0 | 1 | 3 | 0 | 5 |

Georissidae [1/1]

| | | | | | | |
|------------------------------------|----|---|---|---|---|----|
| Georissus crenulatus (Rossi, 1794) | 12 | 6 | 0 | 1 | 0 | 19 |
|------------------------------------|----|---|---|---|---|----|

Hydrochidae [7/3]

| | | | | | | |
|--------------------------------------|---|---|---|---|---|----|
| Hydrochus angustatus Germar, 1824 | 1 | 0 | 0 | 1 | 0 | 2 |
| Hydrochus brevis (Herbst, 1793) | 1 | 0 | 1 | 0 | 0 | 2 |
| Hydrochus elongatus (Schaller, 1783) | 1 | 1 | 7 | 3 | 0 | 12 |

Spercheidae [1/1]

| | | | | | | |
|--|---|---|---|---|---|---|
| Spercheus emarginatus (Schaller, 1783) | 0 | 1 | 0 | 2 | 0 | 3 |
|--|---|---|---|---|---|---|

Hydrophilidae [70/62]

| | | | | | | |
|---|----|----|-----|----|----|-----|
| Anacaena bipustulata (Marsham, 1802) | 1 | 0 | 3 | 0 | 0 | 4 |
| Anacaena globulus (Paykull, 1798) | 64 | 41 | 289 | 66 | 27 | 487 |
| Anacaena limbata (Fabricius, 1792) | 53 | 6 | 89 | 34 | 1 | 183 |
| Anacaena lutescens (Stephens, 1829) | 12 | 4 | 53 | 19 | 2 | 90 |
| Berosus (Berosus) affinis Brullé, 1835 | 3 | 1 | 3 | 1 | 0 | 8 |
| Berosus (Berosus) luridus (Linnaeus, 1761) | 3 | 0 | 5 | 0 | 0 | 8 |
| Berosus (Berosus) signaticollis (Charpentier, 1825) | 1 | 0 | 0 | 2 | 0 | 3 |
| Berosus (Enoplurus) fulvus Kuwert, 1888 | 0 | 1 | 0 | 0 | 0 | 1 |
| Chaetarthria seminulum (Herbst, 1797) | 10 | 11 | 13 | 8 | 0 | 42 |
| Chaetarthria simillima Vorst & Cuppen, 2003 | 3 | 0 | 0 | 0 | 0 | 3 |
| Cymbiodyta marginellus (Fabricius, 1792) | 24 | 3 | 19 | 15 | 0 | 61 |
| Enochrus Thomson, C.G., 1859 | 0 | 0 | 1 | 0 | 0 | 1 |
| Enochrus affinis (Thunberg, 1794) | 7 | 5 | 5 | 4 | 0 | 21 |
| Enochrus bicolor (Fabricius, 1792) | 8 | 1 | 3 | 0 | 0 | 12 |
| Enochrus coarctatus (Gredler, 1863) | 2 | 3 | 5 | 6 | 0 | 16 |
| Enochrus fuscipennis (Thomson, C.G., 1884) | 2 | 0 | 1 | 0 | 0 | 3 |
| Enochrus halophilus (Bedel, 1878) | 6 | 0 | 0 | 0 | 0 | 6 |
| Enochrus melanocephalus (Olivier, 1792) | 3 | 3 | 11 | 5 | 0 | 22 |
| Enochrus ochropterus (Marsham, 1802) | 3 | 1 | 6 | 3 | 0 | 13 |
| Enochrus quadripunctatus (Herbst, 1797) | 1 | 1 | 0 | 1 | 0 | 3 |
| Enochrus testaceus (Fabricius, 1801) | 13 | 3 | 16 | 10 | 0 | 42 |
| Helochares lividus (Forster, 1771) | 17 | 0 | 12 | 7 | 0 | 36 |
| Helochares obscurus (Müller, O.F., 1776) | 3 | 1 | 5 | 0 | 0 | 9 |
| Helochares punctatus Sharp, 1869 | 6 | 4 | 25 | 5 | 0 | 40 |
| Hydrobius fuscipes (Linnaeus, 1758) | 68 | 27 | 146 | 32 | 14 | 287 |
| Hydrochara caraboides (Linnaeus, 1758) | 0 | 0 | 2 | 2 | 0 | 4 |
| Hydrophilus piceus (Linnaeus, 1758) | 0 | 0 | 3 | 0 | 0 | 3 |
| Laccobius atratus Rottenberg, 1874 | 1 | 2 | 0 | 0 | 1 | 4 |
| Laccobius bipunctatus (Fabricius, 1775) | 11 | 11 | 51 | 17 | 3 | 93 |
| Laccobius colon (Stephens, 1829) | 10 | 2 | 29 | 15 | 0 | 56 |
| Laccobius minutus (Linnaeus, 1758) | 14 | 6 | 26 | 7 | 4 | 57 |
| Laccobius sinuatus Motschulsky, 1849 | 9 | 1 | 11 | 3 | 0 | 24 |
| Laccobius striatulus (Fabricius, 1801) | 0 | 5 | 5 | 6 | 3 | 19 |
| Laccobius ytenensis Sharp, 1910 | 0 | 0 | 1 | 0 | 0 | 1 |
| Coelostoma orbiculare (Fabricius, 1775) | 5 | 12 | 11 | 9 | 1 | 38 |
| Cercyon (Cercyon) bifenestratus Küster, 1851 | 0 | 0 | 3 | 0 | 0 | 3 |
| Cercyon (Cercyon) convexiusculus Stephens, 1829 | 25 | 2 | 44 | 21 | 3 | 95 |
| Cercyon (Cercyon) depressus Stephens, 1829 | 3 | 0 | 1 | 0 | 0 | 4 |
| Cercyon (Cercyon) granarius Erichson, 1837 | 2 | 1 | 0 | 0 | 0 | 3 |
| Cercyon (Cercyon) haemorrhoidalis (Fabricius, 1775) | 35 | 13 | 92 | 22 | 13 | 175 |
| Cercyon (Cercyon) impressus (Sturm, 1807) | 14 | 13 | 85 | 20 | 13 | 145 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| Cercyon (Cercyon) lateralis (Marsham, 1802) | 27 | 25 | 93 | 18 | 13 | 176 |
| Cercyon (Cercyon) littoralis (Gyllenhal, 1808) | 16 | 6 | 0 | 0 | 0 | 22 |
| Cercyon (Cercyon) marinus Thomson, C.G., 1853 | 18 | 1 | 17 | 2 | 1 | 39 |
| Cercyon (Cercyon) melanocephalus (Linnaeus, 1758) | 35 | 18 | 152 | 34 | 14 | 253 |
| Cercyon (Cercyon) nigriceps (Marsham, 1802) | 3 | 4 | 4 | 2 | 0 | 13 |
| Cercyon (Cercyon) obsoletus (Gyllenhal, 1808) | 4 | 2 | 16 | 2 | 0 | 24 |
| Cercyon (Cercyon) pygmaeus (Illiger, 1801) | 11 | 8 | 22 | 8 | 3 | 52 |
| Cercyon (Cercyon) quisquilius (Linnaeus, 1761) | 7 | 5 | 23 | 3 | 12 | 50 |
| Cercyon (Cercyon) terminatus (Marsham, 1802) | 11 | 3 | 10 | 2 | 1 | 27 |
| Cercyon (Cercyon) tristis (Illiger, 1801) | 15 | 7 | 8 | 17 | 1 | 48 |
| Cercyon (Cercyon) unipunctatus (Linnaeus, 1758) | 15 | 10 | 32 | 6 | 6 | 69 |
| Cercyon (Paracercyon) analis (Paykull, 1798) | 26 | 5 | 65 | 8 | 0 | 104 |
| Cercyon (Dicyrtocercyon) ustulatus (Preyssl, 1790) | 11 | 5 | 26 | 6 | 2 | 50 |
| Megasternum concinnum (Marsham, 1802) | 86 | 31 | 239 | 76 | 32 | 464 |
| Cryptopleurum crenatum (Kugelann, 1794) | 0 | 0 | 0 | 3 | 0 | 3 |
| Cryptopleurum minutum (Fabricius, 1775) | 29 | 14 | 100 | 18 | 8 | 169 |
| Cryptopleurum subtile Sharp, 1884 | 3 | 1 | 1 | 1 | 0 | 6 |
| Sphaeridium bipustulatum Fabricius, 1781 | 8 | 5 | 14 | 10 | 1 | 38 |
| Sphaeridium lunatum Fabricius, 1792 | 15 | 12 | 34 | 4 | 1 | 66 |
| Sphaeridium marginatum Fabricius, 1787 | 1 | 0 | 2 | 0 | 0 | 3 |
| Sphaeridium scarabaeoides (Linnaeus, 1758) | 9 | 3 | 38 | 8 | 6 | 64 |

Sphaeritidae [1/1]

| | | | | | | |
|--|---|---|---|---|---|---|
| Sphaerites glabratus (Fabricius, 1792) | 0 | 4 | 0 | 1 | 0 | 5 |
|--|---|---|---|---|---|---|

Histeridae [51/32]

| | | | | | | |
|---|----|----|----|----|---|----|
| Abraeus granulum Erichson, 1839 | 0 | 0 | 2 | 0 | 0 | 2 |
| Abraeus perpusillus (Marsham, 1802) | 3 | 7 | 5 | 1 | 1 | 17 |
| Plegaderus vulneratus (Panzer, 1797) | 0 | 0 | 2 | 1 | 0 | 3 |
| Acritus nigricornis (Hoffmann, J., 1803) | 4 | 3 | 6 | 1 | 0 | 14 |
| Aeletes atomarius (Aubé, 1842) | 0 | 1 | 1 | 0 | 0 | 2 |
| Saprinus aeneus (Fabricius, 1775) | 3 | 2 | 5 | 0 | 0 | 10 |
| Saprinus planiusculus Motschulsky, 1849 | 1 | 0 | 0 | 0 | 0 | 1 |
| Saprinus semistriatus (Scriba, 1790) | 9 | 9 | 15 | 5 | 1 | 39 |
| Saprinus virescens (Paykull, 1798) | 1 | 0 | 0 | 2 | 0 | 3 |
| Hypocaccus (Hypocaccus) rugiceps (Duftschmid, 1805) | 1 | 0 | 0 | 0 | 0 | 1 |
| Hypocaccus (Hypocaccus) rugifrons (Paykull, 1798) | 0 | 1 | 1 | 0 | 0 | 2 |
| Hypocaccus (Baeckmanniolus) dimidiatus (Illiger, 1807) | 1 | 2 | 0 | 0 | 0 | 3 |
| Gnathoncus nannetensis (Marseul, 1862) | 3 | 5 | 1 | 2 | 0 | 11 |
| Gnathoncus rotundatus (Kugelann, 1792) | 1 | 0 | 1 | 2 | 0 | 4 |
| Myrmetes paykulli Kanaar, 1979 | 0 | 5 | 0 | 0 | 0 | 5 |
| Dendrophilus punctatus (Herbst, 1792) | 0 | 1 | 0 | 0 | 0 | 1 |
| Dendrophilus pygmaeus (Linnaeus, 1758) | 0 | 0 | 2 | 0 | 0 | 2 |
| Kissister minimus (Aubé, 1850) | 0 | 0 | 1 | 0 | 0 | 1 |
| Carcinops pumilio (Erichson, 1834) | 8 | 2 | 14 | 3 | 1 | 28 |
| Paromalus flavicornis (Herbst, 1792) | 0 | 4 | 9 | 3 | 0 | 16 |
| Paromalus parallelepipedus (Herbst, 1792) | 0 | 0 | 1 | 0 | 0 | 1 |
| Onthophilus striatus (Forster, 1771) | 18 | 10 | 60 | 10 | 1 | 99 |
| Margarinotus (Ptomister) brunneus (Fabricius, 1775) | 14 | 6 | 22 | 6 | 1 | 49 |
| Margarinotus (Ptomister) merdarius (Hoffmann, J., 1803) | 2 | 0 | 14 | 0 | 0 | 16 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|---|---|----|----|---|----|
| Margarinotus (Ptomister) striola (Sahlberg, C.R., 1819) | 2 | 7 | 15 | 10 | 5 | 39 |
| Margarinotus (Paralister) neglectus (Germar, 1813) | 0 | 1 | 0 | 1 | 0 | 2 |
| Margarinotus (Paralister) purpurascens (Herbst, 1792) | 7 | 1 | 2 | 0 | 0 | 10 |
| Margarinotus (Paralister) ventralis (Marseul, 1854) | 8 | 6 | 24 | 5 | 2 | 45 |
| Margarinotus (Promethister) marginatus (Erichson, 1834) | 4 | 2 | 1 | 0 | 0 | 7 |
| Hister unicolor Linnaeus, 1758 | 5 | 4 | 7 | 0 | 2 | 18 |
| Atholus bimaculatus (Linnaeus, 1758) | 2 | 1 | 10 | 2 | 0 | 15 |
| Atholus duodecimstriatus (Schrank, 1781) | 3 | 4 | 16 | 3 | 1 | 27 |

Hydraenidae [30/20]

| | | | | | | |
|--|----|----|----|----|---|-----|
| Hydraena britteni Joy, 1907 | 3 | 1 | 16 | 7 | 1 | 28 |
| Hydraena flavipes Sturm, 1836 | 0 | 2 | 2 | 0 | 0 | 4 |
| Hydraena gracilis Germar, 1824 | 0 | 8 | 4 | 9 | 8 | 29 |
| Hydraena nigrita Germar, 1824 | 1 | 5 | 7 | 9 | 4 | 26 |
| Hydraena palustris Erichson, 1837 | 1 | 1 | 2 | 2 | 1 | 7 |
| Hydraena pulchella Germar, 1824 | 0 | 1 | 2 | 0 | 0 | 3 |
| Hydraena pygmaea Waterhouse, G.R., 1833 | 0 | 3 | 2 | 2 | 0 | 7 |
| Hydraena riparia Kugelann, 1794 | 8 | 5 | 11 | 10 | 1 | 35 |
| Hydraena rufipes Curtis, 1830 | 0 | 3 | 0 | 3 | 2 | 8 |
| Hydraena testacea Curtis, 1830 | 4 | 3 | 10 | 1 | 0 | 18 |
| Limnebius aluta Bedel, 1881 | 1 | 0 | 1 | 3 | 0 | 5 |
| Limnebius nitidus (Marsham, 1802) | 0 | 2 | 7 | 0 | 0 | 9 |
| Limnebius papposus Mulsant, 1844 | 1 | 0 | 5 | 1 | 0 | 7 |
| Limnebius truncatellus (Thunberg, 1794) | 4 | 14 | 34 | 6 | 5 | 63 |
| Enicocerus exsculptus (Germar, 1824) | 3 | 5 | 2 | 7 | 8 | 25 |
| Ochthebius (Asiobates) auriculatus Rey, 1885 | 3 | 0 | 0 | 0 | 0 | 3 |
| Ochthebius (Asiobates) bicolon Germar, 1824 | 3 | 4 | 7 | 8 | 1 | 23 |
| Ochthebius (Asiobates) dilatatus Stephens, 1829 | 20 | 3 | 8 | 2 | 0 | 33 |
| Ochthebius (Homalochthebius) minimus (Fabricius, 1792) | 49 | 15 | 37 | 24 | 5 | 130 |
| Ochthebius (Ochthebius) marinus (Paykull, 1798) | 10 | 1 | 0 | 0 | 0 | 11 |

Ptiliidae [75/57]

| | | | | | | |
|--|----|----|----|----|----|-----|
| Nossidium pilosellum (Marsham, 1802) | 2 | 0 | 0 | 0 | 0 | 2 |
| Ptenidium (Gressnerium) gressneri Erichson, 1845 | 0 | 0 | 0 | 1 | 0 | 1 |
| Ptenidium (Matthewsium) laevigatum Erichson, 1845 | 8 | 8 | 19 | 8 | 1 | 44 |
| Ptenidium (Matthewsium) turgidum Thomson, C.G., 1855 | 0 | 2 | 0 | 1 | 0 | 3 |
| Ptenidium (Wankowizium) intermedium Wankowicz, 1869 | 5 | 1 | 9 | 5 | 1 | 21 |
| Ptenidium (Ptenidium) formicetorum Kraatz, 1851 | 0 | 0 | 1 | 0 | 0 | 1 |
| Ptenidium (Ptenidium) fuscicorne Erichson, 1845 | 13 | 7 | 27 | 4 | 1 | 52 |
| Ptenidium (Ptenidium) punctatum (Gyllenhal, 1827) | 4 | 5 | 7 | 0 | 2 | 18 |
| Ptenidium (Ptenidium) pusillum (Gyllenhal, 1808) | 27 | 5 | 75 | 22 | 8 | 137 |
| Ptenidium (Gillmeisterium) nitidum (Heer, 1841) | 20 | 22 | 93 | 24 | 15 | 174 |
| Euryptilium gillmeisteri Flach, 1889 | 0 | 0 | 0 | 1 | 0 | 1 |
| Euryptilium saxonicum (Gillmeister, 1845) | 0 | 0 | 2 | 0 | 0 | 2 |
| Ptiliola kunzei (Heer, 1841) | 1 | 3 | 10 | 3 | 0 | 17 |
| Ptiliolium (Ptiliolium) fuscum (Erichson, 1845) | 4 | 14 | 13 | 8 | 4 | 43 |
| Ptiliolium (Ptiliolium) marginatum (Aubé, 1850) | 0 | 0 | 1 | 0 | 0 | 1 |
| Ptiliolium (Ptiliolium) sahlbergi (Flach, 1888) | 0 | 0 | 0 | 1 | 0 | 1 |
| Ptiliolium (Ptiliolium) spencei (Allibert, 1844) | 0 | 4 | 13 | 4 | 2 | 23 |
| Ptiliolium (Euptilium) schwarzi (Flach, 1887) | 0 | 0 | 2 | 0 | 0 | 2 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| <i>Actidium aterrimum</i> (Motschulsky, 1845) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Oligella foveolata</i> (Allibert, 1844) | 1 | 0 | 10 | 2 | 0 | 13 |
| <i>Oligella intermedia</i> Besuchet, 1971 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Ptilium horioni</i> Rosskothén, 1934 | 0 | 0 | 4 | 0 | 0 | 4 |
| <i>Ptilium myrmecophilum</i> (Allibert, 1844) | 0 | 5 | 3 | 0 | 0 | 8 |
| <i>Millidium minutissimum</i> (Ljungh, 1804) | 0 | 1 | 3 | 0 | 1 | 5 |
| <i>Ptinella aptera</i> (Guérin-Méneville, 1839) | 0 | 1 | 2 | 0 | 0 | 3 |
| <i>Ptinella britannica</i> Matthews, A., 1858 | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Ptinella cavelli</i> (Broun, 1893) | 2 | 3 | 9 | 3 | 0 | 17 |
| <i>Ptinella denticollis</i> (Fairmaire, 1858) | 0 | 2 | 3 | 2 | 0 | 7 |
| <i>Ptinella errabunda</i> Johnson, 1975 | 0 | 2 | 14 | 3 | 1 | 20 |
| <i>Pteryx suturalis</i> (Heer, 1841) | 0 | 3 | 8 | 6 | 2 | 19 |
| <i>Nephanes titan</i> (Newman, 1834) | 12 | 5 | 22 | 7 | 7 | 53 |
| <i>Smicrus filicornis</i> (Fairmaire & Laboulbène, 1855) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Baeocrara variolosa</i> (Mulsant & Rey, 1861) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>arnoldi</i> Rosskothén, 1935 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>atomaria</i> (De Geer, 1774) | 11 | 12 | 50 | 20 | 4 | 97 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>cephalotes</i> (Allibert, 1844) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>cognata</i> (Matthews, A., 1877) | 23 | 15 | 115 | 31 | 18 | 202 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>danica</i> Sundt, 1958 | 7 | 3 | 30 | 14 | 3 | 57 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>fascicularis</i> (Herbst, 1793) | 30 | 25 | 122 | 42 | 10 | 229 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>henrici</i> (Matthews, A., 1872) | 10 | 0 | 23 | 2 | 0 | 35 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>insularis</i> (Mäklin, 1852) | 6 | 8 | 20 | 4 | 0 | 38 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>intermedia</i> (Gillmeister, 1845) | 8 | 12 | 82 | 27 | 3 | 132 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>josephi</i> (Matthews, A., 1872) | 3 | 0 | 0 | 0 | 0 | 3 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>lucidula</i> Rosskothén, 1935 | 0 | 1 | 3 | 0 | 0 | 4 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>montandonii</i> (Allibert, 1844) | 6 | 12 | 22 | 7 | 0 | 47 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>norvegica</i> Strand, A., 1941 | 0 | 0 | 2 | 2 | 0 | 4 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>parva</i> Rosskothén, 1935 | 1 | 0 | 2 | 1 | 0 | 4 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>pumila</i> (Erichson, 1845) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>rosskotheni</i> Sundt, 1971 | 0 | 0 | 17 | 6 | 0 | 23 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>rugulosa</i> Rosskothén, 1935 | 0 | 2 | 2 | 1 | 0 | 5 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>sericans</i> (Heer, 1841) | 11 | 6 | 37 | 11 | 3 | 68 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>silvatica</i> Rosskothén, 1935 | 1 | 1 | 8 | 0 | 0 | 10 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>sitkaensis</i> (Motschulsky, 1845) | 20 | 2 | 45 | 25 | 4 | 96 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>strandii</i> Sundt, 1958 | 4 | 1 | 7 | 6 | 1 | 19 |
| <i>Acrotrichis</i> (<i>Acrotrichis</i>) <i>thoracica</i> (Waltl, 1838) | 2 | 7 | 15 | 9 | 3 | 36 |
| <i>Acrotrichis</i> (<i>Ctenopteryx</i>) <i>grandicollis</i> (Mannerheim, 1844) | 22 | 19 | 59 | 17 | 6 | 123 |
| <i>Acrotrichis</i> (<i>Ctenopteryx</i>) <i>sanctae-helenae</i> Johnson, 1972 | 0 | 1 | 1 | 0 | 0 | 2 |
| <u>Leiodidae [93/77]</u> | | | | | | |
| <i>Sogda suturalis</i> (Zetterstedt, 1828) | 0 | 3 | 0 | 0 | 0 | 3 |
| <i>Hydnobius latifrons</i> (Curtis, 1840) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Hydnobius punctatus</i> (Sturm, 1807) | 0 | 3 | 4 | 0 | 0 | 7 |
| <i>Hydnobius spinipes</i> (Gyllenhal, 1813) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Triarthron maerkelii</i> Märkel, 1840 | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Leiodes badia</i> (Sturm, 1807) | 0 | 2 | 2 | 0 | 0 | 4 |
| <i>Leiodes calcarata</i> (Erichson, 1845) | 3 | 8 | 29 | 7 | 3 | 50 |
| <i>Leiodes ciliaris</i> (Schmidt, W.L.E., 1841) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Leiodes cinnamomea</i> (Panzer, 1793) | 1 | 2 | 2 | 0 | 0 | 5 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|---|----|----|---|-----|
| <i>Leiodes ferruginea</i> (Fabricius, 1787) | 2 | 3 | 1 | 1 | 1 | 8 |
| <i>Leiodes furva</i> (Erichson, 1845) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Leiodes gallica</i> (Reitter, 1884) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Leiodes gyllenhalii</i> Stephens, 1829 | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Leiodes litura</i> Stephens, 1832 | 1 | 3 | 3 | 1 | 0 | 8 |
| <i>Leiodes lucens</i> (Fairmaire, 1855) | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Leiodes lunicollis</i> (Rye, 1872) | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Leiodes nigrita</i> (Schmidt, W.L.E., 1841) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Leiodes obesa</i> (Schmidt, W.L.E., 1841) | 1 | 0 | 10 | 5 | 0 | 16 |
| <i>Leiodes picea</i> (Panzer, 1797) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Leiodes rufipennis</i> (Paykull, 1798) | 0 | 0 | 4 | 0 | 1 | 5 |
| <i>Leiodes rugosa</i> Stephens, 1829 | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Leiodes strigipennis</i> Daffner, 1983 | 0 | 0 | 3 | 2 | 0 | 5 |
| <i>Leiodes triepkii</i> (Schmidt, W.L.E., 1841) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Liocyrtusa minuta</i> (Ahrens, 1812) | 1 | 1 | 0 | 1 | 0 | 3 |
| <i>Liocyrtusa vittata</i> (Curtis, 1840) | 1 | 1 | 2 | 1 | 2 | 7 |
| <i>Agaricophagus cephalotes</i> Schmidt, W.L.E., 1841 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Colenis immunda</i> (Sturm, 1807) | 0 | 2 | 6 | 3 | 0 | 11 |
| <i>Anisotoma glabra</i> (Fabricius, 1792) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Anisotoma humeralis</i> (Fabricius, 1792) | 5 | 9 | 67 | 10 | 4 | 95 |
| <i>Anisotoma orbicularis</i> (Herbst, 1792) | 0 | 5 | 8 | 1 | 0 | 14 |
| <i>Amphicyllis globus</i> (Fabricius, 1792) | 0 | 3 | 10 | 1 | 0 | 14 |
| <i>Agathidium</i> (Cyphocele) <i>arcticum</i> Thomson, C.G., 1862 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Agathidium</i> (Cyphocele) <i>nigrinum</i> Sturm, 1807 | 2 | 3 | 18 | 4 | 0 | 27 |
| <i>Agathidium</i> (Neoceleble) <i>convexum</i> Sharp, 1866 | 2 | 0 | 0 | 1 | 0 | 3 |
| <i>Agathidium</i> (Neoceleble) <i>marginatum</i> Sturm, 1807 | 2 | 1 | 0 | 0 | 0 | 3 |
| <i>Agathidium</i> (Neoceleble) <i>nigripenne</i> (Fabricius, 1792) | 2 | 5 | 29 | 14 | 3 | 53 |
| <i>Agathidium</i> (Neoceleble) <i>rotundatum</i> (Gyllenhal, 1827) | 0 | 5 | 4 | 0 | 0 | 9 |
| <i>Agathidium</i> (Neoceleble) <i>varians</i> Beck, 1817 | 10 | 6 | 32 | 5 | 1 | 54 |
| <i>Agathidium</i> (Agathidium) <i>atrum</i> (Paykull, 1798) | 1 | 4 | 19 | 3 | 0 | 27 |
| <i>Agathidium</i> (Agathidium) <i>laevigatum</i> Erichson, 1845 | 18 | 3 | 23 | 4 | 1 | 49 |
| <i>Agathidium</i> (Agathidium) <i>pisanum</i> Brisout, 1872 | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Agathidium</i> (Agathidium) <i>seminulum</i> (Linnaeus, 1758) | 0 | 2 | 3 | 5 | 0 | 10 |
| <i>Colon</i> (Eurycolon) <i>latum</i> Kraatz, 1850 | 0 | 3 | 2 | 0 | 0 | 5 |
| <i>Colon</i> (Myloechus) <i>appendiculatum</i> (Sahlberg, C.R., 1822) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Colon</i> (Myloechus) <i>brunneum</i> (Latreille, 1807) | 1 | 2 | 3 | 3 | 0 | 9 |
| <i>Colon</i> (Myloechus) <i>dentipes</i> (Sahlberg, C.R., 1822) | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Colon</i> (Colon) <i>serripes</i> (Sahlberg, C.R., 1822) | 0 | 2 | 1 | 0 | 0 | 3 |
| <i>Nemadus colonoides</i> (Kraatz, 1851) | 1 | 1 | 0 | 1 | 0 | 3 |
| <i>Nargus</i> (Nargus) <i>velox</i> (Spence, 1813) | 21 | 6 | 95 | 17 | 6 | 145 |
| <i>Nargus</i> (Nargus) <i>wilkinii</i> (Spence, 1813) | 3 | 6 | 59 | 5 | 1 | 74 |
| <i>Nargus</i> (Demochrus) <i>anisotomoides</i> (Spence, 1813) | 1 | 4 | 16 | 0 | 0 | 21 |
| <i>Choleva</i> (Cholevopsis) <i>spadicea</i> (Sturm, 1839) | 0 | 2 | 7 | 0 | 0 | 9 |
| <i>Choleva</i> (Choleva) <i>agilis</i> (Illiger, 1798) | 2 | 3 | 9 | 0 | 0 | 14 |
| <i>Choleva</i> (Choleva) <i>angustata</i> (Fabricius, 1781) | 7 | 2 | 12 | 3 | 0 | 24 |
| <i>Choleva</i> (Choleva) <i>cisteloides</i> (Frölich, 1799) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Choleva</i> (Choleva) <i>fagniezi</i> Jeannel, 1922 | 0 | 0 | 10 | 5 | 0 | 15 |
| <i>Choleva</i> (Choleva) <i>glauca</i> Britten, 1918 | 1 | 0 | 2 | 4 | 0 | 7 |
| <i>Choleva</i> (Choleva) <i>jeanneli</i> Britten, 1922 | 7 | 2 | 10 | 5 | 0 | 24 |
| <i>Choleva</i> (Choleva) <i>oblonga</i> Latreille, 1807 | 1 | 1 | 6 | 4 | 0 | 12 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|---|-----|
| <i>Sciodrepoides fumatus</i> (Spence, 1813) | 3 | 6 | 18 | 11 | 2 | 40 |
| <i>Sciodrepoides watsoni</i> (Spence, 1813) | 17 | 14 | 70 | 19 | 3 | 123 |
| <i>Catops chrysomeloides</i> (Panzer, 1798) | 9 | 4 | 10 | 8 | 0 | 31 |
| <i>Catops coracinus</i> Kellner, 1846 | 5 | 5 | 15 | 6 | 3 | 34 |
| <i>Catops fuliginosus</i> Erichson, 1837 | 15 | 4 | 66 | 11 | 0 | 96 |
| <i>Catops fuscus</i> (Panzer, 1794) | 3 | 1 | 4 | 1 | 0 | 9 |
| <i>Catops grandicollis</i> Erichson, 1837 | 11 | 4 | 32 | 8 | 4 | 59 |
| <i>Catops kirbii</i> (Spence, 1813) | 6 | 7 | 21 | 5 | 2 | 41 |
| <i>Catops longulus</i> Kellner, 1846 | 0 | 5 | 7 | 3 | 1 | 16 |
| <i>Catops morio</i> (Fabricius, 1787) | 20 | 8 | 32 | 11 | 2 | 73 |
| <i>Catops nigricans</i> (Spence, 1813) | 17 | 3 | 35 | 6 | 1 | 62 |
| <i>Catops nigrita</i> Erichson, 1837 | 0 | 5 | 43 | 3 | 2 | 53 |
| <i>Catops tristis</i> (Panzer, 1793) | 11 | 12 | 97 | 15 | 2 | 137 |
| <i>Catopidius depressus</i> (Murray, 1856) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Ptomaphagus medius</i> Rey, 1889 | 3 | 3 | 10 | 3 | 1 | 20 |
| <i>Ptomaphagus subvillosus</i> (Goeze, 1777) | 14 | 19 | 109 | 33 | 3 | 178 |
| <i>Ptomaphagus varicornis</i> (Rosenhauer, 1847) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Leptinus testaceus</i> Müller, P.W.J., 1817 | 1 | 5 | 8 | 3 | 0 | 17 |

Scymaenidae [32/18]

| | | | | | | |
|--|----|----|----|----|---|-----|
| <i>Eutheia schaumii</i> Kiesenwetter, 1858 | 0 | 2 | 1 | 0 | 0 | 3 |
| <i>Eutheia scydmaenoides</i> Stephens, 1830 | 0 | 2 | 2 | 1 | 0 | 5 |
| <i>Cephennium gallicum</i> Ganglbauer, 1899 | 2 | 0 | 5 | 0 | 0 | 7 |
| <i>Euconnus</i> (<i>Cladoconnus</i>) <i>denticornis</i> (Müller, P.W.J. & Kunze, 1822) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Euconnus</i> (<i>Euconnus</i>) <i>fimetarius</i> (Chaudoir, 1845) | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Euconnus</i> (<i>Euconnus</i>) <i>hirticollis</i> (Illiger, 1798) | 1 | 2 | 1 | 8 | 0 | 12 |
| <i>Euconnus</i> (<i>Napochus</i>) <i>duboisii</i> Méquignon, 1929 | 2 | 0 | 1 | 0 | 0 | 3 |
| <i>Microscydmus nanus</i> (Schaum, 1844) | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Neuraphes</i> (<i>Neuraphes</i>) <i>angulatus</i> (Müller, P.W.J. & Kunze, 1822) | 0 | 3 | 1 | 0 | 0 | 4 |
| <i>Neuraphes</i> (<i>Neuraphes</i>) <i>elongatulus</i> (Müller, P.W.J. & Kunze, 1822) | 0 | 9 | 27 | 7 | 1 | 44 |
| <i>Neuraphes</i> (<i>Neuraphes</i>) <i>talparum</i> Lokay, 1921 | 0 | 3 | 0 | 1 | 0 | 4 |
| <i>Neuraphes</i> (<i>Pararaphes</i>) <i>plicicollis</i> Reitter, 1879 | 0 | 0 | 0 | 3 | 0 | 3 |
| <i>Scydmorephes helvolus</i> (Schaum, 1844) | 4 | 0 | 5 | 1 | 0 | 10 |
| <i>Scydmorephes sparshalli</i> (Denny, 1825) | 0 | 2 | 0 | 1 | 0 | 3 |
| <i>Stenichnus bicolor</i> (Denny, 1825) | 0 | 2 | 8 | 3 | 0 | 13 |
| <i>Stenichnus collaris</i> (Müller, P.W.J. & Kunze, 1822) | 8 | 23 | 42 | 18 | 2 | 93 |
| <i>Stenichnus scutellaris</i> (Müller, P.W.J. & Kunze, 1822) | 8 | 4 | 9 | 1 | 0 | 22 |
| <i>Scydmaenus</i> (<i>Scydmaenus</i>) <i>tarsatus</i> Müller, P.W.J. & Kunze, 1822 | 19 | 10 | 91 | 18 | 2 | 140 |

Silphidae [21/19]

| | | | | | | |
|--|----|----|----|----|----|-----|
| <i>Nicrodes littoralis</i> (Linnaeus, 1758) | 6 | 5 | 19 | 9 | 28 | 67 |
| <i>Thanatophilus dispar</i> (Herbst, 1793) | 0 | 1 | 0 | 2 | 0 | 3 |
| <i>Thanatophilus rugosus</i> (Linnaeus, 1758) | 27 | 6 | 47 | 22 | 9 | 111 |
| <i>Thanatophilus sinuatus</i> (Fabricius, 1775) | 9 | 4 | 21 | 6 | 1 | 41 |
| <i>Oiceoptoma thoracicum</i> (Linnaeus, 1758) | 9 | 25 | 32 | 8 | 0 | 74 |
| <i>Aclypea opaca</i> (Linnaeus, 1758) | 5 | 7 | 6 | 6 | 0 | 24 |
| <i>Aclypea undata</i> (Müller, O.F., 1776) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Dendroxena quadrimaculata</i> (Scopoli, 1772) | 3 | 2 | 22 | 2 | 0 | 29 |
| <i>Silpha atrata</i> Linnaeus, 1758 | 12 | 27 | 49 | 15 | 7 | 110 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|-----|-----|
| <i>Silpha laevigata</i> Fabricius, 1775 | 0 | 2 | 3 | 0 | 0 | 5 |
| <i>Silpha obscura</i> Linnaeus, 1758 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Silpha tristis</i> Illiger, 1798 | 0 | 1 | 42 | 7 | 1 | 51 |
| <i>Silpha tyrolensis</i> Laicharting, 1781 | 0 | 3 | 5 | 2 | 3 | 13 |
| <i>Nicrophorus humator</i> (Gleditsch, 1767) | 22 | 3 | 87 | 40 | 127 | 279 |
| <i>Nicrophorus interruptus</i> Stephens, 1830 | 0 | 1 | 3 | 0 | 0 | 4 |
| <i>Nicrophorus investigator</i> Zetterstedt, 1824 | 12 | 4 | 33 | 28 | 105 | 182 |
| <i>Nicrophorus vespillo</i> (Linnaeus, 1758) | 10 | 7 | 36 | 15 | 4 | 72 |
| <i>Nicrophorus vespilloides</i> Herbst, 1783 | 6 | 20 | 123 | 6 | 3 | 158 |
| <i>Nicrophorus vestigator</i> Herschel, 1807 | 0 | 4 | 0 | 0 | 0 | 4 |

Staphylinidae [1080/802]

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Acidota crenata</i> (Fabricius, 1792) | 2 | 12 | 12 | 10 | 4 | 40 |
| <i>Acidota cruentata</i> Mannerheim, 1830 | 0 | 3 | 8 | 7 | 0 | 18 |
| <i>Anthobium atrocephalum</i> (Gyllenhal, 1827) | 14 | 16 | 156 | 18 | 1 | 205 |
| <i>Anthobium unicolor</i> (Marsham, 1802) | 34 | 25 | 194 | 48 | 7 | 308 |
| <i>Anthophagus</i> (<i>Anthophagus</i>) <i>alpinus</i> (Paykull, 1790) | 0 | 0 | 0 | 4 | 0 | 4 |
| <i>Anthophagus</i> (<i>Phaganthus</i>) <i>caraboides</i> (Linnaeus, 1758) | 0 | 27 | 4 | 9 | 4 | 44 |
| <i>Deliphrum tectum</i> (Paykull, 1789) | 2 | 3 | 16 | 4 | 2 | 27 |
| <i>Eucnecosum brachypterum</i> (Gravenhorst, 1802) | 0 | 1 | 6 | 8 | 5 | 20 |
| <i>Geodromicus nigrita</i> (Müller, P.W.J., 1821) | 0 | 3 | 4 | 7 | 4 | 18 |
| <i>Lesteva hanseni</i> Lohse, 1953 | 0 | 0 | 0 | 3 | 0 | 3 |
| <i>Lesteva longoelytrata</i> (Goeze, 1777) | 43 | 43 | 125 | 48 | 20 | 279 |
| <i>Lesteva monticola</i> Kiesenwetter, 1847 | 0 | 4 | 2 | 8 | 10 | 24 |
| <i>Lesteva pubescens</i> Mannerheim, 1830 | 1 | 11 | 12 | 5 | 2 | 31 |
| <i>Lesteva punctata</i> Erichson, 1839 | 3 | 9 | 11 | 6 | 2 | 31 |
| <i>Lesteva sicula</i> subsp. <i>heeri</i> Fauvel, 1871 | 37 | 40 | 166 | 55 | 13 | 311 |
| <i>Olophrum consimile</i> (Gyllenhal, 1810) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Olophrum fuscum</i> (Gravenhorst, 1806) | 2 | 3 | 20 | 8 | 6 | 39 |
| <i>Olophrum piceum</i> (Gyllenhal, 1810) | 20 | 33 | 123 | 26 | 10 | 212 |
| <i>Philorinum sordidum</i> (Stephens, 1834) | 13 | 13 | 25 | 6 | 4 | 61 |
| <i>Phyllodrepoidea crenata</i> Ganglbauer, 1895 | 0 | 8 | 11 | 3 | 0 | 22 |
| <i>Eusphalerum luteum</i> (Marsham, 1802) | 1 | 8 | 18 | 6 | 1 | 34 |
| <i>Eusphalerum minutum</i> (Fabricius, 1792) | 8 | 10 | 4 | 36 | 7 | 65 |
| <i>Eusphalerum primulae</i> (Stephens, 1834) | 3 | 12 | 14 | 4 | 0 | 33 |
| <i>Eusphalerum sorbi</i> (Gyllenhal, 1810) | 4 | 20 | 12 | 16 | 4 | 56 |
| <i>Eusphalerum torquatum</i> (Marsham, 1802) | 3 | 4 | 14 | 11 | 8 | 40 |
| <i>Acrolocha minuta</i> (Olivier, 1795) | 0 | 1 | 2 | 0 | 0 | 3 |
| <i>Acrolocha sulcula</i> (Stephens, 1834) | 5 | 8 | 11 | 4 | 2 | 30 |
| <i>Acrulia inflata</i> (Gyllenhal, 1813) | 3 | 6 | 43 | 8 | 0 | 60 |
| <i>Dropephylla devillei</i> Bernhauer, 1902 | 0 | 6 | 7 | 4 | 0 | 17 |
| <i>Dropephylla ioptera</i> (Stephens, 1832) | 6 | 4 | 8 | 7 | 1 | 26 |
| <i>Dropephylla vilis</i> (Erichson, 1840) | 3 | 11 | 43 | 9 | 0 | 66 |
| <i>Hapalareae pygmaea</i> (Paykull, 1800) | 0 | 1 | 7 | 3 | 0 | 11 |
| <i>Micralymma marinum</i> (Ström, 1783) | 4 | 9 | 1 | 0 | 0 | 14 |
| <i>Omalius</i> Gravenhorst, 1802 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Omalius allardi</i> Fairmaire & Brisout, 1859 | 0 | 3 | 4 | 4 | 0 | 11 |
| <i>Omalius caesum</i> Gravenhorst, 1806 | 36 | 9 | 86 | 16 | 2 | 149 |
| <i>Omalius excavatum</i> Stephens, 1834 | 26 | 8 | 33 | 15 | 2 | 84 |
| <i>Omalius exiguum</i> Gyllenhal, 1810 | 0 | 1 | 2 | 0 | 0 | 3 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|---|-----|
| <i>Omalium italicum</i> Bernhauer, 1902 | 7 | 2 | 32 | 4 | 0 | 45 |
| <i>Omalium laeviusculum</i> Gyllenhal, 1827 | 5 | 6 | 0 | 0 | 0 | 11 |
| <i>Omalium laticolle</i> Kraatz, 1858 | 0 | 0 | 3 | 4 | 0 | 7 |
| <i>Omalium oxyacanthae</i> Gravenhorst, 1806 | 2 | 4 | 9 | 6 | 1 | 22 |
| <i>Omalium riparium</i> Thomson, C.G., 1857 | 10 | 3 | 0 | 0 | 0 | 13 |
| <i>Omalium rivulare</i> (Paykull, 1789) | 39 | 19 | 181 | 34 | 7 | 280 |
| <i>Omalium rugatum</i> Mulsant & Rey, 1880 | 0 | 1 | 7 | 3 | 0 | 11 |
| <i>Omalium rugulipenne</i> Rye, 1864 | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Omalium septentrionis</i> Thomson, C.G., 1856 | 0 | 6 | 8 | 3 | 0 | 17 |
| <i>Phloeonomus punctipennis</i> Thomson, C.G., 1867 | 3 | 6 | 24 | 12 | 3 | 48 |
| <i>Phloeonomus pusillus</i> (Gravenhorst, 1806) | 3 | 5 | 20 | 2 | 1 | 31 |
| <i>Phloeostiba plana</i> (Paykull, 1792) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Phyllodrepa floralis</i> (Paykull, 1789) | 13 | 6 | 11 | 5 | 0 | 35 |
| <i>Phyllodrepa puberula</i> Bernhauer, 1903 | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Phyllodrepa salicis</i> (Gyllenhal, 1810) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Xylodromus concinnus</i> (Marsham, 1802) | 14 | 16 | 16 | 8 | 2 | 56 |
| <i>Xylodromus depressus</i> (Gravenhorst, 1802) | 5 | 3 | 7 | 6 | 1 | 22 |
| <i>Xylostiba monilicornis</i> (Gyllenhal, 1810) | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Coryphium angusticolle</i> Stephens, 1834 | 3 | 3 | 27 | 6 | 0 | 39 |
| <i>Euedectus whitei</i> Sharp, 1871 | 0 | 0 | 0 | 3 | 0 | 3 |
| <i>Megarathrus bellevoeyi</i> (Saulcy, 1862) | 4 | 5 | 19 | 0 | 0 | 28 |
| <i>Megarathrus denticollis</i> (Beck, 1817) | 11 | 11 | 80 | 21 | 6 | 129 |
| <i>Megarathrus depressus</i> (Paykull, 1789) | 21 | 19 | 138 | 37 | 3 | 218 |
| <i>Megarathrus prosseni</i> Schatzmayr, 1904 | 1 | 0 | 7 | 0 | 0 | 8 |
| <i>Metopsia clypeata</i> (Müller, P.W.J., 1821) | 24 | 10 | 56 | 28 | 0 | 118 |
| <i>Proteinus atomarius</i> Erichson, 1840 | 0 | 1 | 6 | 2 | 0 | 9 |
| <i>Proteinus brachypterus</i> (Fabricius, 1792) | 10 | 24 | 90 | 22 | 3 | 149 |
| <i>Proteinus crenulatus</i> Pandellé, 1867 | 0 | 3 | 12 | 2 | 0 | 17 |
| <i>Proteinus laevigatus</i> Hochhuth, 1871 | 6 | 1 | 31 | 9 | 1 | 48 |
| <i>Proteinus ovalis</i> Stephens, 1834 | 39 | 14 | 190 | 32 | 3 | 278 |
| <i>Arrhenopeplus tesserula</i> Curtis, 1828 | 0 | 0 | 2 | 3 | 1 | 6 |
| <i>Micropeplus fulvus</i> Erichson, 1840 | 14 | 6 | 46 | 18 | 0 | 84 |
| <i>Micropeplus porcatus</i> (Paykull, 1789) | 2 | 10 | 1 | 4 | 0 | 17 |
| <i>Micropeplus staphylinoides</i> (Marsham, 1802) | 17 | 7 | 33 | 6 | 1 | 64 |
| <i>Trichonyx sulcicollis</i> (Reichenbach, 1816) | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Trimium brevicorne</i> (Reichenbach, 1816) | 0 | 2 | 1 | 1 | 0 | 4 |
| <i>Bibloporus bicolor</i> (Denny, 1825) | 0 | 2 | 7 | 0 | 0 | 9 |
| <i>Bibloporus minutus</i> Raffray, 1914 | 0 | 3 | 1 | 1 | 0 | 5 |
| <i>Biblopectus ambiguus</i> (Reichenbach, 1816) | 1 | 1 | 1 | 4 | 0 | 7 |
| <i>Biblopectus pusillus</i> (Denny, 1825) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Biblopectus tenebrosus</i> (Reitter, 1880) | 0 | 0 | 0 | 3 | 0 | 3 |
| <i>Euplectus bescidicus</i> Reitter, 1881 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Euplectus karstenii</i> (Reichenbach, 1816) | 0 | 0 | 3 | 3 | 0 | 6 |
| <i>Euplectus nanus</i> (Reichenbach, 1816) | 3 | 1 | 3 | 1 | 1 | 9 |
| <i>Euplectus piceus</i> Motschulsky, 1835 | 0 | 2 | 1 | 2 | 0 | 5 |
| <i>Euplectus sanguineus</i> Denny, 1825 | 6 | 4 | 21 | 3 | 2 | 36 |
| <i>Euplectus signatus</i> (Reichenbach, 1816) | 0 | 2 | 10 | 0 | 0 | 12 |
| <i>Plectophloeus nitidus</i> (Fairmaire, 1857) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Batrisodes venustus</i> (Reichenbach, 1816) | 1 | 1 | 0 | 1 | 0 | 3 |
| <i>Bryaxis bulbifer</i> (Reichenbach, 1816) | 10 | 9 | 22 | 35 | 2 | 78 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|-----|----|-----|-----|----|-----|
| <i>Bryaxis curtisii</i> (Leach, 1817) | 0 | 4 | 5 | 7 | 0 | 16 |
| <i>Bryaxis puncticollis</i> (Denny, 1825) | 5 | 12 | 17 | 19 | 1 | 54 |
| <i>Bythinus burrellii</i> Denny, 1825 | 2 | 10 | 7 | 15 | 2 | 36 |
| <i>Bythinus macropalpus</i> Aubé, 1833 | 3 | 2 | 9 | 3 | 0 | 17 |
| <i>Tychus niger</i> (Paykull, 1800) | 14 | 19 | 22 | 13 | 3 | 71 |
| <i>Brachygluta fossulata</i> (Reichenbach, 1816) | 8 | 10 | 27 | 19 | 0 | 64 |
| <i>Brachygluta helferi</i> (Schmidt-Göbel, 1836) | 19 | 0 | 0 | 0 | 0 | 19 |
| <i>Fagniezia impressus</i> (Panzer, 1803) | 0 | 2 | 1 | 3 | 0 | 6 |
| <i>Reichenbachia juncorum</i> (Leach, 1817) | 5 | 4 | 3 | 5 | 0 | 17 |
| <i>Rybaxis longicornis</i> (Leach, 1817) | 11 | 0 | 46 | 8 | 1 | 66 |
| <i>Pselaphaulax dresdensis</i> (Herbst, 1792) | 2 | 0 | 1 | 2 | 0 | 5 |
| <i>Pselaphus heisei</i> Herbst, 1792 | 6 | 9 | 9 | 14 | 3 | 41 |
| <i>Claviger testaceus</i> Preyssler, 1790 | 0 | 3 | 0 | 0 | 0 | 3 |
| <i>Phloeocharis subtilissima</i> Mannerheim, 1830 | 4 | 9 | 29 | 9 | 2 | 53 |
| <i>Cilea silphoides</i> (Linnaeus, 1767) | 8 | 9 | 16 | 5 | 0 | 38 |
| <i>Lamprinodes saginatus</i> (Gravenhorst, 1806) | 8 | 2 | 5 | 0 | 0 | 15 |
| <i>Sepedophilus constans</i> (Fowler, 1888) | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Sepedophilus immaculatus</i> (Stephens, 1832) | 14 | 2 | 16 | 1 | 0 | 33 |
| <i>Sepedophilus littoreus</i> (Linnaeus, 1758) | 3 | 6 | 21 | 2 | 0 | 32 |
| <i>Sepedophilus lusitanicus</i> Hammond, 1973 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Sepedophilus marshami</i> (Stephens, 1832) | 66 | 22 | 209 | 23 | 6 | 326 |
| <i>Sepedophilus nigripennis</i> (Stephens, 1832) | 76 | 20 | 200 | 27 | 7 | 330 |
| <i>Sepedophilus pedicularius</i> (Gravenhorst, 1802) | 1 | 0 | 8 | 10 | 0 | 19 |
| <i>Sepedophilus testaceus</i> (Fabricius, 1792) | 2 | 0 | 0 | 2 | 0 | 4 |
| <i>Tachinus corticinus</i> Gravenhorst, 1802 | 8 | 10 | 21 | 36 | 4 | 79 |
| <i>Tachinus elongatus</i> Gyllenhal, 1810 | 0 | 15 | 8 | 4 | 4 | 31 |
| <i>Tachinus humeralis</i> Gravenhorst, 1802 | 9 | 8 | 46 | 7 | 2 | 72 |
| <i>Tachinus laticollis</i> Gravenhorst, 1802 | 14 | 18 | 100 | 13 | 4 | 149 |
| <i>Tachinus lignorum</i> (Linnaeus, 1758) | 0 | 1 | 3 | 3 | 0 | 7 |
| <i>Tachinus marginellus</i> (Fabricius, 1781) | 33 | 27 | 207 | 54 | 17 | 338 |
| <i>Tachinus pallipes</i> (Gravenhorst, 1806) | 4 | 10 | 50 | 8 | 1 | 73 |
| <i>Tachinus proximus</i> Kraatz, 1855 | 2 | 5 | 20 | 7 | 1 | 35 |
| <i>Tachinus rufipennis</i> Gyllenhal, 1810 | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Tachinus rufipes</i> (Linnaeus, 1758) | 73 | 57 | 471 | 108 | 17 | 726 |
| <i>Tachinus subterraneus</i> (Linnaeus, 1758) | 9 | 4 | 33 | 8 | 1 | 55 |
| <i>Tachyporus</i> Gravenhorst, 1802 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Tachyporus atriceps</i> Stephens, 1832 | 9 | 7 | 11 | 6 | 0 | 33 |
| <i>Tachyporus chrysomelinus</i> (Linnaeus, 1758) | 20 | 3 | 218 | 26 | 3 | 270 |
| <i>Tachyporus dispar</i> (Paykull, 1789) | 65 | 68 | 338 | 73 | 15 | 559 |
| <i>Tachyporus formosus</i> Matthews, A.H., 1838 | 0 | 1 | 0 | 1 | 1 | 3 |
| <i>Tachyporus hypnorum</i> (Fabricius, 1775) | 139 | 85 | 584 | 98 | 18 | 924 |
| <i>Tachyporus nitidulus</i> (Fabricius, 1781) | 52 | 39 | 237 | 38 | 7 | 373 |
| <i>Tachyporus obtusus</i> (Linnaeus, 1767) | 81 | 51 | 449 | 63 | 15 | 659 |
| <i>Tachyporus pallidus</i> Sharp, 1871 | 38 | 24 | 116 | 25 | 2 | 205 |
| <i>Tachyporus pusillus</i> Gravenhorst, 1806 | 8 | 4 | 32 | 9 | 3 | 56 |
| <i>Tachyporus quadriscopulatus</i> Pandellé, 1869 | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Tachyporus solutus</i> Erichson, 1839 | 22 | 8 | 256 | 2 | 1 | 289 |
| <i>Tachyporus tersus</i> Erichson, 1839 | 0 | 10 | 8 | 5 | 1 | 24 |
| <i>Tachyporus transversalis</i> Gravenhorst, 1806 | 6 | 6 | 28 | 4 | 0 | 44 |
| <i>Bolitobius cingulatus</i> (Mannerheim, 1830) | 5 | 8 | 28 | 6 | 1 | 48 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Ischnosoma splendidum</i> (Gravenhorst, 1806) | 0 | 0 | 8 | 2 | 0 | 10 |
| <i>Lordithon exoletus</i> (Erichson, 1839) | 2 | 5 | 9 | 1 | 0 | 17 |
| <i>Lordithon lunulatus</i> (Linnaeus, 1761) | 4 | 12 | 34 | 10 | 5 | 65 |
| <i>Lordithon thoracicus</i> (Fabricius, 1777) | 4 | 12 | 58 | 18 | 2 | 94 |
| <i>Lordithon trinotatus</i> (Erichson, 1839) | 8 | 16 | 73 | 11 | 3 | 111 |
| <i>Mycetoporus angularis</i> Mulsant & Rey, 1853 | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Mycetoporus baudueri</i> Mulsant & Rey, 1875 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Mycetoporus clavicornis</i> (Stephens, 1832) | 3 | 14 | 23 | 5 | 2 | 47 |
| <i>Mycetoporus despectus</i> Strand, A., 1969 | 0 | 0 | 4 | 1 | 1 | 6 |
| <i>Mycetoporus lepidus</i> (Gravenhorst, 1806) | 8 | 25 | 45 | 21 | 3 | 102 |
| <i>Mycetoporus longulus</i> Mannerheim, 1830 | 4 | 8 | 10 | 10 | 2 | 34 |
| <i>Mycetoporus nigricollis</i> Stephens, 1835 | 20 | 16 | 38 | 14 | 1 | 89 |
| <i>Mycetoporus piceolus</i> Rey, 1883 | 2 | 1 | 0 | 0 | 0 | 3 |
| <i>Mycetoporus punctus</i> (Gravenhorst, 1806) | 2 | 3 | 4 | 0 | 0 | 9 |
| <i>Mycetoporus rufescens</i> (Stephens, 1832) | 2 | 12 | 15 | 4 | 2 | 35 |
| <i>Parabolitobius inclinans</i> (Gravenhorst, 1806) | 0 | 9 | 6 | 4 | 0 | 19 |
| <i>Trichophya pilicornis</i> (Gyllenhal, 1810) | 0 | 3 | 4 | 0 | 0 | 7 |
| <i>Habrocerus capillaricornis</i> (Gravenhorst, 1806) | 33 | 13 | 108 | 40 | 8 | 202 |
| <i>Deinopsis erosa</i> (Stephens, 1832) | 4 | 0 | 4 | 2 | 0 | 10 |
| <i>Gymnusa brevicollis</i> (Paykull, 1800) | 3 | 9 | 9 | 2 | 0 | 23 |
| <i>Gymnusa variegata</i> Kiesenwetter, 1845 | 2 | 3 | 7 | 0 | 2 | 14 |
| <i>Cypha discoidea</i> (Erichson, 1839) | 3 | 0 | 0 | 0 | 0 | 3 |
| <i>Cypha laeviuscula</i> (Mannerheim, 1830) | 15 | 8 | 22 | 5 | 1 | 51 |
| <i>Cypha longicornis</i> (Paykull, 1800) | 33 | 24 | 96 | 9 | 2 | 164 |
| <i>Cypha pulicaria</i> (Erichson, 1839) | 4 | 3 | 5 | 1 | 0 | 13 |
| <i>Cypha punctum</i> (Motschulsky, 1857) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Cypha seminulum</i> (Erichson, 1839) | 0 | 3 | 0 | 1 | 0 | 4 |
| <i>Oligota apicata</i> (Erichson, 1837) | 5 | 3 | 5 | 2 | 2 | 17 |
| <i>Holobus flavicornis</i> (Lacordaire, 1835) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Oligota granaria</i> Erichson, 1837 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Oligota inflata</i> (Mannerheim, 1830) | 11 | 13 | 19 | 2 | 0 | 45 |
| <i>Oligota parva</i> Kraatz, 1862 | 8 | 1 | 14 | 1 | 0 | 24 |
| <i>Oligota picipes</i> (Stephens, 1832) | 4 | 2 | 3 | 0 | 0 | 9 |
| <i>Oligota pumilio</i> Kiesenwetter, 1858 | 1 | 0 | 1 | 1 | 0 | 3 |
| <i>Oligota punctulata</i> Heer, 1839 | 4 | 1 | 39 | 8 | 4 | 56 |
| <i>Oligota pusillima</i> (Gravenhorst, 1806) | 9 | 3 | 9 | 1 | 0 | 22 |
| <i>Myllaena brevicornis</i> (Matthews, A.H., 1838) | 7 | 13 | 39 | 15 | 12 | 86 |
| <i>Myllaena dubia</i> (Gravenhorst, 1806) | 15 | 4 | 22 | 2 | 4 | 47 |
| <i>Myllaena elongata</i> (Matthews, A.H., 1838) | 3 | 2 | 1 | 0 | 0 | 6 |
| <i>Myllaena gracilis</i> (Matthews, A.H., 1838) | 3 | 0 | 0 | 2 | 0 | 5 |
| <i>Myllaena graeca</i> Kraatz, 1858 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Myllaena infuscata</i> Kraatz, 1853 | 8 | 2 | 8 | 9 | 1 | 28 |
| <i>Myllaena intermedia</i> Erichson, 1837 | 3 | 2 | 5 | 1 | 0 | 11 |
| <i>Myllaena kraatzi</i> Sharp, 1871 | 2 | 4 | 2 | 0 | 1 | 9 |
| <i>Myllaena minuta</i> (Gravenhorst, 1806) | 6 | 1 | 7 | 10 | 2 | 26 |
| <i>Diglotta mersa</i> (Haliday, 1837) | 3 | 0 | 0 | 0 | 0 | 3 |
| <i>Hygronoma dimidiata</i> (Gravenhorst, 1806) | 21 | 5 | 41 | 12 | 3 | 82 |
| <i>Encephalus complicans</i> Stephens, 1832 | 5 | 5 | 22 | 8 | 1 | 41 |
| <i>Gyrophaena affinis</i> Mannerheim, 1830 | 4 | 5 | 16 | 5 | 0 | 30 |
| <i>Gyrophaena manca</i> Erichson, 1839 | 3 | 9 | 32 | 10 | 1 | 55 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|---|-----|
| <i>Gyrophana bihamata</i> Thomson,C.G.,1867 | 2 | 0 | 8 | 2 | 0 | 12 |
| <i>Gyrophana congrua</i> Erichson,1837 | 1 | 2 | 1 | 0 | 0 | 4 |
| <i>Gyrophana fasciata</i> (Marsham,1802) | 3 | 8 | 21 | 12 | 1 | 45 |
| <i>Gyrophana gentilis</i> Erichson,1839 | 4 | 13 | 41 | 11 | 3 | 72 |
| <i>Gyrophana joyi</i> Wendeler,1924 | 0 | 0 | 0 | 1 | 1 | 2 |
| <i>Gyrophana joyioides</i> Wusthoff,1937 | 0 | 0 | 7 | 5 | 0 | 12 |
| <i>Agaricochara latissima</i> (Stephens,1832) | 2 | 5 | 43 | 7 | 2 | 59 |
| <i>Gyrophana lucidula</i> Erichson,1837 | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Gyrophana minima</i> Erichson,1837 | 1 | 4 | 4 | 6 | 0 | 15 |
| <i>Gyrophana munsteri</i> Strand,A.,1935 | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Gyrophana nana</i> (Paykull,1800) | 2 | 0 | 1 | 1 | 1 | 5 |
| <i>Gyrophana poweri</i> Crotch,1867 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Gyrophana strictula</i> Erichson,1839 | 1 | 0 | 4 | 0 | 0 | 5 |
| <i>Gyrophana williamsi</i> Strand,A.,1935 | 0 | 0 | 1 | 1 | 0 | 2 |
| <i>Placusa depressa</i> Maklin,1845 | 0 | 1 | 3 | 0 | 0 | 4 |
| <i>Placusa tachyporoides</i> (Waltl,1838) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Homalota plana</i> (Gyllenhal,1810) | 2 | 7 | 15 | 7 | 1 | 32 |
| <i>Anomognathus cuspidatus</i> (Erichson,1839) | 1 | 4 | 8 | 0 | 1 | 14 |
| <i>Phytosus balticus</i> Kraatz,1859 | 4 | 2 | 0 | 0 | 0 | 6 |
| <i>Phytosus spinifer</i> Curtis,1838 | 3 | 1 | 0 | 0 | 0 | 4 |
| <i>Arena tabida</i> (Kiesenwetter,1850) | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Thecturota marchii</i> (Dodero,1922) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Leptusa fumida</i> (Erichson,1839) | 11 | 45 | 217 | 21 | 6 | 300 |
| <i>Leptusa norvegica</i> Strand,A.,1941 | 0 | 3 | 1 | 0 | 0 | 4 |
| <i>Leptusa pulchella</i> (Mannerheim,1830) | 0 | 9 | 15 | 4 | 2 | 30 |
| <i>Leptusa ruficollis</i> (Erichson,1839) | 3 | 19 | 31 | 8 | 1 | 62 |
| <i>Heterota plumbea</i> (Waterhouse,G.R.,1858) | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Bolitochara bella</i> Markel,1844 | 2 | 0 | 1 | 1 | 0 | 4 |
| <i>Bolitochara lucida</i> (Gravenhorst,1802) | 2 | 5 | 16 | 7 | 0 | 30 |
| <i>Bolitochara mulsanti</i> Sharp,1875 | 0 | 3 | 5 | 0 | 1 | 9 |
| <i>Bolitochara obliqua</i> Erichson,1837 | 14 | 23 | 139 | 23 | 6 | 205 |
| <i>Bolitochara pulchra</i> (Gravenhorst,1806) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Autalia impressa</i> (Olivier,1795) | 4 | 12 | 116 | 16 | 3 | 151 |
| <i>Autalia longicornis</i> Scheerpeltz,1947 | 1 | 2 | 22 | 7 | 1 | 33 |
| <i>Autalia puncticollis</i> Sharp,1864 | 0 | 0 | 0 | 2 | 1 | 3 |
| <i>Autalia rivularis</i> (Gravenhorst,1802) | 23 | 15 | 115 | 26 | 7 | 186 |
| <i>Cordalia obscura</i> (Gravenhorst,1802) | 0 | 3 | 1 | 2 | 0 | 6 |
| <i>Falagria caesa</i> Erichson,1837 | 1 | 4 | 16 | 1 | 0 | 22 |
| <i>Falagria sulcatula</i> (Gravenhorst,1806) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Falagrioma thoracica</i> (Stephens,1832) | 0 | 1 | 2 | 0 | 0 | 3 |
| <i>Myrmecopora sulcata</i> (Kiesenwetter,1850) | 3 | 2 | 0 | 0 | 0 | 5 |
| <i>Myrmecopora uvida</i> (Erichson,1840) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Thinonoma atra</i> (Gravenhorst,1806) | 31 | 3 | 38 | 8 | 1 | 81 |
| <i>Tachyusa coarctata</i> (Erichson,1837) | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Tachyusa constricta</i> (Erichson,1837) | 3 | 11 | 1 | 5 | 0 | 20 |
| <i>Ischnopoda leucopus</i> (Marsham,1802) | 4 | 11 | 5 | 4 | 0 | 24 |
| <i>Ischnopoda umbratica</i> (Erichson,1837) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Gnypeta caerulea</i> (Sahlberg,C.R.,1831) | 0 | 1 | 0 | 4 | 4 | 9 |
| <i>Gnypeta carbonaria</i> (Mannerheim,1830) | 11 | 8 | 26 | 5 | 0 | 50 |
| <i>Gnypeta ripicola</i> (Kiesenwetter,1844) | 0 | 1 | 2 | 0 | 0 | 3 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| <i>Gnypeta rubrior</i> Tottenham,1939 | 4 | 1 | 8 | 0 | 0 | 13 |
| <i>Dasygnypeta velata</i> (Erichson,1837) | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Callicerus obscurus</i> Gravenhorst,1802 | 6 | 3 | 7 | 3 | 0 | 19 |
| <i>Callicerus rigidicornis</i> (Erichson,1839) | 1 | 1 | 15 | 3 | 1 | 21 |
| <i>Schistoglossa aubei</i> (Brisout,1860) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Schistoglossa curtipennis</i> (Sharp,1869) | 1 | 0 | 7 | 0 | 0 | 8 |
| <i>Schistoglossa gemina</i> (Erichson,1837) | 3 | 0 | 3 | 2 | 0 | 8 |
| <i>Boreophilia eremita</i> (Rye,1866) | 4 | 8 | 27 | 8 | 3 | 50 |
| <i>Dacrila fallax</i> (Kraatz,1856) | 6 | 0 | 3 | 0 | 0 | 9 |
| <i>Hydrosmeeta delicatula</i> (Sharp,1869) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Hydrosmeeta eximia</i> (Sharp,1869) | 1 | 1 | 0 | 1 | 0 | 3 |
| <i>Hydrosmeeta fragilis</i> (Kraatz,1854) | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Hydrosmeeta subtilissima</i> (Kraatz,1854) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Aloconota cambrica</i> (Wollaston,1855) | 0 | 3 | 0 | 2 | 0 | 5 |
| <i>Aloconota currax</i> (Kraatz,1856) | 2 | 11 | 3 | 1 | 2 | 19 |
| <i>Aloconota gregaria</i> (Erichson,1839) | 83 | 50 | 139 | 39 | 7 | 318 |
| <i>Aloconota insecta</i> (Thomson,C.G.,1856) | 2 | 3 | 13 | 3 | 2 | 23 |
| <i>Aloconota sulcifrons</i> (Stephens,1832) | 5 | 1 | 22 | 1 | 2 | 31 |
| <i>Aloconota languida</i> (Erichson,1837) | 3 | 0 | 0 | 1 | 1 | 5 |
| <i>Pycnota paradoxa</i> (Mulsant & Rey,1861) | 3 | 0 | 0 | 1 | 0 | 4 |
| <i>Amischa analis</i> (Gravenhorst,1802) | 39 | 59 | 194 | 34 | 10 | 336 |
| <i>Amischa bifoveolata</i> (Mannerheim,1830) | 21 | 11 | 18 | 6 | 3 | 59 |
| <i>Amischa decipiens</i> (Sharp,1869) | 20 | 6 | 38 | 7 | 2 | 73 |
| <i>Amischa forcipata</i> Mulsant & Rey,1873 | 4 | 0 | 5 | 2 | 0 | 11 |
| <i>Amischa nigrofusca</i> (Stephens,1829) | 3 | 1 | 11 | 2 | 2 | 19 |
| <i>Amidobia talpa</i> (Heer,1841) | 2 | 7 | 6 | 0 | 0 | 15 |
| <i>Nehemitropia lividipennis</i> (Mannerheim,1830) | 22 | 11 | 53 | 14 | 0 | 100 |
| <i>Notothecta confusa</i> (Markel,1844) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Notothecta flavipes</i> (Gravenhorst,1806) | 1 | 7 | 14 | 0 | 0 | 22 |
| <i>Lyprocorrhe anceps</i> (Erichson,1837) | 1 | 5 | 5 | 1 | 0 | 12 |
| <i>Neohilara subterranea</i> (Mulsant & Rey,1853) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Brundinia meridionalis</i> (Mulsant & Rey,1853) | 10 | 0 | 0 | 0 | 0 | 10 |
| <i>Dochmonota clancula</i> (Erichson,1837) | 2 | 0 | 12 | 6 | 0 | 20 |
| <i>Geostiba circellaris</i> (Gravenhorst,1806) | 25 | 27 | 221 | 53 | 11 | 337 |
| <i>Ousipalia caesula</i> (Erichson,1839) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Dinaraea aequata</i> (Erichson,1837) | 16 | 12 | 59 | 13 | 3 | 103 |
| <i>Dinaraea angustula</i> (Gyllenhal,1810) | 13 | 4 | 31 | 3 | 1 | 52 |
| <i>Dinaraea linearis</i> (Gravenhorst,1802) | 0 | 1 | 3 | 0 | 0 | 4 |
| <i>Dadobia immersa</i> (Erichson,1837) | 2 | 1 | 4 | 4 | 0 | 11 |
| <i>Plataraea brunnea</i> (Fabricius,1798) | 8 | 5 | 26 | 2 | 1 | 42 |
| <i>Liogluta granigera</i> (Kiesenwetter,1850) | 1 | 8 | 11 | 2 | 0 | 22 |
| <i>Liogluta longiuscula</i> (Gravenhorst,1802) | 9 | 4 | 9 | 2 | 1 | 25 |
| <i>Liogluta alpestris</i> (Heer,1839) | 4 | 46 | 9 | 1 | 1 | 61 |
| <i>Liogluta microptera</i> Thomson,C.G.,1867 | 6 | 4 | 56 | 10 | 3 | 79 |
| <i>Liogluta pagana</i> (Erichson,1839) | 3 | 1 | 0 | 3 | 0 | 7 |
| <i>Philhygra arctica</i> (Thomson,C.G.,1856) | 0 | 6 | 1 | 1 | 0 | 8 |
| <i>Philhygra britteni</i> Joy,1913 | 10 | 1 | 5 | 0 | 0 | 16 |
| <i>Philhygra debilis</i> (Erichson,1837) | 8 | 0 | 8 | 0 | 1 | 17 |
| <i>Philhygra deformis</i> (Kraatz,1856) | 1 | 0 | 0 | 0 | 1 | 2 |
| <i>Philhygra elongatula</i> (Gravenhorst,1802) | 30 | 31 | 39 | 23 | 9 | 132 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|-----|----|------|
| <i>Philhygra fallaciosa</i> (Sharp,1869) | 1 | 2 | 1 | 0 | 0 | 4 |
| <i>Philhygra gyllenhali</i> (Thomson,C.G.,1856) | 7 | 2 | 10 | 2 | 0 | 21 |
| <i>Philhygra hygrobia</i> (Thomson,C.G.,1856) | 3 | 1 | 4 | 2 | 0 | 10 |
| <i>Philhygra hygrotopora</i> (Kraatz,1856) | 6 | 6 | 12 | 4 | 4 | 32 |
| <i>Philhygra luridipennis</i> (Mannerheim,1830) | 6 | 3 | 8 | 5 | 0 | 22 |
| <i>Philhygra malleus</i> (Joy,1913) | 27 | 3 | 30 | 5 | 2 | 67 |
| <i>Philhygra melanocera</i> (Thomson,C.G.,1856) | 10 | 4 | 44 | 4 | 0 | 62 |
| <i>Philhygra obtusangula</i> (Joy,1913) | 1 | 4 | 13 | 2 | 3 | 23 |
| <i>Philhygra palustris</i> (Kiesenwetter,1844) | 10 | 6 | 8 | 2 | 0 | 26 |
| <i>Philhygra terminalis</i> (Gravenhorst,1806) | 4 | 0 | 0 | 0 | 0 | 4 |
| <i>Philhygra volans</i> (Scriba,1859) | 5 | 0 | 12 | 1 | 1 | 19 |
| <i>Parameotica difficilis</i> (Brisout,1860) | 17 | 0 | 0 | 0 | 0 | 17 |
| <i>Dilacra luteipes</i> (Erichson,1837) | 5 | 1 | 3 | 4 | 0 | 13 |
| <i>Dilacra vilis</i> (Erichson,1837) | 1 | 0 | 4 | 6 | 0 | 11 |
| <i>Enalodroma hepatica</i> (Erichson,1839) | 1 | 2 | 5 | 6 | 2 | 16 |
| <i>Bessobia excellens</i> (Kraatz,1856) | 2 | 5 | 2 | 6 | 0 | 15 |
| <i>Bessobia fungivora</i> (Thomson,C.G.,1867) | 4 | 4 | 3 | 5 | 0 | 16 |
| <i>Bessobia monticola</i> (Thomson,C.G.,1852) | 1 | 2 | 6 | 3 | 0 | 12 |
| <i>Bessobia occulta</i> (Erichson,1837) | 0 | 3 | 3 | 0 | 1 | 7 |
| <i>Atheta divisa</i> (Markel,1844) | 1 | 0 | 0 | 1 | 0 | 2 |
| <i>Atheta euryptera</i> (Stephens,1832) | 2 | 2 | 2 | 1 | 0 | 7 |
| <i>Atheta harwoodi</i> (Williams,1930) | 6 | 4 | 77 | 11 | 5 | 103 |
| <i>Atheta liturata</i> (Stephens,1832) | 0 | 5 | 5 | 14 | 6 | 30 |
| <i>Atheta vaga</i> (Heer,1839) | 4 | 12 | 52 | 12 | 1 | 81 |
| <i>Atheta nigritula</i> (Gravenhorst,1802) | 2 | 0 | 5 | 0 | 0 | 7 |
| <i>Anopleta corvina</i> (Thomson,C.G.,1856) | 0 | 4 | 7 | 0 | 2 | 13 |
| <i>Anopleta puberula</i> (Sharp,1869) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Microdota amicula</i> (Stephens,1832) | 32 | 20 | 95 | 30 | 9 | 186 |
| <i>Microdota atricolor</i> (Sharp,1869) | 5 | 4 | 19 | 4 | 2 | 34 |
| <i>Microdota benickiella</i> (Brundin,1948) | 0 | 0 | 1 | 2 | 0 | 3 |
| <i>Microdota boreella</i> (Brundin,1948) | 0 | 0 | 2 | 1 | 0 | 3 |
| <i>Microdota excelsa</i> (Bernhauer,1911) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Microdota indubia</i> (Sharp,1869) | 10 | 6 | 39 | 15 | 3 | 73 |
| <i>Microdota liliputana</i> (Brisout,1860) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Microdota minuscula</i> (Brisout,1860) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Microdota spatuloides</i> (Benick,1939) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Microdota subtilis</i> (Scriba,1866) | 0 | 0 | 2 | 2 | 0 | 4 |
| <i>Cadaverota cadaverina</i> (Brisout,1860) | 0 | 9 | 18 | 0 | 0 | 27 |
| <i>Alaobia gagatina</i> (Baudi,1848) | 3 | 12 | 16 | 1 | 1 | 33 |
| <i>Alaobia hybrida</i> (Sharp,1869) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Alaobia pallidicornis</i> (Thomson,C.G.,1856) | 3 | 4 | 25 | 9 | 1 | 42 |
| <i>Alaobia sodalis</i> (Erichson,1837) | 1 | 12 | 16 | 6 | 3 | 38 |
| <i>Alaobia taxiceroides</i> Munster,1932 | 0 | 1 | 5 | 0 | 0 | 6 |
| <i>Oreostiba tibialis</i> (Heer,1839) | 0 | 0 | 0 | 6 | 3 | 9 |
| <i>Alaobia trinotata</i> (Kraatz,1856) | 36 | 28 | 132 | 23 | 3 | 222 |
| <i>Coprothassa melanaria</i> (Mannerheim,1830) | 4 | 1 | 0 | 0 | 0 | 5 |
| <i>Mocyta amplicollis</i> (Mulsant & Rey,1873) | 13 | 5 | 23 | 3 | 1 | 45 |
| <i>Mocyta clientula</i> (Erichson,1839) | 3 | 4 | 5 | 3 | 0 | 15 |
| <i>Mocyta fungi</i> (Gravenhorst,1806) | 86 | 91 | 681 | 114 | 29 | 1001 |
| <i>Mocyta gilvicollis</i> Scheerpeltz | 0 | 2 | 1 | 0 | 0 | 3 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Mocyta orbata</i> (Erichson,1837) | 4 | 1 | 16 | 2 | 1 | 24 |
| <i>Mocyta orphana</i> (Erichson,1837) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Acrotona aterrima</i> (Gravenhorst,1802) | 30 | 13 | 125 | 17 | 8 | 193 |
| <i>Acrotona benicki</i> (Allen,1940) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Acrotona exigua</i> (Erichson,1837) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Acrotona muscorum</i> (Brisout,1860) | 11 | 9 | 31 | 16 | 5 | 72 |
| <i>Acrotona obfusca</i> (Gravenhorst,1802) | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Acrotona parvula</i> (Mannerheim,1830) | 11 | 10 | 42 | 13 | 9 | 85 |
| <i>Acrotona pygmaea</i> (Gravenhorst,1802) | 2 | 0 | 31 | 9 | 0 | 42 |
| <i>Rhagocneme subsinuata</i> (Erichson,1839) | 1 | 2 | 10 | 1 | 0 | 14 |
| <i>Datomicra canescens</i> (Sharp,1869) | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Datomicra celata</i> (Erichson,1837) | 38 | 28 | 251 | 28 | 14 | 359 |
| <i>Datomicra dadopora</i> (Thomson,C.G.,1867) | 4 | 4 | 49 | 10 | 2 | 69 |
| <i>Datomicra nigra</i> (Kraatz,1856) | 46 | 16 | 131 | 38 | 9 | 240 |
| <i>Datomicra sordidula</i> (Erichson,1837) | 7 | 6 | 29 | 6 | 3 | 51 |
| <i>Datomicra zosteræ</i> (Thomson,C.G.,1856) | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Atheta aquatica</i> (Thomson,C.G.,1852) | 3 | 7 | 65 | 9 | 2 | 86 |
| <i>Atheta aquatilis</i> (Thomson,C.G.,1867) | 2 | 9 | 47 | 1 | 3 | 62 |
| <i>Atheta brunneipennis</i> (Thomson,C.G.,1852) | 2 | 5 | 18 | 5 | 1 | 31 |
| <i>Atheta castanoptera</i> (Mannerheim,1830) | 3 | 25 | 128 | 23 | 4 | 183 |
| <i>Atheta graminicola</i> (Gravenhorst,1806) | 33 | 29 | 199 | 46 | 10 | 317 |
| <i>Atheta hypnorum</i> (Kiesenwetter,1850) | 1 | 15 | 15 | 3 | 0 | 34 |
| <i>Atheta incognita</i> (Sharp,1869) | 0 | 5 | 2 | 0 | 0 | 7 |
| <i>Atheta aeneicollis</i> (Sharp,1869) | 35 | 35 | 99 | 24 | 4 | 197 |
| <i>Atheta triangulum</i> (Kraatz,1856) | 18 | 5 | 24 | 10 | 3 | 60 |
| <i>Atheta xanthopus</i> (Thomson,C.G.,1856) | 4 | 4 | 30 | 4 | 2 | 44 |
| <i>Atheta basicornis</i> (Mulsant & Rey,1852) | 0 | 0 | 1 | 2 | 1 | 4 |
| <i>Dalotia coriaria</i> (Kraatz,1856) | 7 | 6 | 48 | 10 | 0 | 71 |
| <i>Atheta crassicornis</i> (Fabricius,1793) | 35 | 54 | 238 | 50 | 18 | 395 |
| <i>Atheta diversa</i> (Sharp,1869) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Atheta fungicola</i> (Thomson,C.G.,1852) | 0 | 4 | 6 | 1 | 0 | 11 |
| <i>Atheta intermedia</i> (Thomson,C.G.,1852) | 2 | 0 | 3 | 1 | 0 | 6 |
| <i>Mycetota laticollis</i> (Stephens,1832) | 41 | 13 | 199 | 38 | 4 | 295 |
| <i>Atheta nidicola</i> (Johansen,1914) | 0 | 1 | 2 | 2 | 0 | 5 |
| <i>Atheta oblita</i> (Erichson,1839) | 1 | 2 | 2 | 0 | 0 | 5 |
| <i>Atheta paracrassicornis</i> Brundin,1954 | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Atheta pilicornis</i> (Thomson,C.G.,1852) | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Atheta ravilla</i> (Erichson,1839) | 12 | 8 | 42 | 8 | 2 | 72 |
| <i>Atheta britanniae</i> (Bernhauer & Scheerpeltz,1926) | 3 | 6 | 19 | 4 | 2 | 34 |
| <i>Atheta strandiella</i> (Brundin,1954) | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Dimetrota atramentaria</i> (Gyllenhal,1810) | 23 | 37 | 168 | 27 | 19 | 274 |
| <i>Dimetrota cauta</i> (Erichson,1837) | 4 | 0 | 1 | 1 | 0 | 6 |
| <i>Dimetrota cinnamoptera</i> (Thomson,C.G.,1856) | 0 | 12 | 29 | 3 | 6 | 50 |
| <i>Dimetrota ischnocera</i> (Thomson,C.G.,1870) | 2 | 6 | 26 | 3 | 1 | 38 |
| <i>Dimetrota laevana</i> (Mulsant & Rey,1852) | 0 | 5 | 10 | 5 | 2 | 22 |
| <i>Badura macrocera</i> (Thomson,C.G.,1856) | 4 | 2 | 15 | 6 | 2 | 29 |
| <i>Dimetrota marcida</i> (Erichson,1837) | 0 | 7 | 79 | 4 | 1 | 91 |
| <i>Dimetrota nigripes</i> (Thomson,C.G.,1856) | 13 | 7 | 48 | 2 | 6 | 76 |
| <i>Dimetrota aeneipennis</i> (Thomson,C.G.,1856) | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Dimetrota setigera</i> (Sharp,1869) | 4 | 3 | 28 | 2 | 1 | 38 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|---|-----|
| <i>Chaetida longicornis</i> (Gravenhorst,1802) | 31 | 24 | 142 | 31 | 9 | 237 |
| <i>Brundinia marina</i> (Mulsant & Rey,1853) | 19 | 0 | 0 | 0 | 0 | 19 |
| <i>Thinobaena vestita</i> (Gravenhorst,1806) | 34 | 13 | 0 | 0 | 0 | 47 |
| <i>Alevonota rufotestacea</i> (Kraatz,1856) | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Halobrecta algae</i> (Hardy,1851) | 3 | 0 | 0 | 0 | 0 | 3 |
| <i>Halobrecta flavipes</i> Thomson,C.G.,1861 | 6 | 2 | 0 | 0 | 0 | 8 |
| <i>Alianta incana</i> (Erichson,1837) | 8 | 1 | 32 | 5 | 2 | 48 |
| <i>Pachnida nigella</i> (Erichson,1837) | 21 | 1 | 53 | 6 | 1 | 82 |
| <i>Thamiaraea cinnamomea</i> (Gravenhorst,1802) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Thamiaraea hospita</i> (Markel,1844) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Drusilla canaliculata</i> (Fabricius,1787) | 33 | 43 | 67 | 10 | 5 | 158 |
| <i>Pella humeralis</i> (Gravenhorst,1802) | 2 | 3 | 5 | 0 | 1 | 11 |
| <i>Lomechusa emarginata</i> (Paykull,1789) | 0 | 0 | 6 | 0 | 0 | 6 |
| <i>Dinarda dentata</i> (Gravenhorst,1806) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Dinarda maerkeli</i> Kiesenwetter,1843 | 1 | 7 | 1 | 0 | 0 | 9 |
| <i>Trichiusa immigrata</i> Lohse,1984 | 8 | 0 | 35 | 0 | 1 | 44 |
| <i>Phloeopora corticalis</i> (Gravenhorst,1802) | 1 | 2 | 7 | 2 | 1 | 13 |
| <i>Phloeopora testacea</i> (Mannerheim,1830) | 3 | 10 | 37 | 7 | 0 | 57 |
| <i>Ilyobates nigricollis</i> (Paykull,1800) | 1 | 0 | 1 | 3 | 0 | 5 |
| <i>Ilyobates propinquus</i> (Aube,1850) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Ilyobates bennetti</i> Donisthorpe,1914 | 4 | 0 | 2 | 2 | 0 | 8 |
| <i>Calodera aethiops</i> (Gravenhorst,1802) | 23 | 5 | 16 | 6 | 1 | 51 |
| <i>Calodera nigrita</i> Mannerheim,1830 | 2 | 1 | 1 | 1 | 1 | 6 |
| <i>Calodera protensa</i> Mannerheim,1830 | 0 | 0 | 0 | 2 | 0 | 2 |
| <i>Calodera riparia</i> Erichson,1837 | 4 | 0 | 10 | 2 | 0 | 16 |
| <i>Tetralaucopora longitarsis</i> (Erichson,1837) | 24 | 17 | 25 | 11 | 3 | 80 |
| <i>Tetralaucopora rubicunda</i> (Erichson,1837) | 1 | 1 | 0 | 0 | 1 | 3 |
| <i>Amarochara umbrosa</i> (Erichson,1837) | 0 | 2 | 1 | 0 | 0 | 3 |
| <i>Ocalea badia</i> Erichson,1837 | 1 | 3 | 1 | 1 | 0 | 6 |
| <i>Ocalea latipennis</i> Sharp,1870 | 1 | 1 | 2 | 0 | 0 | 4 |
| <i>Ocalea picata</i> (Stephens,1832) | 4 | 19 | 45 | 11 | 6 | 85 |
| <i>Meotica anglica</i> Benick in Muona,1991 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Meotica filiformis</i> (Motschulsky,1860) | 4 | 0 | 10 | 2 | 0 | 16 |
| <i>Meotica exilis</i> (Knoch in Gravenhorst,1806) | 4 | 0 | 0 | 1 | 1 | 6 |
| <i>Meotica exillima</i> Sharp,1915 | 1 | 1 | 3 | 1 | 0 | 6 |
| <i>Ocyusa picina</i> (Aube,1850) | 1 | 1 | 7 | 1 | 0 | 10 |
| <i>Cousya longitarsis</i> (Thomson,C.G.,1867) | 0 | 11 | 0 | 0 | 0 | 11 |
| <i>Ocyusa maura</i> (Erichson,1837) | 9 | 0 | 25 | 5 | 0 | 39 |
| <i>Cousya nitidiventris</i> (Fagel,1958) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Mniusa incrassata</i> (Mulsant & Rey,1852) | 6 | 44 | 29 | 10 | 4 | 93 |
| <i>Oxypoda alternans</i> (Gravenhorst,1802) | 8 | 27 | 125 | 26 | 3 | 189 |
| <i>Oxypoda flavicornis</i> Kraatz,1856 | 3 | 0 | 4 | 1 | 0 | 8 |
| <i>Oxypoda annularis</i> Mannerheim,1830 | 6 | 25 | 22 | 6 | 0 | 59 |
| <i>Oxypoda brachyptera</i> (Stephens,1832) | 15 | 0 | 4 | 2 | 1 | 22 |
| <i>Oxypoda elongatula</i> Aube,1850 | 30 | 23 | 241 | 35 | 8 | 337 |
| <i>Oxypoda exoleta</i> Erichson,1839 | 8 | 1 | 3 | 0 | 0 | 12 |
| <i>Oxypoda formiceticola</i> Markel,1841 | 1 | 5 | 9 | 0 | 0 | 15 |
| <i>Oxypoda haemorrhoea</i> (Mannerheim,1830) | 32 | 19 | 114 | 16 | 7 | 188 |
| <i>Oxypoda induta</i> Mulsant & Rey,1861 | 1 | 2 | 1 | 0 | 0 | 4 |
| <i>Oxypoda islandica</i> Kraatz,1857 | 0 | 0 | 0 | 2 | 0 | 2 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Oxypoda lentula</i> Erichson,1837 | 20 | 1 | 5 | 0 | 0 | 26 |
| <i>Oxypoda acuminata</i> (Stephens,1832) | 3 | 7 | 22 | 3 | 3 | 38 |
| <i>Oxypoda longipes</i> Mulsant & Rey,1861 | 2 | 0 | 1 | 0 | 0 | 3 |
| <i>Oxypoda nigricornis</i> Motschulsky,1860 | 0 | 0 | 3 | 1 | 0 | 4 |
| <i>Oxypoda opaca</i> (Gravenhorst,1802) | 23 | 15 | 72 | 16 | 2 | 128 |
| <i>Oxypoda procerula</i> Mannerheim,1830 | 4 | 18 | 28 | 2 | 1 | 53 |
| <i>Oxypoda carbonaria</i> (Heer,1841) | 2 | 5 | 19 | 1 | 0 | 27 |
| <i>Oxypoda spectabilis</i> Markel,1844 | 1 | 6 | 2 | 2 | 0 | 11 |
| <i>Oxypoda brevicornis</i> (Stephens,1832) | 18 | 13 | 41 | 4 | 1 | 77 |
| <i>Ischnoglossa prolixa</i> (Gravenhorst,1802) | 1 | 0 | 11 | 0 | 0 | 12 |
| <i>Dexiogyia corticina</i> (Erichson,1837) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Thiasophila angulata</i> (Erichson,1837) | 3 | 8 | 9 | 0 | 0 | 20 |
| <i>Crataraea suturalis</i> (Mannerheim,1830) | 1 | 1 | 5 | 2 | 1 | 10 |
| <i>Haploglossa gentilis</i> (Markel,1844) | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Haploglossa nidicola</i> (Fairmaire,1852) | 5 | 1 | 2 | 2 | 1 | 11 |
| <i>Haploglossa villosula</i> (Stephens,1832) | 2 | 7 | 1 | 3 | 1 | 14 |
| <i>Tinotus morion</i> (Gravenhorst,1802) | 10 | 13 | 21 | 9 | 2 | 55 |
| <i>Aleochara bilineata</i> Gyllenhal,1810 | 7 | 4 | 5 | 0 | 0 | 16 |
| <i>Aleochara binotata</i> Kraatz,1858 | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Aleochara bipustulata</i> (Linnaeus,1761) | 21 | 9 | 28 | 5 | 1 | 64 |
| <i>Aleochara brevipennis</i> Gravenhorst,1806 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Aleochara cuniculorum</i> Kraatz,1858 | 5 | 0 | 1 | 2 | 1 | 9 |
| <i>Aleochara curtula</i> (Goeze,1777) | 22 | 15 | 46 | 18 | 11 | 112 |
| <i>Aleochara discipennis</i> Mulsant & Rey,1853 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Aleochara kamila</i> Likovsky,1984 | 0 | 1 | 3 | 1 | 0 | 5 |
| <i>Aleochara fumata</i> Gravenhorst,1802 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Aleochara funebris</i> Wollaston,1864 | 1 | 2 | 24 | 0 | 2 | 29 |
| <i>Aleochara grisea</i> Kraatz,1856 | 12 | 3 | 0 | 0 | 0 | 15 |
| <i>Aleochara inconspicua</i> Aube,1850 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Aleochara intricata</i> Mannerheim,1830 | 5 | 4 | 2 | 2 | 0 | 13 |
| <i>Aleochara lanuginosa</i> Gravenhorst,1802 | 24 | 37 | 108 | 29 | 17 | 215 |
| <i>Aleochara lygaea</i> Kraatz,1862 | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Aleochara moerens</i> Gyllenhal,1827 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Aleochara moesta</i> Gravenhorst,1802 | 1 | 1 | 1 | 2 | 0 | 5 |
| <i>Aleochara obscurella</i> Gravenhorst,1806 | 12 | 9 | 0 | 0 | 0 | 21 |
| <i>Aleochara punctatella</i> Motschulsky,1858 | 5 | 3 | 0 | 0 | 0 | 8 |
| <i>Aleochara ruficornis</i> Gravenhorst,1802 | 3 | 9 | 15 | 3 | 1 | 31 |
| <i>Aleochara spadicea</i> (Erichson,1837) | 8 | 3 | 1 | 5 | 0 | 17 |
| <i>Aleochara sparsa</i> Heer,1839 | 8 | 10 | 7 | 2 | 3 | 30 |
| <i>Aleochara verna</i> Say,1836 | 5 | 2 | 3 | 3 | 0 | 13 |
| <i>Aleochara villosa</i> Mannerheim,1830 | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Scaphidium quadrimaculatum</i> Olivier, 1790 | 1 | 16 | 35 | 2 | 0 | 54 |
| <i>Scaphisoma agaricinum</i> (Linnaeus, 1758) | 3 | 12 | 46 | 13 | 2 | 76 |
| <i>Scaphisoma boleti</i> (Panzer, 1793) | 8 | 12 | 10 | 6 | 0 | 36 |
| <i>Siagonium quadricorne</i> Kirby, 1815 | 3 | 6 | 25 | 6 | 1 | 41 |
| <i>Deleaster dichrous</i> (Gravenhorst, 1802) | 1 | 6 | 3 | 5 | 5 | 20 |
| <i>Syntomium aeneum</i> (Müller, P.W.J., 1821) | 6 | 24 | 60 | 19 | 1 | 110 |
| <i>Coprophilus striatulus</i> (Fabricius, 1792) | 5 | 6 | 7 | 3 | 0 | 21 |
| <i>Aploderus caelatus</i> (Gravenhorst, 1802) | 1 | 4 | 4 | 1 | 1 | 11 |
| <i>Bledius (Astycops) subterraneus</i> Erichson, 1839 | 0 | 14 | 3 | 7 | 2 | 26 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|-----|----|-----|
| <i>Bledius</i> (<i>Bledius</i>) <i>limicola</i> Tottenham, 1940 | 6 | 0 | 1 | 0 | 0 | 7 |
| <i>Bledius</i> (<i>Bledius</i>) <i>spectabilis</i> Kraatz, 1857 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Bledius</i> (<i>Bledius</i>) <i>tricornis</i> (Herbst, 1784) | 3 | 1 | 0 | 0 | 0 | 4 |
| <i>Bledius</i> (<i>Dicarenus</i>) <i>fergussoni</i> Joy, 1912 | 1 | 3 | 0 | 0 | 0 | 4 |
| <i>Bledius</i> (<i>Elbidus</i>) <i>bicornis</i> (Germar, 1822) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>annae</i> Sharp, 1911 | 2 | 4 | 2 | 11 | 1 | 20 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>defensus</i> Fauvel, 1872 | 0 | 2 | 1 | 9 | 2 | 14 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>dissimilis</i> Erichson, 1840 | 9 | 0 | 0 | 0 | 0 | 9 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>erraticus</i> Erichson, 1839 | 0 | 2 | 0 | 0 | 2 | 4 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>femorialis</i> (Gyllenhal, 1827) | 0 | 3 | 0 | 0 | 0 | 3 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>fuscipes</i> Rye, 1865 | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>gallicus</i> (Gravenhorst, 1806) | 4 | 5 | 5 | 1 | 1 | 16 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>longulus</i> Erichson, 1839 | 2 | 4 | 0 | 1 | 2 | 9 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>occidentalis</i> Bondroit, 1907 | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>opacus</i> (Block, 1799) | 8 | 4 | 4 | 0 | 0 | 16 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>pallipes</i> (Gravenhorst, 1806) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>praetermissus</i> Williams, 1929 | 6 | 0 | 1 | 0 | 0 | 7 |
| <i>Bledius</i> (<i>Hesperophilus</i>) <i>terebrans</i> (Schiödte, 1866) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Carpelimus</i> <i>bilineatus</i> Stephens, 1834 | 25 | 13 | 25 | 15 | 2 | 80 |
| <i>Carpelimus</i> <i>corticinus</i> (Gravenhorst, 1806) | 24 | 7 | 61 | 23 | 12 | 127 |
| <i>Carpelimus</i> <i>elongatulus</i> (Erichson, 1839) | 17 | 10 | 52 | 31 | 4 | 114 |
| <i>Carpelimus</i> <i>foveolatus</i> (Sahlberg, C.R., 1832) | 9 | 0 | 0 | 0 | 0 | 9 |
| <i>Carpelimus</i> <i>fuliginosus</i> (Gravenhorst, 1802) | 1 | 0 | 2 | 0 | 0 | 3 |
| <i>Carpelimus</i> <i>gracilis</i> (Mannerheim, 1830) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Carpelimus</i> <i>halophilus</i> (Kiesenwetter, 1844) | 3 | 0 | 0 | 1 | 0 | 4 |
| <i>Carpelimus</i> <i>impressus</i> (Boisduval & Lacordaire, 1835) | 1 | 1 | 1 | 1 | 0 | 4 |
| <i>Carpelimus</i> <i>manchuricus</i> subsp. <i>subtilicornis</i> (Roubal, 1946) | 1 | 1 | 1 | 7 | 0 | 10 |
| <i>Carpelimus</i> <i>obesus</i> (Kiesenwetter, 1844) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Carpelimus</i> <i>pusillus</i> (Gravenhorst, 1802) | 8 | 8 | 14 | 3 | 1 | 34 |
| <i>Carpelimus</i> <i>rivularis</i> (Motschulsky, 1860) | 25 | 8 | 48 | 18 | 7 | 106 |
| <i>Carpelimus</i> <i>zealandicus</i> (Sharp, 1900) | 6 | 2 | 10 | 3 | 0 | 21 |
| <i>Ochtheophilus</i> <i>andalusiacus</i> (Fagel, 1957) | 0 | 1 | 0 | 0 | 1 | 2 |
| <i>Ochtheophilus</i> <i>angustior</i> (Bernhauer, 1943) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Ochtheophilus</i> <i>aureus</i> (Fauvel, 1871) | 0 | 3 | 2 | 2 | 1 | 8 |
| <i>Ochtheophilus</i> <i>omalinus</i> (Erichson, 1840) | 0 | 14 | 1 | 4 | 3 | 22 |
| <i>Thinobius</i> <i>bicolor</i> Joy, 1911 | 0 | 2 | 0 | 1 | 0 | 3 |
| <i>Thinobius</i> <i>crinifer</i> Smetana, 1959 | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Thinodromus</i> <i>arcuatus</i> (Stephens, 1834) | 1 | 7 | 1 | 4 | 2 | 15 |
| <i>Anotylus</i> <i>complanatus</i> (Erichson, 1839) | 14 | 20 | 44 | 8 | 4 | 90 |
| <i>Anotylus</i> <i>fairmairei</i> (Pandellé, 1867) | 0 | 1 | 2 | 0 | 0 | 3 |
| <i>Anotylus</i> <i>insecatus</i> (Gravenhorst, 1806) | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Anotylus</i> <i>inustus</i> (Gravenhorst, 1806) | 9 | 11 | 61 | 1 | 1 | 83 |
| <i>Anotylus</i> <i>maritimus</i> Thomson, C.G., 1861 | 2 | 2 | 0 | 0 | 0 | 4 |
| <i>Anotylus</i> <i>mutator</i> (Lohse, 1963) | 2 | 4 | 21 | 5 | 0 | 32 |
| <i>Anotylus</i> <i>nitidulus</i> (Gravenhorst, 1802) | 1 | 8 | 7 | 4 | 1 | 21 |
| <i>Anotylus</i> <i>rugosus</i> (Fabricius, 1775) | 96 | 45 | 346 | 101 | 27 | 615 |
| <i>Anotylus</i> <i>saulcyi</i> (Pandellé, 1867) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Anotylus</i> <i>sculpturatus</i> (Gravenhorst, 1806) | 38 | 35 | 218 | 69 | 16 | 376 |
| <i>Anotylus</i> <i>tetracarınatus</i> (Block, 1799) | 37 | 52 | 196 | 56 | 30 | 371 |
| <i>Oxytelus</i> <i>fulvipes</i> Erichson, 1839 | 0 | 0 | 4 | 7 | 0 | 11 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| <i>Oxytelus laqueatus</i> (Marshall, 1802) | 15 | 25 | 111 | 38 | 14 | 203 |
| <i>Oxytelus migrator</i> Fauvel, 1904 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Oxytelus piceus</i> (Linnaeus, 1767) | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Oxytelus sculptus</i> Gravenhorst, 1806 | 5 | 4 | 19 | 3 | 2 | 33 |
| <i>Platystethus</i> (<i>Craetopycrus</i>) <i>cornutus</i> (Gravenhorst, 1802) | 28 | 13 | 41 | 15 | 3 | 100 |
| <i>Platystethus</i> (<i>Craetopycrus</i>) <i>nitens</i> (Sahlberg, C.R., 1832) | 10 | 2 | 5 | 1 | 0 | 18 |
| <i>Platystethus</i> (<i>Craetopycrus</i>) <i>nodifrons</i> (Mannerheim, 1830) | 24 | 3 | 1 | 2 | 4 | 34 |
| <i>Platystethus</i> (<i>Platystethus</i>) <i>arenarius</i> (Fourcroy, 1785) | 23 | 28 | 98 | 17 | 12 | 178 |
| <i>Oxyporus rufus</i> (Linnaeus, 1758) | 13 | 6 | 16 | 4 | 0 | 39 |
| <i>Stenus</i> (<i>Hemistenus</i>) <i>aceris</i> Stephens, 1833 | 1 | 5 | 6 | 1 | 2 | 15 |
| <i>Stenus</i> (<i>Hemistenus</i>) <i>fuscicornis</i> Erichson, 1840 | 2 | 2 | 16 | 2 | 1 | 23 |
| <i>Stenus</i> (<i>Hemistenus</i>) <i>geniculatus</i> Gravenhorst, 1806 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Stenus</i> (<i>Hemistenus</i>) <i>impressus</i> Germar, 1824 | 45 | 59 | 315 | 76 | 19 | 514 |
| <i>Stenus</i> (<i>Hemistenus</i>) <i>ochropus</i> Kiesenwetter, 1858 | 1 | 0 | 2 | 1 | 0 | 4 |
| <i>Stenus</i> (<i>Hemistenus</i>) <i>ossium</i> Stephens, 1833 | 10 | 6 | 16 | 5 | 0 | 37 |
| <i>Stenus</i> (<i>Hemistenus</i>) <i>pallipes</i> Gravenhorst, 1802 | 9 | 0 | 8 | 1 | 0 | 18 |
| <i>Stenus</i> (<i>Hemistenus</i>) <i>palustris</i> Erichson, 1839 | 9 | 2 | 6 | 5 | 0 | 22 |
| <i>Stenus</i> (<i>Hemistenus</i>) <i>subaeneus</i> Erichson, 1840 | 6 | 3 | 11 | 1 | 0 | 21 |
| <i>Stenus</i> (<i>Hypostenus</i>) <i>cicindeloides</i> (Schaller, 1783) | 26 | 9 | 47 | 20 | 0 | 102 |
| <i>Stenus</i> (<i>Hypostenus</i>) <i>fulvicornis</i> Stephens, 1833 | 25 | 15 | 114 | 30 | 4 | 188 |
| <i>Stenus</i> (<i>Hypostenus</i>) <i>kiesenwetteri</i> Rosenhauer, 1856 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Stenus</i> (<i>Hypostenus</i>) <i>latifrons</i> Erichson, 1839 | 13 | 7 | 34 | 8 | 1 | 63 |
| <i>Stenus</i> (<i>Hypostenus</i>) <i>oscillator</i> Rye, 1870 | 0 | 0 | 8 | 0 | 0 | 8 |
| <i>Stenus</i> (<i>Hypostenus</i>) <i>similis</i> (Herbst, 1784) | 61 | 58 | 201 | 49 | 6 | 375 |
| <i>Stenus</i> (<i>Hypostenus</i>) <i>solutus</i> Erichson, 1840 | 1 | 1 | 17 | 2 | 0 | 21 |
| <i>Stenus</i> (<i>Hypostenus</i>) <i>tarsalis</i> Ljungh, 1810 | 20 | 6 | 7 | 5 | 1 | 39 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>bifoveolatus</i> Gyllenhal, 1827 | 13 | 6 | 47 | 22 | 1 | 89 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>binotatus</i> Ljungh, 1804 | 2 | 10 | 22 | 1 | 0 | 35 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>brevipennis</i> Thomson, C.G., 1851 | 1 | 8 | 6 | 4 | 5 | 24 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>flavipes</i> Stephens, 1833 | 13 | 23 | 33 | 42 | 5 | 116 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>nitidiusculus</i> Stephens, 1833 | 14 | 17 | 83 | 42 | 6 | 162 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>niveus</i> Fauvel, 1865 | 2 | 0 | 1 | 0 | 0 | 3 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>pallitarsis</i> Stephens, 1833 | 39 | 7 | 41 | 14 | 2 | 103 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>picipennis</i> Erichson, 1840 | 5 | 4 | 25 | 6 | 1 | 41 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>picipes</i> Stephens, 1833 | 17 | 12 | 54 | 28 | 6 | 117 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>pubescens</i> Stephens, 1833 | 10 | 4 | 35 | 5 | 0 | 54 |
| <i>Stenus</i> (<i>Metatesnus</i>) <i>umbratilis</i> Casey, 1884 | 0 | 0 | 0 | 5 | 1 | 6 |
| <i>Stenus</i> (<i>Stenus</i>) <i>argus</i> Gravenhorst, 1806 | 10 | 2 | 0 | 3 | 0 | 15 |
| <i>Stenus</i> (<i>Stenus</i>) <i>asphaltinus</i> Erichson, 1840 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Stenus</i> (<i>Stenus</i>) <i>ater</i> Mannerheim, 1830 | 3 | 0 | 4 | 2 | 0 | 9 |
| <i>Stenus</i> (<i>Stenus</i>) <i>atratus</i> Erichson, 1839 | 4 | 5 | 9 | 5 | 0 | 23 |
| <i>Stenus</i> (<i>Stenus</i>) <i>biguttatus</i> (Linnaeus, 1758) | 14 | 11 | 13 | 9 | 5 | 52 |
| <i>Stenus</i> (<i>Stenus</i>) <i>bimaculatus</i> Gyllenhal, 1810 | 37 | 19 | 77 | 57 | 8 | 198 |
| <i>Stenus</i> (<i>Stenus</i>) <i>boops</i> Ljungh, 1810 | 38 | 13 | 79 | 29 | 3 | 162 |
| <i>Stenus</i> (<i>Stenus</i>) <i>canaliculatus</i> Gyllenhal, 1827 | 14 | 1 | 28 | 15 | 1 | 59 |
| <i>Stenus</i> (<i>Stenus</i>) <i>carbonarius</i> Gyllenhal, 1827 | 22 | 2 | 16 | 7 | 0 | 47 |
| <i>Stenus</i> (<i>Stenus</i>) <i>clavicornis</i> (Scopoli, 1763) | 67 | 34 | 240 | 44 | 10 | 395 |
| <i>Stenus</i> (<i>Stenus</i>) <i>comma</i> LeConte, 1863 | 5 | 4 | 8 | 5 | 0 | 22 |
| <i>Stenus</i> (<i>Stenus</i>) <i>europaeus</i> Puthz, 1966 | 22 | 3 | 13 | 9 | 1 | 48 |
| <i>Stenus</i> (<i>Stenus</i>) <i>fuscipes</i> Gravenhorst, 1802 | 0 | 0 | 1 | 1 | 0 | 2 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| Stenus (Stenus) guttula Müller, P.W.J., 1821 | 15 | 14 | 9 | 10 | 7 | 55 |
| Stenus (Stenus) guynemeri Jacquelin du Val, 1850 | 0 | 5 | 20 | 7 | 9 | 41 |
| Stenus (Stenus) incrassatus Erichson, 1839 | 1 | 0 | 1 | 0 | 0 | 2 |
| Stenus (Stenus) junco (Paykull, 1789) | 79 | 27 | 186 | 71 | 10 | 373 |
| Stenus (Stenus) lustrator Erichson, 1839 | 2 | 2 | 6 | 0 | 0 | 10 |
| Stenus (Stenus) melanarius Stephens, 1833 | 4 | 0 | 12 | 4 | 0 | 20 |
| Stenus (Stenus) melanopus (Marsham, 1802) | 4 | 0 | 2 | 0 | 0 | 6 |
| Stenus (Stenus) morio Gravenhorst, 1806 | 1 | 1 | 0 | 0 | 0 | 2 |
| Stenus (Stenus) nanus Stephens, 1833 | 6 | 11 | 54 | 9 | 4 | 84 |
| Stenus (Stenus) nitens Stephens, 1833 | 3 | 1 | 23 | 1 | 0 | 28 |
| Stenus (Stenus) providus Erichson, 1839 | 25 | 25 | 93 | 36 | 9 | 188 |
| Stenus (Stenus) pusillus Stephens, 1833 | 21 | 10 | 49 | 25 | 5 | 110 |
| Stenus (Tesnus) brunnipes Stephens, 1833 | 67 | 22 | 191 | 63 | 14 | 357 |
| Stenus (Tesnus) crassus Stephens, 1833 | 3 | 1 | 8 | 5 | 0 | 17 |
| Stenus (Tesnus) formicetorum Mannerheim, 1843 | 2 | 2 | 1 | 0 | 0 | 5 |
| Stenus (Tesnus) nigritulus Gyllenhal, 1827 | 9 | 3 | 7 | 1 | 1 | 21 |
| Dianous coerulescens (Gyllenhal, 1810) | 1 | 17 | 21 | 18 | 18 | 75 |
| Euaesthetus bipunctatus (Ljungh, 1804) | 2 | 1 | 7 | 14 | 2 | 26 |
| Euaesthetus laeviusculus Mannerheim, 1844 | 2 | 4 | 17 | 4 | 0 | 27 |
| Euaesthetus ruficapillus Lacordaire, 1835 | 16 | 4 | 26 | 6 | 3 | 55 |
| Pseudopsis sulcata Newman, 1834 | 1 | 3 | 4 | 3 | 0 | 11 |
| Achenium depressum (Gravenhorst, 1802) | 1 | 0 | 0 | 0 | 0 | 1 |
| Achenium humile (Nicolai, 1822) | 1 | 0 | 0 | 0 | 0 | 1 |
| Astenus (Astenognathus) procerus (Gravenhorst, 1806) | 0 | 0 | 1 | 0 | 0 | 1 |
| Astenus (Astenognathus) pulchellus (Heer, 1839) | 6 | 1 | 10 | 2 | 0 | 19 |
| Astenus (Astenus) lyonessius (Joy, 1908) | 14 | 1 | 5 | 3 | 0 | 23 |
| Lathrobium (Tetartopeus) quadratum (Paykull, 1789) | 16 | 3 | 24 | 5 | 0 | 48 |
| Lathrobium (Tetartopeus) terminatum Gravenhorst, 1802 | 6 | 6 | 71 | 8 | 3 | 94 |
| Lathrobium (Tetartopeus) zetterstedti Rye, 1872 | 0 | 1 | 3 | 1 | 2 | 7 |
| Lathrobium (Lathrobium) brunnipes (Fabricius, 1792) | 49 | 36 | 272 | 92 | 25 | 474 |
| Lathrobium (Lathrobium) elongatum (Linnaeus, 1767) | 9 | 4 | 13 | 9 | 3 | 38 |
| Lathrobium (Lathrobium) fovulum Stephens, 1833 | 1 | 0 | 2 | 5 | 0 | 8 |
| Lathrobium (Lathrobium) fulvipenne (Gravenhorst, 1806) | 25 | 21 | 75 | 19 | 3 | 143 |
| Lathrobium (Lathrobium) geminum Kraatz, 1857 | 13 | 8 | 58 | 9 | 5 | 93 |
| Lathrobium (Lathrobium) impressum Heer, 1841 | 7 | 1 | 11 | 13 | 0 | 32 |
| Lathrobium (Lathrobium) longulum Gravenhorst, 1802 | 7 | 2 | 10 | 1 | 1 | 21 |
| Lathrobium (Lathrobium) pallidipenne Hichhuth, 1851 | 1 | 4 | 1 | 1 | 0 | 7 |
| Lathrobium (Lathrobium) rufipenne Gyllenhal, 1813 | 0 | 0 | 1 | 0 | 0 | 1 |
| Lithocharis nigriceps Kraatz, 1859 | 15 | 2 | 40 | 7 | 1 | 65 |
| Lithocharis ochracea (Gravenhorst, 1802) | 11 | 2 | 20 | 5 | 2 | 40 |
| Lobrathium multipunctum (Gravenhorst, 1802) | 1 | 6 | 5 | 1 | 1 | 14 |
| Medon apicalis (Kraatz, 1857) | 1 | 0 | 0 | 0 | 0 | 1 |
| Medon brunneus (Erichson, 1839) | 0 | 1 | 2 | 0 | 0 | 3 |
| Medon castaneus (Gravenhorst, 1802) | 0 | 0 | 1 | 0 | 0 | 1 |
| Medon fuscus (Mannerheim, 1830) | 0 | 4 | 0 | 0 | 0 | 4 |
| Medon pocofer (Peyron, 1857) | 0 | 1 | 0 | 0 | 0 | 1 |
| Ochtheophilum fracticorne (Paykull, 1800) | 4 | 7 | 11 | 6 | 1 | 29 |
| Paederus fuscipes Curtis, 1826 | 0 | 0 | 1 | 0 | 0 | 1 |
| Paederus littoralis Gravenhorst, 1802 | 1 | 3 | 1 | 1 | 0 | 6 |
| Paederus riparius (Linnaeus, 1758) | 3 | 8 | 3 | 1 | 0 | 15 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Pseudomedon obscurellus</i> (Erichson, 1840) | 0 | 0 | 2 | 1 | 0 | 3 |
| <i>Pseudomedon obsoletus</i> (Nordmann, 1837) | 0 | 0 | 7 | 0 | 0 | 7 |
| <i>Rugilus erichsoni</i> (Fauvel, 1867) | 13 | 5 | 28 | 17 | 0 | 63 |
| <i>Rugilus orbiculatus</i> (Paykull, 1789) | 46 | 19 | 158 | 29 | 2 | 254 |
| <i>Rugilus rufipes</i> Germar, 1835 | 44 | 7 | 82 | 28 | 4 | 165 |
| <i>Rugilus subtilis</i> (Erichson, 1840) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Scopaeus sulcicollis</i> (Stephens, 1833) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Sunius bicolor</i> (Olivier, 1795) | 4 | 0 | 1 | 0 | 0 | 5 |
| <i>Sunius melanocephalus</i> (Fabricius, 1792) | 3 | 1 | 0 | 1 | 0 | 5 |
| <i>Sunius propinquus</i> (Brisout, 1867) | 15 | 2 | 16 | 0 | 0 | 33 |
| <i>Bisnius cephalotes</i> (Gravenhorst, 1802) | 12 | 8 | 57 | 14 | 3 | 94 |
| <i>Bisnius fimetarius</i> (Gravenhorst, 1802) | 37 | 22 | 272 | 36 | 8 | 375 |
| <i>Bisnius nigriventris</i> (Thomson, C.G., 1867) | 4 | 1 | 8 | 3 | 2 | 18 |
| <i>Bisnius parvus</i> (Sharp, 1874) | 4 | 0 | 13 | 2 | 0 | 19 |
| <i>Bisnius puella</i> (Nordmann, 1837) | 2 | 3 | 29 | 10 | 2 | 46 |
| <i>Bisnius sordidus</i> (Gravenhorst, 1802) | 22 | 5 | 67 | 11 | 0 | 105 |
| <i>Bisnius subuliformis</i> (Gravenhorst, 1802) | 1 | 0 | 3 | 2 | 0 | 6 |
| <i>Cafius fucicola</i> Curtis, 1830 | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Cafius xantholoma</i> (Gravenhorst, 1806) | 13 | 9 | 2 | 0 | 0 | 24 |
| <i>Erichsonius cinerascens</i> (Gravenhorst, 1802) | 2 | 4 | 10 | 0 | 0 | 16 |
| <i>Gabrius appendiculatus</i> Sharp, 1910 | 1 | 3 | 8 | 4 | 1 | 17 |
| <i>Gabrius astutoides</i> (Strand, A., 1946) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Gabrius bishopi</i> Sharp, 1910 | 12 | 0 | 4 | 4 | 1 | 21 |
| <i>Gabrius breviventer</i> (Sperk, 1835) | 22 | 17 | 93 | 35 | 3 | 170 |
| <i>Gabrius nigritulus</i> (Gravenhorst, 1802) | 6 | 7 | 14 | 4 | 1 | 32 |
| <i>Gabrius osseticus</i> (Kolenati, 1846) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Gabrius piliger</i> Mulsant & Rey, 1876 | 5 | 0 | 36 | 5 | 2 | 48 |
| <i>Gabrius splendidulus</i> (Gravenhorst, 1802) | 6 | 13 | 80 | 14 | 6 | 119 |
| <i>Gabrius trossulus</i> (Nordmann, 1837) | 17 | 6 | 29 | 26 | 1 | 79 |
| <i>Gabrius velox</i> Sharp, 1910 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Gabronthus thermarum</i> (Aubé, 1850) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Neobisnius lathrobioides</i> (Baudi, 1848) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Neobisnius procerulus</i> (Gravenhorst, 1806) | 1 | 0 | 2 | 0 | 0 | 3 |
| <i>Neobisnius prolixus</i> (Erichson, 1840) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Neobisnius villosulus</i> (Stephens, 1833) | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Philonthus addendus</i> Sharp, 1867 | 2 | 5 | 10 | 2 | 2 | 21 |
| <i>Philonthus albipes</i> (Gravenhorst, 1802) | 8 | 3 | 22 | 3 | 1 | 37 |
| <i>Philonthus atratus</i> (Gravenhorst, 1802) | 3 | 4 | 5 | 0 | 0 | 12 |
| <i>Philonthus carbonarius</i> (Gravenhorst, 1802) | 31 | 47 | 160 | 39 | 7 | 284 |
| <i>Philonthus cognatus</i> Stephens, 1832 | 33 | 34 | 132 | 55 | 8 | 262 |
| <i>Philonthus concinnus</i> (Gravenhorst, 1802) | 12 | 1 | 21 | 5 | 0 | 39 |
| <i>Philonthus corruscus</i> (Gravenhorst, 1802) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Philonthus corvinus</i> Erichson, 1839 | 3 | 3 | 0 | 0 | 0 | 6 |
| <i>Philonthus cruentatus</i> (Gmelin in Linnaeus, 1790) | 4 | 2 | 7 | 1 | 0 | 14 |
| <i>Philonthus debilis</i> (Gravenhorst, 1802) | 4 | 2 | 30 | 3 | 0 | 39 |
| <i>Philonthus decorus</i> (Gravenhorst, 1802) | 7 | 22 | 176 | 34 | 15 | 254 |
| <i>Philonthus discoideus</i> (Gravenhorst, 1802) | 4 | 5 | 10 | 2 | 0 | 21 |
| <i>Philonthus ebeninus</i> (Gravenhorst, 1802) | 0 | 4 | 5 | 2 | 0 | 11 |
| <i>Philonthus fumarius</i> (Gravenhorst, 1806) | 0 | 1 | 0 | 6 | 0 | 7 |
| <i>Philonthus intermedius</i> (Lacordaire, 1835) | 12 | 10 | 38 | 5 | 2 | 67 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| <i>Philonthus jurgans</i> Tottenham, 1937 | 2 | 2 | 6 | 2 | 0 | 12 |
| <i>Philonthus laminatus</i> (Creutzer, 1799) | 17 | 16 | 84 | 27 | 8 | 152 |
| <i>Philonthus longicornis</i> Stephens, 1832 | 11 | 5 | 10 | 5 | 1 | 32 |
| <i>Philonthus mannerheimi</i> Fauvel, 1869 | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Philonthus marginatus</i> (Müller, O.F., 1764) | 16 | 14 | 73 | 27 | 6 | 136 |
| <i>Philonthus micans</i> (Gravenhorst, 1802) | 19 | 14 | 7 | 6 | 2 | 48 |
| <i>Philonthus micantoides</i> Benick & Lohse, 1956 | 11 | 2 | 12 | 8 | 0 | 33 |
| <i>Philonthus nigrita</i> (Gravenhorst, 1806) | 2 | 9 | 4 | 4 | 0 | 19 |
| <i>Philonthus parvicornis</i> (Gravenhorst, 1802) | 0 | 0 | 3 | 2 | 0 | 5 |
| <i>Philonthus politus</i> (Linnaeus, 1758) | 14 | 5 | 43 | 9 | 3 | 74 |
| <i>Philonthus quisquiliarius</i> (Gyllenhal, 1810) | 26 | 9 | 34 | 14 | 6 | 89 |
| <i>Philonthus rectangulus</i> Sharp, 1874 | 4 | 5 | 23 | 7 | 2 | 41 |
| <i>Philonthus rotundicollis</i> (Ménétriés, 1832) | 2 | 10 | 23 | 5 | 0 | 40 |
| <i>Philonthus rubripennis</i> Stephens, 1832 | 0 | 20 | 0 | 2 | 0 | 22 |
| <i>Philonthus rufipes</i> (Stephens, 1832) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Philonthus sanguinolentus</i> (Gravenhorst, 1802) | 9 | 13 | 26 | 2 | 0 | 50 |
| <i>Philonthus spinipes</i> Sharp, 1874 | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Philonthus splendens</i> (Fabricius, 1792) | 16 | 10 | 37 | 11 | 11 | 85 |
| <i>Philonthus succicola</i> Thomson, C.G., 1860 | 17 | 9 | 90 | 19 | 8 | 143 |
| <i>Philonthus tenuicornis</i> Mulsant & Rey, 1853 | 11 | 7 | 74 | 11 | 2 | 105 |
| <i>Philonthus umbratilis</i> (Gravenhorst, 1802) | 8 | 3 | 28 | 2 | 0 | 41 |
| <i>Philonthus varians</i> (Paykull, 1789) | 50 | 35 | 158 | 49 | 19 | 311 |
| <i>Philonthus ventralis</i> (Gravenhorst, 1802) | 1 | 3 | 6 | 3 | 0 | 13 |
| <i>Remus sericeus</i> Holme, 1837 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Creophilus maxillosus</i> (Linnaeus, 1758) | 13 | 8 | 29 | 4 | 6 | 60 |
| <i>Dinothenarus pubescens</i> (De Geer, 1774) | 3 | 1 | 5 | 3 | 0 | 12 |
| <i>Ocypus</i> (Matidus) <i>brunnipes</i> (Fabricius, 1781) | 25 | 13 | 50 | 14 | 3 | 105 |
| <i>Ocypus</i> (Matidus) <i>nitens</i> (Schrank, 1781) | 1 | 1 | 4 | 0 | 0 | 6 |
| <i>Ocypus</i> (<i>Ocypus</i>) <i>olens</i> (Müller, O.F., 1764) | 33 | 8 | 132 | 28 | 1 | 202 |
| <i>Ocypus</i> (<i>Ocypus</i>) <i>ophthalmicus</i> (Scopoli, 1763) | 0 | 0 | 8 | 0 | 0 | 8 |
| <i>Ocypus</i> (<i>Pseudocypus</i>) <i>aeneocephalus</i> (De Geer, 1774) | 5 | 13 | 40 | 32 | 7 | 97 |
| <i>Ocypus</i> (<i>Pseudocypus</i>) <i>fortunatarum</i> (Wollaston, 1871) | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Ocypus</i> (<i>Pseudocypus</i>) <i>fuscatus</i> (Gravenhorst, 1802) | 2 | 2 | 7 | 1 | 0 | 12 |
| <i>Ontholestes murinus</i> (Linnaeus, 1758) | 3 | 0 | 14 | 1 | 0 | 18 |
| <i>Ontholestes tessellatus</i> (Geoffroy, 1785) | 1 | 2 | 20 | 7 | 1 | 31 |
| <i>Platydracus fulvipes</i> (Scopoli, 1763) | 1 | 0 | 1 | 0 | 1 | 3 |
| <i>Platydracus latebricola</i> (Gravenhorst, 1806) | 0 | 4 | 0 | 0 | 0 | 4 |
| <i>Platydracus stercorarius</i> (Olivier, 1795) | 8 | 6 | 20 | 5 | 6 | 45 |
| <i>Staphylinus caesareus</i> Cederhjelm, 1798 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Staphylinus dimidiaticornis</i> Gemminger, 1851 | 0 | 1 | 0 | 0 | 1 | 2 |
| <i>Staphylinus erythropterus</i> Linnaeus, 1758 | 2 | 11 | 0 | 1 | 3 | 17 |
| <i>Tasgius</i> (<i>Rayacheila</i>) <i>globulifer</i> (Geoffroy, 1785) | 7 | 7 | 21 | 6 | 2 | 43 |
| <i>Tasgius</i> (<i>Rayacheila</i>) <i>melanarius</i> (Heer, 1839) | 9 | 4 | 46 | 20 | 2 | 81 |
| <i>Tasgius</i> (<i>Rayacheila</i>) <i>morsitans</i> (Rossi, 1790) | 8 | 4 | 11 | 1 | 2 | 26 |
| <i>Tasgius</i> (<i>Rayacheila</i>) <i>winkleri</i> Bernhauer, 1906 | 3 | 3 | 3 | 0 | 0 | 9 |
| <i>Tasgius</i> (<i>Tasgius</i>) <i>ater</i> (Gravenhorst, 1802) | 24 | 2 | 22 | 2 | 0 | 50 |
| <i>Tasgius</i> (<i>Tasgius</i>) <i>pedator</i> (Gravenhorst, 1802) | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Heterothops binotatus</i> (Gravenhorst, 1802) | 5 | 2 | 0 | 2 | 0 | 9 |
| <i>Heterothops dissimilis</i> (Gravenhorst, 1802) | 6 | 3 | 11 | 1 | 0 | 21 |
| <i>Heterothops minutus</i> Wollaston, 1860 | 3 | 0 | 10 | 1 | 0 | 14 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| Heterothops niger Kraatz, 1868 | 3 | 1 | 2 | 1 | 0 | 7 |
| Heterothops praeius Erichson, 1839 | 1 | 1 | 1 | 2 | 0 | 5 |
| Quedius (Distichalius) cinctus (Paykull, 1790) | 9 | 3 | 20 | 10 | 0 | 42 |
| Quedius (Microsaurus) brevicornis (Thomson, C.G., 1860) | 4 | 0 | 0 | 1 | 0 | 5 |
| Quedius (Microsaurus) brevis Erichson, 1840 | 0 | 4 | 2 | 0 | 1 | 7 |
| Quedius (Microsaurus) cruentus (Olivier, 1795) | 10 | 12 | 59 | 15 | 3 | 99 |
| Quedius (Microsaurus) fulgidus (Fabricius, 1792) | 1 | 2 | 1 | 1 | 0 | 5 |
| Quedius (Microsaurus) invreae Gridelli, 1924 | 0 | 2 | 9 | 3 | 0 | 14 |
| Quedius (Microsaurus) lateralis (Gravenhorst, 1802) | 1 | 7 | 30 | 4 | 1 | 43 |
| Quedius (Microsaurus) longicornis Kraatz, 1857 | 3 | 3 | 4 | 1 | 0 | 11 |
| Quedius (Microsaurus) maurus (Sahlberg, C.R., 1830) | 3 | 9 | 17 | 3 | 0 | 32 |
| Quedius (Microsaurus) mesomelinus (Marsham, 1802) | 16 | 11 | 89 | 14 | 5 | 135 |
| Quedius (Microsaurus) nigrocaeruleus Fauvel, 1874 | 3 | 0 | 0 | 2 | 0 | 5 |
| Quedius (Microsaurus) puncticollis (Thomson, C.G., 1867) | 1 | 4 | 1 | 6 | 1 | 13 |
| Quedius (Microsaurus) scitus (Gravenhorst, 1806) | 7 | 6 | 24 | 7 | 2 | 46 |
| Quedius (Microsaurus) truncicola Fairmaire & Laboulbène, 1856 | 0 | 2 | 4 | 1 | 0 | 7 |
| Quedius (Microsaurus) xanthopus Erichson, 1839 | 0 | 9 | 3 | 0 | 1 | 13 |
| Quedius (Quedionuchus) plagiatus Mannerheim, 1843 | 0 | 5 | 8 | 11 | 1 | 25 |
| Quedius (Quedius) curtipennis Bernhauer, 1908 | 19 | 23 | 165 | 58 | 11 | 276 |
| Quedius (Quedius) fuliginosus (Gravenhorst, 1802) | 14 | 9 | 56 | 21 | 4 | 104 |
| Quedius (Quedius) levicollis (Brullé, 1832) | 12 | 6 | 47 | 13 | 4 | 82 |
| Quedius (Quedius) molochinus (Gravenhorst, 1806) | 15 | 24 | 83 | 25 | 12 | 159 |
| Quedius (Quedius) simplicifrons Fairmaire, 1861 | 16 | 0 | 0 | 0 | 0 | 16 |
| Quedius (Raphirus) auricomus Kiesenwetter, 1850 | 0 | 5 | 6 | 6 | 5 | 22 |
| Quedius (Raphirus) boopoides Munster, 1923 | 4 | 0 | 4 | 1 | 1 | 10 |
| Quedius (Raphirus) boops (Gravenhorst, 1802) | 14 | 19 | 43 | 13 | 7 | 96 |
| Quedius (Raphirus) fulvicollis (Stephens, 1833) | 0 | 5 | 5 | 4 | 1 | 15 |
| Quedius (Raphirus) fumatus (Stephens, 1833) | 2 | 11 | 41 | 12 | 1 | 67 |
| Quedius (Raphirus) humeralis Stephens, 1832 | 4 | 0 | 4 | 1 | 0 | 9 |
| Quedius (Raphirus) maurorufus (Gravenhorst, 1806) | 29 | 7 | 127 | 44 | 7 | 214 |
| Quedius (Raphirus) nemoralis Baudi, 1848 | 4 | 0 | 21 | 1 | 0 | 26 |
| Quedius (Raphirus) nigriceps Kraatz, 1857 | 1 | 14 | 29 | 3 | 0 | 47 |
| Quedius (Raphirus) nitipennis (Stephens, 1833) | 8 | 8 | 14 | 13 | 6 | 49 |
| Quedius (Raphirus) persimilis Mulsant & Rey, 1876 | 2 | 0 | 23 | 3 | 1 | 29 |
| Quedius (Raphirus) picipes (Mannerheim, 1830) | 6 | 9 | 16 | 6 | 1 | 38 |
| Quedius (Raphirus) riparius Kellner, 1843 | 1 | 1 | 1 | 0 | 0 | 3 |
| Quedius (Raphirus) schatzmayri Gridelli, 1922 | 4 | 5 | 19 | 6 | 0 | 34 |
| Quedius (Raphirus) scintillans (Gravenhorst, 1806) | 11 | 2 | 58 | 22 | 3 | 96 |
| Quedius (Raphirus) semiaeneus (Stephens, 1833) | 8 | 5 | 7 | 2 | 1 | 23 |
| Quedius (Raphirus) semiobscurus (Marsham, 1802) | 15 | 6 | 25 | 8 | 0 | 54 |
| Quedius (Raphirus) umbrinus Erichson, 1839 | 5 | 12 | 12 | 13 | 5 | 47 |
| Atrecus affinis (Paykull, 1789) | 5 | 19 | 92 | 18 | 5 | 139 |
| Othius angustus Stephens, 1833 | 4 | 5 | 33 | 7 | 7 | 56 |
| Othius laeviusculus Stephens, 1833 | 10 | 3 | 18 | 1 | 2 | 34 |
| Othius punctulatus (Goeze, 1777) | 13 | 14 | 118 | 41 | 9 | 195 |
| Othius subuliformis Stephens, 1833 | 17 | 29 | 101 | 19 | 12 | 178 |
| Gauropterus fulgidus (Fabricius, 1787) | 2 | 1 | 3 | 3 | 0 | 9 |
| Gyrohypnus angustatus Stephens, 1833 | 19 | 13 | 39 | 18 | 1 | 90 |
| Gyrohypnus atratus (Heer, 1839) | 0 | 3 | 0 | 0 | 0 | 3 |
| Gyrohypnus fracticornis (Müller, O.F., 1776) | 23 | 11 | 91 | 19 | 4 | 148 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| Gyrophypnus punctulatus (Paykull, 1789) | 1 | 1 | 47 | 8 | 1 | 58 |
| Leptacinus batychrus (Gyllenhal, 1827) | 0 | 4 | 14 | 0 | 0 | 18 |
| Leptacinus formicetorum Märkel, 1841 | 0 | 5 | 7 | 0 | 0 | 12 |
| Leptacinus intermedius Donisthorpe, 1936 | 1 | 0 | 1 | 2 | 0 | 4 |
| Leptacinus pusillus (Stephens, 1833) | 11 | 3 | 64 | 10 | 1 | 89 |
| Megalinus glabratus (Gravenhorst, 1802) | 34 | 12 | 25 | 10 | 0 | 81 |
| Nudobius lentus (Gravenhorst, 1806) | 2 | 9 | 12 | 7 | 1 | 31 |
| Phacophallus parumpunctatus (Gyllenhal, 1827) | 6 | 2 | 8 | 1 | 0 | 17 |
| Xantholinus (Acanthophallus) laevigatus Jacobsen, 1849 | 0 | 0 | 1 | 1 | 0 | 2 |
| Xantholinus (Purrolinus) elegans (Olivier, 1795) | 8 | 1 | 17 | 1 | 1 | 28 |
| Xantholinus (Purrolinus) tricolor (Fabricius, 1787) | 1 | 0 | 0 | 0 | 0 | 1 |
| Xantholinus (Xantholinus) gallicus Coiffait, 1956 | 0 | 3 | 0 | 0 | 1 | 4 |
| Xantholinus (Xantholinus) linearis (Olivier, 1795) | 85 | 41 | 281 | 56 | 17 | 480 |
| Xantholinus (Xantholinus) longiventris Heer, 1839 | 20 | 20 | 66 | 27 | 0 | 133 |

Lucanidae [4/3]

| | | | | | | |
|--|----|----|----|----|---|-----|
| Sinodendron cylindricum (Linnaeus, 1758) | 8 | 24 | 98 | 25 | 5 | 160 |
| Lucanus cervus (Linnaeus, 1758) | 1 | 2 | 3 | 3 | 0 | 9 |
| Dorcus parallelipedus (Linnaeus, 1758) | 22 | 3 | 36 | 7 | 0 | 68 |

Trogidae [3/2]

| | | | | | | |
|---------------------------------|---|---|----|---|---|----|
| Trox sabulosus (Linnaeus, 1758) | 0 | 0 | 0 | 1 | 0 | 1 |
| Trox scaber (Linnaeus, 1767) | 0 | 1 | 10 | 2 | 0 | 13 |

Bolboceratidae [1/1]

| | | | | | | |
|----------------------------------|---|---|---|---|---|---|
| Odonteus armiger (Scopoli, 1772) | 0 | 0 | 2 | 0 | 0 | 2 |
|----------------------------------|---|---|---|---|---|---|

Geotrupidae [7/6]

| | | | | | | |
|---|----|----|----|---|---|----|
| Typhaeus typhoeus (Linnaeus, 1758) | 13 | 11 | 3 | 1 | 1 | 29 |
| Anoplotrupes stercorosus (Hartmann in Scriba, 1791) | 0 | 5 | 7 | 0 | 1 | 13 |
| Geotrupes mutator (Marsham, 1802) | 2 | 0 | 0 | 0 | 0 | 2 |
| Geotrupes spiniger (Marsham, 1802) | 10 | 1 | 12 | 3 | 1 | 27 |
| Geotrupes stercorarius (Linnaeus, 1758) | 6 | 6 | 40 | 6 | 8 | 66 |
| Trypocopris vernalis (Linnaeus, 1758) | 1 | 4 | 0 | 0 | 1 | 6 |

Scarabiidae [83/57]

| | | | | | | |
|--|----|----|----|----|-----|-----|
| Onthophagus (Paleonthophagus) fracticornis (Preyssler, 1790) | 0 | 0 | 1 | 0 | 0 | 1 |
| Onthophagus (Paleonthophagus) joannae Goljan, 1953 | 0 | 0 | 0 | 1 | 0 | 1 |
| Onthophagus (Paleonthophagus) nuchicornis (Linnaeus, 1758) | 0 | 1 | 0 | 0 | 0 | 1 |
| Onthophagus (Paleonthophagus) similis (Scriba, 1790) | 0 | 0 | 3 | 0 | 0 | 3 |
| Onthophagus (Paleonthophagus) vacca (Linnaeus, 1767) | 0 | 0 | 1 | 0 | 0 | 1 |
| Aegialia (Aegialia) arenaria (Fabricius, 1787) | 12 | 5 | 0 | 0 | 0 | 17 |
| Aegialia (Psammoporus) insularis Pittino, 2006 | 1 | 6 | 1 | 3 | 3 | 14 |
| Euheptaulacus villosus (Gyllenhal in Schönherr, 1806) | 0 | 1 | 1 | 2 | 0 | 4 |
| Heptaulacus testudinarius (Fabricius, 1775) | 0 | 2 | 0 | 2 | 0 | 4 |
| Oxyomus sylvestris (Scopoli, 1763) | 0 | 2 | 3 | 0 | 0 | 5 |
| Aphodius (Acrossus) depressus (Kugelann, 1792) | 8 | 11 | 30 | 11 | 4 | 64 |
| Aphodius (Acrossus) luridus (Fabricius, 1775) | 4 | 5 | 12 | 2 | 0 | 23 |
| Aphodius (Acrossus) rufipes (Linnaeus, 1758) | 20 | 15 | 92 | 41 | 161 | 329 |
| Aphodius (Agoliinus) lapponum Gyllenhal, 1808 | 0 | 6 | 23 | 6 | 2 | 37 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|----|----|----|-----|
| Aphodius (Agrilinus) ater (De Geer, 1774) | 19 | 21 | 61 | 19 | 10 | 130 |
| Aphodius (Agrilinus) constans Duftschmid, 1805 | 0 | 5 | 3 | 5 | 3 | 16 |
| Aphodius (Agrilinus) rufus (Moll, 1782) | 11 | 6 | 25 | 3 | 11 | 56 |
| Aphodius (Agrilinus) sordidus (Fabricius, 1775) | 1 | 1 | 4 | 0 | 0 | 6 |
| Aphodius (Ammoecius) brevis (Erichson, 1848) | 0 | 0 | 0 | 1 | 0 | 1 |
| Aphodius (Aphodius) fimetarius (Linnaeus, 1758) | 7 | 17 | 43 | 5 | 5 | 77 |
| Aphodius (Aphodius) foetens (Fabricius, 1787) | 4 | 6 | 11 | 5 | 0 | 26 |
| Aphodius (Aphodius) foetidus (Herbst, 1783) | 3 | 3 | 0 | 0 | 1 | 7 |
| Aphodius (Bodilus) ictericus (Laicharting, 1781) | 2 | 0 | 0 | 1 | 0 | 3 |
| Aphodius (Calomosternus) granarius (Linnaeus, 1767) | 9 | 6 | 10 | 2 | 1 | 28 |
| Aphodius (Chilothorax) conspurcatus (Linnaeus, 1758) | 0 | 3 | 5 | 6 | 0 | 14 |
| Aphodius (Chilothorax) distinctus (Müller, O.F., 1776) | 5 | 4 | 8 | 2 | 0 | 19 |
| Aphodius (Chilothorax) paykulli Bedel, 1907 | 1 | 0 | 7 | 0 | 0 | 8 |
| Aphodius (Colobopterus) erraticus (Linnaeus, 1758) | 2 | 3 | 0 | 1 | 0 | 6 |
| Aphodius (Esymus) merdarius (Fabricius, 1775) | 3 | 4 | 3 | 1 | 0 | 11 |
| Aphodius (Esymus) pusillus (Herbst, 1789) | 2 | 2 | 8 | 4 | 2 | 18 |
| Aphodius (Euorodalus) coenosus (Panzer, 1798) | 0 | 0 | 1 | 0 | 0 | 1 |
| Aphodius (Eupleurus) subterraneus (Linnaeus, 1758) | 1 | 1 | 4 | 2 | 0 | 8 |
| Aphodius (Limarus) zenkeri Germar, 1813 | 1 | 0 | 3 | 0 | 0 | 4 |
| Aphodius (Liothorax) plagiatus (Linnaeus, 1767) | 4 | 0 | 0 | 0 | 0 | 4 |
| Aphodius (Melinopterus) prodromus (Brahm, 1790) | 18 | 18 | 48 | 12 | 5 | 101 |
| Aphodius (Melinopterus) sphacelatus (Panzer, 1798) | 7 | 16 | 68 | 8 | 6 | 105 |
| Aphodius (Nimbus) contaminatus (Herbst, 1783) | 9 | 12 | 46 | 6 | 4 | 77 |
| Aphodius (Nimbus) obliteratus Panzer, 1823 | 1 | 2 | 2 | 5 | 0 | 10 |
| Aphodius (Otophorus) haemorrhoidalis (Linnaeus, 1758) | 6 | 5 | 9 | 2 | 1 | 23 |
| Aphodius (Plagiogonus) arenarius (Olivier, 1789) | 1 | 0 | 4 | 2 | 1 | 8 |
| Aphodius (Planolinus) borealis Gyllenhal, 1827 | 0 | 3 | 5 | 2 | 2 | 12 |
| Aphodius (Planolinus) fasciatus (Olivier, 1789) | 0 | 7 | 2 | 1 | 5 | 15 |
| Aphodius (Sigorus) porcus (Fabricius, 1792) | 0 | 1 | 0 | 1 | 0 | 2 |
| Aphodius (Teuchestes) fossor (Linnaeus, 1758) | 19 | 20 | 40 | 8 | 9 | 96 |
| Aphodius (Volinus) sticticus (Panzer, 1798) | 1 | 0 | 1 | 2 | 1 | 5 |
| Psammodius asper (Fabricius, 1775) | 0 | 1 | 0 | 0 | 0 | 1 |
| Melolontha hippocastani Fabricius, 1801 | 0 | 0 | 1 | 0 | 0 | 1 |
| Melolontha melolontha (Linnaeus, 1758) | 5 | 9 | 41 | 23 | 65 | 143 |
| Polyphylla fullo (Linnaeus, 1758) | 0 | 0 | 1 | 0 | 0 | 1 |
| Amphimallon solstitiale (Linnaeus, 1758) | 1 | 4 | 1 | 0 | 0 | 6 |
| Serica brunnea (Linnaeus, 1758) | 9 | 6 | 39 | 20 | 20 | 94 |
| Hoplia philanthus (Füessly, 1775) | 0 | 5 | 3 | 10 | 1 | 19 |
| Anomala dubia (Scopoli, 1763) | 15 | 3 | 1 | 0 | 0 | 19 |
| Phyllopertha horticola (Linnaeus, 1758) | 12 | 4 | 9 | 1 | 0 | 26 |
| Oxythyrea funesta (Poda, 1761) | 0 | 0 | 2 | 0 | 0 | 2 |
| Cetonia aurata (Linnaeus, 1758) | 1 | 3 | 1 | 0 | 0 | 5 |
| Protaetia (Potosia) metallica (Herbst, 1782) | 0 | 1 | 0 | 0 | 0 | 1 |
| <u>Clambidae [10/10]</u> | | | | | | |
| Calyptomerus dubius (Marsham, 1802) | 18 | 3 | 16 | 3 | 0 | 40 |
| Clambus Fischer von Waldheim, 1823/4 | 0 | 0 | 1 | 0 | 0 | 1 |
| Clambus armadillo (De Geer, 1774) | 7 | 8 | 4 | 3 | 2 | 24 |
| Clambus evae Endrödy-Younga, 1960 | 0 | 1 | 0 | 0 | 1 | 2 |
| Clambus gibbulus (LeConte, 1850) | 0 | 0 | 4 | 0 | 0 | 4 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--------------------------------------|----|---|----|---|---|----|
| Clambus nigrellus Reitter, 1914 | 0 | 0 | 0 | 4 | 1 | 5 |
| Clambus nigriclavus Stephens, 1835 | 0 | 0 | 0 | 4 | 0 | 4 |
| Clambus pubescens Redtenbacher, 1849 | 12 | 4 | 17 | 1 | 1 | 35 |
| Clambus punctulum (Beck, 1817) | 3 | 3 | 17 | 1 | 0 | 24 |
| Clambus simsoni Blackburn, 1902 | 0 | 0 | 4 | 0 | 0 | 4 |

Scirtidae [20/18]

| | | | | | | |
|--|----|----|-----|----|----|-----|
| Elodes elongata Tournier, 1868 | 0 | 0 | 2 | 2 | 1 | 5 |
| Elodes minuta (Linnaeus, 1767) | 4 | 11 | 12 | 9 | 3 | 39 |
| Elodes pseudominuta Klausnitzer, 1971 | 1 | 12 | 31 | 1 | 4 | 49 |
| Elodes tricuspis Nyholm, 1985 | 0 | 0 | 0 | 1 | 0 | 1 |
| Odeles marginata (Fabricius, 1798) | 2 | 13 | 41 | 7 | 10 | 73 |
| Microcara testacea (Linnaeus, 1767) | 40 | 9 | 68 | 29 | 2 | 148 |
| Cyphon coarctatus Paykull, 1799 | 44 | 53 | 172 | 65 | 13 | 347 |
| Cyphon hilaris Nyholm, 1844 | 4 | 7 | 30 | 4 | 1 | 46 |
| Cyphon laevipennis Tournier, 1868 | 25 | 3 | 59 | 8 | 1 | 96 |
| Cyphon ochraceus Stephens, 1830 | 16 | 13 | 27 | 12 | 1 | 69 |
| Cyphon padi (Linnaeus, 1758) | 18 | 16 | 81 | 27 | 6 | 148 |
| Cyphon palustris Thomson, C.G., 1855 | 2 | 6 | 22 | 4 | 5 | 39 |
| Cyphon pubescens (Fabricius, 1792) | 1 | 0 | 1 | 1 | 0 | 3 |
| Cyphon punctipennis Sharp, 1873 | 1 | 2 | 0 | 0 | 0 | 3 |
| Cyphon variabilis (Thunberg, 1787) | 14 | 12 | 87 | 18 | 2 | 133 |
| Prionocyphon serricornis (Müller, P.W.J., 1821) | 0 | 5 | 6 | 1 | 0 | 12 |
| Hydrocyphon deflexicollis (Müller, P.W.J., 1821) | 0 | 4 | 0 | 0 | 3 | 7 |
| Scirtes hemisphaericus (Linnaeus, 1758) | 12 | 3 | 18 | 12 | 1 | 46 |

Dascillidae [1/1]

| | | | | | | |
|-------------------------------------|---|----|----|----|---|----|
| Dascillus cervinus (Linnaeus, 1758) | 5 | 12 | 11 | 14 | 6 | 48 |
|-------------------------------------|---|----|----|----|---|----|

Buprestidae [14/6]

| | | | | | | |
|---|---|---|---|---|---|----|
| Agrilus (Agrilus) viridis (Linnaeus, 1758) | 0 | 2 | 0 | 0 | 0 | 2 |
| Agrilus (Anambus) angustulus (Illiger, 1803) | 3 | 2 | 1 | 0 | 0 | 6 |
| Agrilus (Anambus) laticornis (Illiger, 1803) | 1 | 7 | 7 | 2 | 0 | 17 |
| Aphanisticus pusillus (Olivier, 1790) | 0 | 1 | 0 | 0 | 0 | 1 |
| Trachys minuta (Linnaeus, 1758) | 2 | 0 | 0 | 0 | 0 | 2 |
| Trachys troglodytes Gyllenhal in Schoenherr, 1817 | 1 | 1 | 0 | 0 | 0 | 2 |

Byrrhidae [13/11]

| | | | | | | |
|---|---|----|----|----|---|----|
| Simplocaria maculosa Erichson, 1847 | 0 | 0 | 0 | 1 | 0 | 1 |
| Simplocaria semistriata (Fabricius, 1794) | 8 | 8 | 39 | 14 | 1 | 70 |
| Morychus aeneus (Fabricius, 1775) | 1 | 0 | 0 | 0 | 2 | 3 |
| Cytilus sericeus (Forster, 1771) | 3 | 15 | 31 | 14 | 3 | 66 |
| Byrrhus arietinus Steffahn, 1842 | 0 | 1 | 4 | 1 | 0 | 6 |
| Byrrhus fasciatus (Forster, 1770) | 2 | 7 | 15 | 0 | 3 | 27 |
| Byrrhus pilula (Linnaeus, 1758) | 6 | 15 | 45 | 16 | 2 | 84 |
| Byrrhus pustulatus (Forster, 1770) | 2 | 5 | 7 | 1 | 1 | 16 |
| Porcinolus murinus (Fabricius, 1794) | 1 | 1 | 1 | 0 | 0 | 3 |
| Curimopsis nigrita (Palm, 1934) | 0 | 0 | 11 | 0 | 0 | 11 |
| Curimopsis setigera (Illiger, 1798) | 3 | 0 | 0 | 0 | 0 | 3 |

Elmidae [12/8]

| | | | | | | |
|---|---|----|----|----|----|----|
| <i>Elmis aenea</i> (Müller, P.W.J., 1806) | 4 | 15 | 17 | 18 | 19 | 73 |
| <i>Esolus parallelepipedus</i> (Müller, P.W.J., 1806) | 0 | 10 | 3 | 9 | 7 | 29 |
| <i>Limnius volckmari</i> (Panzer, 1793) | 0 | 12 | 6 | 18 | 14 | 50 |
| <i>Normandia nitens</i> (Müller, P.W.J., 1817) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Oulimnius troglodytes</i> (Gyllenhal, 1827) | 0 | 0 | 0 | 5 | 0 | 5 |
| <i>Oulimnius tuberculatus</i> (Müller, P.W.J., 1806) | 1 | 11 | 6 | 16 | 5 | 39 |
| <i>Riolus cupreus</i> (Müller, P.W.J., 1806) | 0 | 3 | 1 | 1 | 1 | 6 |
| <i>Riolus subviolaceus</i> (Müller, P.W.J., 1817) | 0 | 5 | 0 | 2 | 1 | 8 |

Dryopidae [9/6]

| | | | | | | |
|---|---|----|----|----|---|----|
| <i>Pomatinus substriatus</i> (Müller, P.W.J., 1806) | 0 | 2 | 1 | 0 | 0 | 3 |
| <i>Dryops</i> (<i>Dryops</i>) <i>auriculatus</i> (Fourcroy, 1785) | 3 | 1 | 2 | 1 | 2 | 9 |
| <i>Dryops</i> (<i>Dryops</i>) <i>ernesti</i> des Gozis, 1886 | 3 | 12 | 17 | 17 | 6 | 55 |
| <i>Dryops</i> (<i>Dryops</i>) <i>griseus</i> (Erichson, 1847) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Dryops</i> (<i>Dryops</i>) <i>luridus</i> (Erichson, 1847) | 5 | 1 | 8 | 3 | 1 | 18 |
| <i>Dryops</i> (<i>Yrdops</i>) <i>nitidulus</i> (Heer, 1841) | 0 | 0 | 1 | 2 | 3 | 6 |

Limnichidae [1/1]

| | | | | | | |
|---|---|---|---|---|---|---|
| <i>Limnichus pygmaeus</i> (Sturm, 1807) | 0 | 1 | 6 | 0 | 0 | 7 |
|---|---|---|---|---|---|---|

Heteroceridae [9/5]

| | | | | | | |
|---|----|---|---|---|---|----|
| <i>Augyles maritimus</i> (Guérin-Méneville, 1844) | 7 | 0 | 3 | 1 | 0 | 11 |
| <i>Heterocerus fenestratus</i> (Thunberg, 1784) | 18 | 2 | 5 | 5 | 3 | 33 |
| <i>Heterocerus flexuosus</i> Stephens, 1828 | 3 | 1 | 4 | 1 | 0 | 9 |
| <i>Heterocerus fuscus</i> Kiesenwetter, 1843 | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Heterocerus marginatus</i> (Fabricius, 1787) | 8 | 6 | 7 | 8 | 0 | 29 |

Psephenidae [1/1]

| | | | | | | |
|--------------------------------------|---|---|---|---|---|----|
| <i>Eubria palustris</i> Germar, 1818 | 0 | 7 | 1 | 4 | 0 | 12 |
|--------------------------------------|---|---|---|---|---|----|

Eucnemidae [6/4]

| | | | | | | |
|---|---|---|----|----|---|----|
| <i>Melasis buprestoides</i> (Linnaeus, 1761) | 0 | 2 | 27 | 13 | 0 | 42 |
| <i>Epiphanis cornutus</i> Eschscholtz, 1829 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Microrhagus pygmaeus</i> (Fabricius, 1792) | 0 | 4 | 0 | 0 | 0 | 4 |
| <i>Eucnemis capucina</i> Ahrens, 1812 | 0 | 0 | 1 | 0 | 0 | 1 |

Throscidae [5/3]

| | | | | | | |
|---|---|----|----|---|---|----|
| <i>Trixagus carinifrons</i> (de Bonvouloir, 1859) | 1 | 1 | 15 | 1 | 0 | 18 |
| <i>Trixagus dermestoides</i> (Linnaeus, 1767) | 3 | 17 | 56 | 9 | 2 | 87 |
| <i>Trixagus obtusus</i> (Curtis, 1827) | 0 | 0 | 2 | 0 | 2 | 4 |

Elateridae [73/38]

| | | | | | | |
|--|----|----|----|----|----|-----|
| <i>Agrypnus murinus</i> (Linnaeus, 1758) | 20 | 3 | 4 | 1 | 0 | 28 |
| <i>Hypnoidus riparius</i> (Fabricius, 1792) | 11 | 16 | 79 | 23 | 24 | 153 |
| <i>Actenicerus sjaelandicus</i> (Müller, O.F., 1764) | 1 | 7 | 2 | 8 | 0 | 18 |
| <i>Anostirus castaneus</i> (Linnaeus, 1758) | 0 | 0 | 0 | 6 | 0 | 6 |
| <i>Ctenicera cuprea</i> (Fabricius, 1775) | 1 | 19 | 69 | 21 | 13 | 123 |
| <i>Ctenicera pectinicornis</i> (Linnaeus, 1758) | 3 | 8 | 13 | 6 | 3 | 33 |
| <i>Aplotarsus incanus</i> (Gyllenhal, 1827) | 25 | 47 | 87 | 21 | 14 | 194 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| Paraphotistus impressus (Fabricius, 1792) | 0 | 2 | 0 | 3 | 0 | 5 |
| Paraphotistus nigricornis (Panzer, 1799) | 24 | 2 | 2 | 1 | 0 | 29 |
| Prosternon tessellatum (Linnaeus, 1758) | 8 | 6 | 28 | 1 | 0 | 43 |
| Selatosomus aeneus (Linnaeus, 1758) | 4 | 2 | 12 | 7 | 1 | 26 |
| Cidnopus aeruginosus (Olivier, 1790) | 7 | 4 | 4 | 3 | 2 | 20 |
| Kibunea minuta (Linnaeus, 1758) | 24 | 15 | 93 | 4 | 2 | 138 |
| Denticollis linearis (Linnaeus, 1758) | 9 | 29 | 110 | 22 | 13 | 183 |
| Athous (Athous) haemorrhoidalis (Fabricius, 1801) | 58 | 72 | 547 | 60 | 19 | 756 |
| Athous (Athous) vittatus (Fabricius, 1792) | 4 | 13 | 23 | 7 | 3 | 50 |
| Athous (Exanathrotus) subfuscus (Müller, O.F., 1764) | 0 | 0 | 0 | 1 | 0 | 1 |
| Athous (Orthathous) bicolor (Goeze, 1777) | 12 | 3 | 32 | 3 | 0 | 50 |
| Athous (Orthathous) campyloides Newman, 1833 | 0 | 0 | 1 | 2 | 0 | 3 |
| Stenagostus rhombeus (Olivier, 1790) | 0 | 0 | 0 | 1 | 0 | 1 |
| Hemicrepidius hirtus (Herbst, 1784) | 23 | 36 | 62 | 18 | 1 | 140 |
| Adrastus pallens (Fabricius, 1792) | 18 | 22 | 78 | 16 | 10 | 144 |
| Agriotes acuminatus (Stephens, 1830) | 24 | 43 | 155 | 27 | 7 | 256 |
| Agriotes lineatus (Linnaeus, 1767) | 21 | 12 | 7 | 3 | 0 | 43 |
| Agriotes obscurus (Linnaeus, 1758) | 44 | 36 | 158 | 39 | 13 | 290 |
| Agriotes pallidulus (Illiger, 1807) | 32 | 86 | 508 | 80 | 31 | 737 |
| Agriotes sputator (Linnaeus, 1758) | 21 | 14 | 23 | 6 | 3 | 67 |
| Dalopius marginatus (Linnaeus, 1758) | 14 | 82 | 232 | 56 | 12 | 396 |
| Ampedus balteatus (Linnaeus, 1758) | 9 | 18 | 41 | 3 | 2 | 73 |
| Ampedus nigerrimus (Boisduval & Lacordaire, 1835) | 0 | 0 | 1 | 0 | 0 | 1 |
| Ampedus nigrinus (Herbst, 1784) | 0 | 1 | 1 | 1 | 0 | 3 |
| Ampedus pomorum (Herbst, 1784) | 0 | 4 | 1 | 0 | 0 | 5 |
| Ischnodes sanguinicollis (Panzer, 1793) | 0 | 1 | 0 | 0 | 0 | 1 |
| Sericus brunneus (Linnaeus, 1758) | 2 | 4 | 12 | 5 | 1 | 24 |
| Melanotus villosus (Geoffroy in Fourcroy, 1785) | 12 | 22 | 59 | 19 | 4 | 116 |
| Flautiauxellus maritimus (Curtis, 1840) | 1 | 0 | 0 | 3 | 6 | 10 |
| Oedostethus quadripustulatus (Fabricius, 1792) | 12 | 4 | 10 | 1 | 1 | 28 |
| Zorochochroa minimus (Boisduval & Lacordaire, 1835) | 0 | 4 | 12 | 15 | 22 | 53 |

Lycidae [4/2]

| | | | | | | |
|---------------------------------------|---|----|----|---|---|----|
| Pyropterus nigroruber (De Geer, 1774) | 2 | 5 | 25 | 2 | 0 | 34 |
| Platycis minutus (Fabricius, 1787) | 0 | 15 | 7 | 5 | 0 | 27 |

Lampyridae [3/1]

| | | | | | | |
|-------------------------------------|---|----|----|----|---|-----|
| Lampyris noctiluca (Linnaeus, 1758) | 6 | 11 | 94 | 16 | 6 | 133 |
|-------------------------------------|---|----|----|----|---|-----|

Cantharidae [41/38]

| | | | | | | |
|---|----|----|-----|----|----|-----|
| Podabrus alpinus (Paykull, 1798) | 2 | 14 | 29 | 13 | 7 | 65 |
| Ancistronycha abdominalis (Fabricius, 1798) | 2 | 12 | 14 | 3 | 0 | 31 |
| Cantharis cryptica Ashe, 1947 | 32 | 34 | 295 | 43 | 24 | 428 |
| Cantharis decipiens Baudi, 1871 | 16 | 26 | 206 | 28 | 8 | 284 |
| Cantharis figurata Mannerheim, 1843 | 0 | 0 | 7 | 3 | 1 | 11 |
| Cantharis fusca Linnaeus, 1758 | 2 | 0 | 4 | 3 | 1 | 10 |
| Cantharis lateralis Linnaeus, 1758 | 31 | 11 | 80 | 12 | 1 | 135 |
| Cantharis livida Linnaeus, 1758 | 11 | 14 | 102 | 13 | 4 | 144 |
| Cantharis nigra (De Geer, 1774) | 45 | 30 | 187 | 31 | 2 | 295 |
| Cantharis nigricans (Müller, O.F., 1776) | 32 | 38 | 295 | 41 | 21 | 427 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| <i>Cantharis pallida</i> Goeze, 1777 | 5 | 9 | 26 | 8 | 2 | 50 |
| <i>Cantharis paludosa</i> Fallén, 1807 | 1 | 14 | 20 | 12 | 7 | 54 |
| <i>Cantharis pellucida</i> Fabricius, 1792 | 9 | 32 | 139 | 17 | 12 | 209 |
| <i>Cantharis rufa</i> Linnaeus, 1758 | 33 | 19 | 133 | 19 | 2 | 206 |
| <i>Cantharis rustica</i> Fallén, 1807 | 10 | 13 | 59 | 10 | 6 | 98 |
| <i>Cantharis thoracica</i> (Olivier, 1790) | 18 | 6 | 25 | 8 | 2 | 59 |
| <i>Rhagonycha elongata</i> (Fallén, 1807) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Rhagonycha fulva</i> (Scopoli, 1763) | 79 | 51 | 753 | 57 | 19 | 959 |
| <i>Rhagonycha lignosa</i> (Müller, O.F., 1764) | 8 | 29 | 154 | 15 | 6 | 212 |
| <i>Rhagonycha limbata</i> Thomson, C.G., 1864 | 36 | 35 | 242 | 31 | 9 | 353 |
| <i>Rhagonycha lutea</i> (Müller, O.F., 1764) | 2 | 6 | 0 | 1 | 0 | 9 |
| <i>Rhagonycha testacea</i> (Linnaeus, 1758) | 5 | 23 | 64 | 26 | 12 | 130 |
| <i>Rhagonycha translucida</i> (Krynicky, 1832) | 0 | 8 | 5 | 1 | 0 | 14 |
| <i>Crudosilis ruficollis</i> (Fabricius, 1775) | 3 | 0 | 4 | 0 | 0 | 7 |
| <i>Malthinus flaveolus</i> (Herbst, 1786) | 10 | 21 | 97 | 11 | 2 | 141 |
| <i>Malthinus frontalis</i> (Marsham, 1802) | 2 | 6 | 12 | 8 | 1 | 29 |
| <i>Malthinus seriepunctatus</i> Kiesenwetter, 1852 | 3 | 6 | 9 | 0 | 0 | 18 |
| <i>Malthodes crassicornis</i> (Mäklin, 1846) | 0 | 2 | 0 | 1 | 0 | 3 |
| <i>Malthodes dispar</i> (Germar, 1824) | 7 | 3 | 3 | 12 | 4 | 29 |
| <i>Malthodes fibulatus</i> Kiesenwetter, 1852 | 0 | 4 | 0 | 3 | 0 | 7 |
| <i>Malthodes flavoguttatus</i> Kiesenwetter, 1852 | 0 | 7 | 6 | 4 | 0 | 17 |
| <i>Malthodes fuscus</i> (Waltl, 1838) | 2 | 6 | 15 | 7 | 0 | 30 |
| <i>Malthodes guttifer</i> Kiesenwetter, 1852 | 0 | 4 | 4 | 2 | 2 | 12 |
| <i>Malthodes marginatus</i> (Latreille, 1806) | 2 | 22 | 47 | 16 | 8 | 95 |
| <i>Malthodes maurus</i> (Laporte, 1840) | 0 | 4 | 0 | 1 | 0 | 5 |
| <i>Malthodes minimus</i> (Linnaeus, 1758) | 29 | 22 | 48 | 8 | 3 | 110 |
| <i>Malthodes mysticus</i> Kiesenwetter, 1852 | 0 | 33 | 24 | 13 | 7 | 77 |
| <i>Malthodes pumilus</i> (Brébisson, 1835) | 6 | 9 | 24 | 7 | 1 | 47 |

Derodontidae [1/1]

| | | | | | | |
|---|---|---|---|---|---|---|
| <i>Laricobius erichsonii</i> Rosenhauer, 1846 | 0 | 2 | 0 | 0 | 0 | 2 |
|---|---|---|---|---|---|---|

Dermestidae [40/19]

| | | | | | | |
|--|---|---|----|---|---|----|
| <i>Dermestes</i> (<i>Dermestinus</i>) <i>frischii</i> Kugelann, 1792 | 0 | 2 | 1 | 0 | 0 | 3 |
| <i>Dermestes</i> (<i>Dermestinus</i>) <i>maculatus</i> De Geer, 1774 | 3 | 3 | 6 | 3 | 0 | 15 |
| <i>Dermestes</i> (<i>Dermestinus</i>) <i>murinus</i> Linnaeus, 1758 | 7 | 2 | 5 | 1 | 0 | 15 |
| <i>Dermestes</i> (<i>Dermestinus</i>) <i>undulatus</i> Brahm, 1790 | 2 | 1 | 0 | 0 | 0 | 3 |
| <i>Dermestes</i> (<i>Dermestes</i>) <i>ater</i> De Geer, 1774 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Dermestes</i> (<i>Dermestes</i>) <i>haemorrhoidalis</i> Küster, 1852 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Dermestes</i> (<i>Dermestes</i>) <i>lardarius</i> Linnaeus, 1758 | 4 | 5 | 15 | 1 | 0 | 25 |
| <i>Dermestes</i> (<i>Dermestes</i>) <i>peruvianus</i> Laporte de Castelnau, 1840 | 0 | 0 | 7 | 0 | 0 | 7 |
| <i>Attagenus pelli</i> (Linnaeus, 1758) | 9 | 4 | 30 | 3 | 2 | 48 |
| <i>Attagenus unicolor</i> (Brahm, 1791) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Trogoderma granarium</i> Everts, 1898 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Trogoderma inclusum</i> LeConte, 1854 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Reesa vespulae</i> (Milliron, 1939) | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Megatoma undata</i> (Linnaeus, 1758) | 5 | 2 | 3 | 4 | 0 | 14 |
| <i>Ctesias serra</i> (Fabricius, 1792) | 0 | 9 | 3 | 4 | 0 | 16 |
| <i>Anthrenus</i> (<i>Florilinus</i>) <i>museorum</i> (Linnaeus, 1761) | 1 | 3 | 6 | 0 | 0 | 10 |
| <i>Anthrenus</i> (<i>Helocerus</i>) <i>fuscus</i> Olivier, 1789 | 6 | 9 | 36 | 5 | 0 | 56 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|----|----|---|-----|
| <i>Anthrenus (Nathrenus) verbasci</i> (Linnaeus, 1767) | 6 | 3 | 22 | 5 | 0 | 36 |
| <i>Trinodes hirtus</i> (Fabricius, 1781) | 0 | 0 | 1 | 0 | 0 | 1 |
| <u>Bostrichidae [5/5]</u> | | | | | | |
| <i>Bostrichus capucinus</i> (Linnaeus, 1758) | 1 | 0 | 2 | 1 | 0 | 4 |
| <i>Rhyzopertha dominica</i> (Fabricius, 1792) | 2 | 1 | 2 | 0 | 0 | 5 |
| <i>Lyctus brunneus</i> (Stephens, 1830) | 0 | 3 | 6 | 0 | 0 | 9 |
| <i>Lyctus linearis</i> (Goeze, 1777) | 0 | 3 | 1 | 0 | 0 | 4 |
| <i>Lyctus planicollis</i> LeConte, 1858 | 0 | 0 | 1 | 0 | 0 | 1 |
| <u>Anobiidae [53/38]</u> | | | | | | |
| <i>Hedobia (Ptinomorphus) imperialis</i> (Linnaeus, 1767) | 8 | 6 | 12 | 6 | 0 | 32 |
| <i>Gibbium aequinoctiale</i> Boieldieu, 1854 | 0 | 0 | 6 | 0 | 0 | 6 |
| <i>Gibbium psylloides</i> (de Czenpinski, 1778) | 1 | 1 | 7 | 0 | 0 | 9 |
| <i>Stethomezium squamosum</i> Hinton, 1943 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Mezium affine</i> Boieldieu, 1856 | 0 | 1 | 1 | 1 | 0 | 3 |
| <i>Trigonogenius globulus</i> Solier, 1849 | 1 | 2 | 1 | 1 | 0 | 5 |
| <i>Niptus hololeucus</i> (Faldermann, 1835) | 4 | 5 | 14 | 5 | 0 | 28 |
| <i>Tipnus unicolor</i> (Piller & Mitterpacher, 1783) | 4 | 3 | 6 | 2 | 0 | 15 |
| <i>Pseudeurostus hilleri</i> (Reitter, 1877) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Ptinus clavipes</i> Panzer, 1792 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Ptinus fur</i> (Linnaeus, 1758) | 12 | 5 | 11 | 1 | 0 | 29 |
| <i>Ptinus lichenum</i> Marsham, 1802 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Ptinus palliatus</i> Perris, 1847 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Ptinus pusillus</i> Sturm, 1837 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Ptinus sexpunctatus</i> Panzer, 1792 | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Ptinus subpilosus</i> Sturm, 1837 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Ptinus tectus</i> Boieldieu, 1856 | 5 | 5 | 20 | 3 | 0 | 33 |
| <i>Grynobius planus</i> (Fabricius, 1787) | 4 | 17 | 9 | 6 | 1 | 37 |
| <i>Dryophilus pusillus</i> (Gyllenhal, 1808) | 5 | 22 | 14 | 3 | 5 | 49 |
| <i>Ochina ptinoides</i> (Marsham, 1802) | 0 | 2 | 2 | 3 | 0 | 7 |
| <i>Xestobium rufovillosum</i> (De Geer, 1774) | 5 | 8 | 6 | 6 | 1 | 26 |
| <i>Ernobius mollis</i> (Linnaeus, 1758) | 5 | 8 | 13 | 3 | 2 | 31 |
| <i>Ernobius nigrinus</i> (Sturm, 1837) | 1 | 2 | 0 | 0 | 0 | 3 |
| <i>Stegobium paniceum</i> (Linnaeus, 1758) | 2 | 4 | 28 | 3 | 0 | 37 |
| <i>Gastrallus immarginatus</i> (Müller, P.W.J., 1821) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Anobium</i> Fabricius, 1775 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Anobium fulvicorne</i> (Sturm, 1837) | 6 | 5 | 18 | 8 | 0 | 37 |
| <i>Anobium inexpectatum</i> Lohse, 1954 | 1 | 0 | 4 | 1 | 1 | 7 |
| <i>Anobium punctatum</i> (De Geer, 1774) | 16 | 12 | 30 | 13 | 1 | 72 |
| <i>Hadrobregmus denticollis</i> (Creutzer in Panzer, 1796) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Ptilinus pectinicornis</i> (Linnaeus, 1758) | 17 | 32 | 72 | 20 | 3 | 144 |
| <i>Lasioderma serricorne</i> (Fabricius, 1792) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Dorcatoma chrysomelina</i> (Sturm, 1837) | 1 | 3 | 4 | 9 | 0 | 17 |
| <i>Dorcatoma dresdensis</i> Herbst, 1792 | 4 | 3 | 2 | 0 | 0 | 9 |
| <i>Dorcatoma flavicornis</i> (Fabricius, 1792) | 1 | 4 | 4 | 5 | 0 | 14 |
| <i>Dorcatoma substriata</i> Hummel, 1829 | 0 | 4 | 1 | 0 | 0 | 5 |
| <i>Caenocara bovistae</i> (Hoffmann, J.J., 1803) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Anitys rubens</i> (Hoffmann, J.J., 1803) | 0 | 1 | 1 | 3 | 0 | 5 |

Lymexilidae [2/1]

| | | | | | | |
|--|---|----|----|---|---|----|
| Hylecoetus dermestoides (Linnaeus, 1761) | 5 | 17 | 31 | 7 | 0 | 60 |
|--|---|----|----|---|---|----|

Phloiophilidae [1/1]

| | | | | | | |
|---------------------------------------|---|---|---|---|---|---|
| Phloiophilus edwardsii Stephens, 1830 | 1 | 5 | 0 | 1 | 0 | 7 |
|---------------------------------------|---|---|---|---|---|---|

Trogossitidae [5/4]

| | | | | | | |
|---|---|---|---|---|---|----|
| Thymalus limbatus (Fabricius, 1787) | 0 | 5 | 2 | 6 | 0 | 13 |
| Lophocateres pusillus (Klug, 1832) | 0 | 0 | 1 | 0 | 0 | 1 |
| Tenebroides mauritanicus (Linnaeus, 1758) | 1 | 2 | 4 | 0 | 0 | 7 |
| Nemozoma elongatum (Linnaeus, 1761) | 1 | 0 | 1 | 0 | 0 | 2 |

Cleridae [14/8]

| | | | | | | |
|---|----|---|----|---|---|----|
| Tillus elongatus (Linnaeus, 1758) | 0 | 3 | 0 | 0 | 0 | 3 |
| Opilo mollis (Linnaeus, 1758) | 0 | 1 | 0 | 0 | 0 | 1 |
| Thanasimus formicarius (Linnaeus, 1758) | 9 | 7 | 8 | 2 | 0 | 26 |
| Tarsostenus univittatus (Rossi, 1792) | 0 | 0 | 1 | 0 | 0 | 1 |
| Korynetes caeruleus (De Geer, 1775) | 2 | 1 | 2 | 1 | 0 | 6 |
| Necrobia ruficollis (Fabricius, 1775) | 3 | 3 | 3 | 2 | 0 | 11 |
| Necrobia rufipes (De Geer, 1775) | 8 | 5 | 9 | 1 | 0 | 23 |
| Necrobia violacea (Linnaeus, 1758) | 12 | 4 | 17 | 5 | 2 | 40 |

Dasytidae [9/8]

| | | | | | | |
|---|---|----|---|---|---|----|
| Aplocnemus impressus (Marsham, 1802) | 0 | 1 | 2 | 0 | 0 | 3 |
| Aplocnemus nigricornis (Fabricius, 1792) | 0 | 6 | 0 | 1 | 0 | 7 |
| Dasytes aeratus Stephens, 1830 | 4 | 15 | 7 | 9 | 3 | 38 |
| Dasytes niger (Linnaeus, 1761) | 0 | 0 | 1 | 0 | 0 | 1 |
| Dasytes plumbeus (Müller, O.F., 1776) | 0 | 2 | 3 | 0 | 0 | 5 |
| Dasytes virens (Marsham, 1802) | 0 | 0 | 2 | 0 | 0 | 2 |
| Psilothrix viridicoeruleus (Geoffroy in Fourcroy, 1785) | 0 | 0 | 1 | 0 | 0 | 1 |
| Dolichosoma lineare (Rossi, 1794) | 4 | 0 | 0 | 0 | 0 | 4 |

Malachiidae [16/4]

| | | | | | | |
|---|----|----|-----|----|---|-----|
| Ebaeus pedicularius (Linnaeus, 1758) | 0 | 0 | 1 | 0 | 0 | 1 |
| Cordylepherus viridis (Fabricius, 1787) | 1 | 0 | 1 | 2 | 0 | 4 |
| Malachius bipustulatus (Linnaeus, 1758) | 27 | 31 | 210 | 32 | 8 | 308 |
| Anthocomus rufus (Herbst, 1784) | 5 | 0 | 42 | 0 | 0 | 47 |

Sphindidae [2/2]

| | | | | | | |
|--|---|---|----|---|---|----|
| Sphindus dubius (Gyllenhal, 1808) | 0 | 7 | 1 | 0 | 0 | 8 |
| Aspidiphorus orbiculatus (Gyllenhal, 1808) | 2 | 9 | 18 | 2 | 0 | 31 |

Kateretidae [9/8]

| | | | | | | |
|--|----|----|-----|----|----|-----|
| Kateretes pedicularius (Linnaeus, 1758) | 1 | 3 | 4 | 7 | 0 | 15 |
| Kateretes pusillus (Thunberg, 1794) | 13 | 9 | 1 | 6 | 2 | 31 |
| Kateretes rufilabris (Latreille, 1807) | 26 | 10 | 31 | 15 | 6 | 88 |
| Brachypterus glaber (Stephens, 1832) | 57 | 31 | 230 | 34 | 12 | 364 |
| Brachypterus urticae (Fabricius, 1792) | 68 | 45 | 298 | 42 | 20 | 473 |
| Brachypterolus linariae (Stephens, 1830) | 0 | 0 | 4 | 0 | 0 | 4 |
| Brachypterolus pulicarius (Linnaeus, 1758) | 4 | 0 | 28 | 2 | 1 | 35 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|-----|-----|-----|-----|----|------|
| <i>Brachypterolus vestitus</i> (Kiesenwetter, 1850) | 0 | 0 | 2 | 0 | 0 | 2 |
| <u>Nitidulidae [88/68]</u> | | | | | | |
| <i>Carpophilus dimidiatus</i> (Fabricius, 1792) | 1 | 0 | 3 | 0 | 0 | 4 |
| <i>Carpophilus hemipterus</i> (Linnaeus, 1758) | 1 | 1 | 3 | 2 | 0 | 7 |
| <i>Carpophilus ligneus</i> Murray, 1864 | 0 | 1 | 4 | 1 | 0 | 6 |
| <i>Carpophilus marginellus</i> Motschulsky, 1858 | 6 | 0 | 28 | 1 | 1 | 36 |
| <i>Carpophilus mutilatus</i> Erichson, 1843 | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Carpophilus obsoletus</i> Erichson, 1843 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Carpophilus sexpustulatus</i> (Fabricius, 1792) | 1 | 1 | 15 | 4 | 0 | 21 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>aestiva</i> (Linnaeus, 1758) | 30 | 42 | 153 | 26 | 13 | 264 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>angustula</i> Sturm, 1844 | 0 | 2 | 3 | 1 | 0 | 6 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>biguttata</i> (Thunberg, 1784) | 13 | 22 | 85 | 17 | 5 | 142 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>fuscicollis</i> (Stephens, 1832) | 0 | 1 | 0 | 2 | 0 | 3 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>guttata</i> (Olivier, 1811) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>longula</i> Erichson, 1845 | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>marseuli</i> Reitter, 1872 | 3 | 4 | 9 | 3 | 0 | 19 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>melina</i> Erichson, 1843 | 4 | 7 | 11 | 1 | 0 | 23 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>pallescens</i> (Stephens, 1832) | 5 | 5 | 12 | 2 | 0 | 24 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>rufomarginata</i> (Stephens, 1830) | 0 | 3 | 8 | 0 | 0 | 11 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>silacea</i> (Herbst, 1783) | 3 | 13 | 13 | 8 | 1 | 38 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>terminalis</i> Mannerheim, 1843 | 0 | 3 | 1 | 0 | 0 | 4 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>thoracica</i> Tournier, 1872 | 0 | 3 | 0 | 1 | 0 | 4 |
| <i>Epuraea</i> (<i>Epuraea</i>) <i>variegata</i> (Herbst, 1793) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Epuraea</i> (<i>Epuraeanella</i>) <i>limbata</i> (Fabricius, 1787) | 2 | 1 | 1 | 0 | 0 | 4 |
| <i>Epuraea</i> (<i>Micruria</i>) <i>melanocephala</i> (Marsham, 1802) | 17 | 22 | 140 | 15 | 5 | 199 |
| <i>Pria dulcamarae</i> (Scopoli, 1763) | 7 | 1 | 18 | 6 | 0 | 32 |
| <i>Meligethes aeneus</i> (Fabricius, 1775) | 169 | 129 | 992 | 112 | 56 | 1458 |
| <i>Meligethes atramentarius</i> Förster, 1849 | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Meligethes atratus</i> (Olivier, 1790) | 7 | 12 | 60 | 10 | 0 | 89 |
| <i>Meligethes bidens</i> Brisout de Barneville, 1863 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Meligethes brevis</i> Sturm, 1845 | 4 | 5 | 0 | 1 | 0 | 10 |
| <i>Meligethes brunnicornis</i> Sturm, 1845 | 3 | 6 | 10 | 1 | 0 | 20 |
| <i>Meligethes carinulatus</i> Förster, 1849 | 19 | 9 | 28 | 1 | 2 | 59 |
| <i>Meligethes difficilis</i> (Heer, 1841) | 12 | 4 | 25 | 0 | 0 | 41 |
| <i>Meligethes flavimanus</i> Stephens, 1830 | 4 | 10 | 3 | 3 | 2 | 22 |
| <i>Meligethes fulvipes</i> Brisout de Barneville, 1863 | 5 | 3 | 1 | 1 | 0 | 10 |
| <i>Meligethes gagathinus</i> Erichson, 1845 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Meligethes incanus</i> Sturm, 1845 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Meligethes kunzei</i> Erichson, 1845 | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Meligethes lugubris</i> Sturm, 1845 | 2 | 0 | 1 | 0 | 0 | 3 |
| <i>Meligethes morosus</i> Erichson, 1845 | 11 | 4 | 11 | 0 | 1 | 27 |
| <i>Meligethes nigrescens</i> Stephens, 1830 | 35 | 14 | 67 | 9 | 1 | 126 |
| <i>Meligethes obscurus</i> Erichson, 1845 | 2 | 7 | 8 | 4 | 0 | 21 |
| <i>Meligethes ochropus</i> Sturm, 1845 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Meligethes ovatus</i> Sturm, 1845 | 4 | 2 | 5 | 0 | 0 | 11 |
| <i>Meligethes pedicularius</i> (Gyllenhal, 1808) | 7 | 2 | 14 | 1 | 2 | 26 |
| <i>Meligethes planiusculus</i> (Heer, 1841) | 0 | 1 | 8 | 0 | 0 | 9 |
| <i>Meligethes rotundicollis</i> Brisout de Barneville, 1863 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Meligethes ruficornis</i> (Marsham, 1802) | 1 | 0 | 0 | 0 | 0 | 1 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|----|----|---|-----|
| Meligethes solidus (Kugelann, 1794) | 0 | 0 | 2 | 3 | 0 | 5 |
| Meligethes subrugosus (Gyllenhal, 1808) | 7 | 0 | 0 | 0 | 0 | 7 |
| Meligethes viridescens (Fabricius, 1787) | 4 | 14 | 45 | 6 | 2 | 71 |
| Nitidula bipunctata (Linnaeus, 1758) | 4 | 5 | 9 | 5 | 2 | 25 |
| Nitidula flavomaculata Rossi, 1790 | 0 | 1 | 0 | 0 | 0 | 1 |
| Nitidula rufipes (Linnaeus, 1767) | 1 | 0 | 8 | 3 | 0 | 12 |
| Omosita colon (Linnaeus, 1758) | 12 | 1 | 15 | 6 | 1 | 35 |
| Omosita depressa (Linnaeus, 1758) | 0 | 3 | 12 | 3 | 0 | 18 |
| Omosita discoidea (Fabricius, 1775) | 11 | 7 | 35 | 13 | 1 | 67 |
| Soronia grisea (Linnaeus, 1758) | 7 | 8 | 24 | 9 | 1 | 49 |
| Soronia punctatissima (Illiger, 1794) | 0 | 4 | 5 | 6 | 0 | 15 |
| Cychramus luteus (Fabricius, 1787) | 1 | 11 | 13 | 2 | 0 | 27 |
| Pocadius adustus Reitter, 1888 | 0 | 4 | 8 | 0 | 0 | 12 |
| Pocadius ferrugineus (Fabricius, 1775) | 3 | 3 | 13 | 2 | 1 | 22 |
| Thalycra fervida (Olivier, 1790) | 0 | 0 | 1 | 2 | 0 | 3 |
| Cryptarcha strigata (Fabricius, 1787) | 0 | 0 | 2 | 2 | 0 | 4 |
| Cryptarcha undata (Olivier, 1790) | 0 | 0 | 0 | 3 | 0 | 3 |
| Glischrochilus (Glischrochilus) quadripunctatus (Linnaeus, 1758) | 4 | 8 | 7 | 2 | 1 | 22 |
| Glischrochilus (Librodor) hortensis (Geoffroy in Fourcroy, 1785) | 28 | 28 | 81 | 34 | 3 | 174 |
| Glischrochilus (Librodor) quadriguttatus (Fabricius, 1777) | 1 | 3 | 5 | 3 | 0 | 12 |
| Pityophagus ferrugineus (Linnaeus, 1761) | 0 | 2 | 1 | 0 | 0 | 3 |

Monotomidae [23/20]

| | | | | | | |
|---|----|----|-----|----|---|-----|
| Rhizophagus (Eurhizophagus) depressus (Fabricius, 1792) | 1 | 3 | 11 | 3 | 0 | 18 |
| Rhizophagus (Rhizophagus) bipustulatus (Fabricius, 1792) | 10 | 16 | 74 | 23 | 2 | 125 |
| Rhizophagus (Rhizophagus) dispar (Paykull, 1800) | 12 | 24 | 183 | 34 | 2 | 255 |
| Rhizophagus (Rhizophagus) ferrugineus (Paykull, 1800) | 2 | 6 | 5 | 2 | 0 | 15 |
| Rhizophagus (Rhizophagus) nitidulus (Fabricius, 1798) | 0 | 3 | 24 | 6 | 0 | 33 |
| Rhizophagus (Rhizophagus) oblongicollis Blatch & Horner, 1892 | 0 | 1 | 0 | 0 | 0 | 1 |
| Rhizophagus (Rhizophagus) parallelocollis Gyllenhal, 1827 | 1 | 4 | 6 | 0 | 0 | 11 |
| Rhizophagus (Rhizophagus) perforatus Erichson, 1845 | 3 | 4 | 28 | 7 | 0 | 42 |
| Rhizophagus (Rhizophagus) picipes (Olivier, 1790) | 1 | 0 | 7 | 3 | 0 | 11 |
| Rhizophagus (Anomophagus) cribratus Gyllenhal, 1827 | 0 | 0 | 4 | 4 | 0 | 8 |
| Cyanostolus aeneus (Richter, 1820) | 0 | 1 | 0 | 0 | 0 | 1 |
| Monotoma (Gyrocecis) angusticollis (Gyllenhal, 1827) | 0 | 3 | 0 | 0 | 0 | 3 |
| Monotoma (Gyrocecis) conicicollis Guérin-Ménéville, 1837 | 0 | 4 | 6 | 0 | 0 | 10 |
| Monotoma (Gyrocecis) quadrifoveolata Aubé, 1838 | 0 | 1 | 0 | 0 | 0 | 1 |
| Monotoma (Monotoma) bicolor Villa & Villa, 1835 | 5 | 2 | 27 | 5 | 1 | 40 |
| Monotoma (Monotoma) brevicollis Aubé, 1838 | 0 | 0 | 1 | 0 | 0 | 1 |
| Monotoma (Monotoma) longicollis (Gyllenhal, 1827) | 8 | 7 | 14 | 3 | 0 | 32 |
| Monotoma (Monotoma) picipes Herbst, 1793 | 12 | 8 | 33 | 9 | 2 | 64 |
| Monotoma (Monotoma) spinicollis Aubé, 1838 | 4 | 0 | 6 | 3 | 0 | 13 |
| Monotoma (Monotoma) testacea Motschulsky, 1845 | 0 | 0 | 6 | 2 | 0 | 8 |

Silvanidae [12/9]

| | | | | | | |
|--|---|---|----|---|---|----|
| Uleiota planata (Linnaeus, 1761) | 6 | 0 | 0 | 0 | 0 | 6 |
| Cryptomorpha desjardinsii (Guérin-Ménéville, 1844) | 0 | 0 | 1 | 0 | 0 | 1 |
| Psammoecus bipunctatus (Fabricius, 1792) | 3 | 0 | 5 | 0 | 0 | 8 |
| Nausibius clavicornis (Kugelann, 1794) | 0 | 0 | 1 | 0 | 0 | 1 |
| Ahasverus advena (Waltl, 1834) | 8 | 2 | 18 | 3 | 0 | 31 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|---|---|---|---|---|---|
| <i>Oryzaephilus mercator</i> (Fauvel, 1889) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Oryzaephilus surinamensis</i> (Linnaeus, 1758) | 2 | 0 | 3 | 0 | 0 | 5 |
| <i>Silvanus bidentatus</i> (Fabricius, 1792) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Silvanus unidentatus</i> (Olivier, 1790) | 0 | 2 | 2 | 0 | 0 | 4 |

Cucujidae [2/2]

| | | | | | | |
|--|---|---|---|---|---|----|
| <i>Pediacus depressus</i> (Herbst, 1797) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Pediacus dermestoides</i> (Fabricius, 1792) | 0 | 9 | 4 | 2 | 1 | 16 |

Laemophloeidae [11/2]

| | | | | | | |
|--|---|---|----|---|---|----|
| <i>Cryptolestes ferrugineus</i> (Stephens, 1831) | 3 | 5 | 16 | 3 | 0 | 27 |
| <i>Cryptolestes pusillus</i> (Schönherr, 1817) | 0 | 0 | 3 | 0 | 0 | 3 |

Phalacridae [15/11]

| | | | | | | |
|---|-----|----|-----|----|---|-----|
| <i>Phalacrus caricis</i> Sturm, 1807 | 5 | 4 | 8 | 1 | 0 | 18 |
| <i>Phalacrus corruscus</i> (Panzer, 1796) | 1 | 0 | 4 | 0 | 0 | 5 |
| <i>Phalacrus fimetarius</i> (Fabricius, 1775) | 21 | 2 | 18 | 3 | 0 | 44 |
| <i>Phalacrus substriatus</i> Gyllenhal, 1813 | 0 | 1 | 2 | 0 | 0 | 3 |
| <i>Olibrus aeneus</i> (Fabricius, 1792) | 33 | 5 | 63 | 10 | 1 | 112 |
| <i>Olibrus affinis</i> (Sturm, 1807) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Olibrus corticalis</i> (Panzer, 1796) | 6 | 0 | 2 | 0 | 0 | 8 |
| <i>Olibrus liquidus</i> Erichson, 1845 | 5 | 0 | 6 | 1 | 0 | 12 |
| <i>Olibrus pygmaeus</i> (Sturm, 1807) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Stilbus oblongus</i> (Erichson, 1845) | 4 | 0 | 9 | 0 | 0 | 13 |
| <i>Stilbus testaceus</i> (Panzer, 1796) | 111 | 40 | 158 | 26 | 8 | 343 |

Cryptophagidae [103/73]

| | | | | | | |
|---|----|---|----|----|---|-----|
| <i>Telmatophilus caricis</i> (Olivier, 1790) | 6 | 7 | 40 | 7 | 0 | 60 |
| <i>Telmatophilus schoenherrii</i> (Gyllenhal, 1808) | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Telmatophilus typhae</i> (Fallén, 1802) | 22 | 5 | 77 | 10 | 3 | 117 |
| <i>Paramecosoma melanocephalum</i> (Herbst, 1793) | 0 | 6 | 5 | 8 | 2 | 21 |
| <i>Henoticus serratus</i> (Gyllenhal, 1808) | 2 | 7 | 11 | 3 | 0 | 23 |
| <i>Cryptophagus (Cryptophagus) acutangulus</i> Gyllenhal, 1827 | 5 | 5 | 19 | 3 | 0 | 32 |
| <i>Cryptophagus (Cryptophagus) angustus</i> Ganglbauer, 1899 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Cryptophagus (Cryptophagus) badius</i> Sturm, 1845 | 1 | 1 | 0 | 4 | 0 | 6 |
| <i>Cryptophagus (Cryptophagus) cellaris</i> (Scopoli, 1763) | 1 | 4 | 3 | 1 | 0 | 9 |
| <i>Cryptophagus (Cryptophagus) corticinus</i> Thomson, C.G., 1863 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Cryptophagus (Cryptophagus) dentatus</i> (Herbst, 1793) | 8 | 9 | 58 | 12 | 4 | 91 |
| <i>Cryptophagus (Cryptophagus) distinguendus</i> Sturm, 1845 | 9 | 1 | 18 | 7 | 1 | 36 |
| <i>Cryptophagus (Cryptophagus) immixtus</i> Rey, 1889 | 1 | 3 | 2 | 2 | 0 | 8 |
| <i>Cryptophagus (Cryptophagus) labilis</i> Erichson, 1846 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Cryptophagus (Cryptophagus) lapponicus</i> Gyllenhal, 1827 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Cryptophagus (Cryptophagus) laticollis</i> Lucas, 1846 | 7 | 6 | 19 | 2 | 4 | 38 |
| <i>Cryptophagus (Cryptophagus) lycoperdi</i> (Scopoli, 1763) | 5 | 1 | 31 | 5 | 0 | 42 |
| <i>Cryptophagus (Cryptophagus) obsoletus</i> Reitter, 1879 | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Cryptophagus (Cryptophagus) pallidus</i> Sturm, 1845 | 4 | 4 | 4 | 1 | 0 | 13 |
| <i>Cryptophagus (Cryptophagus) pilosus</i> Gyllenhal, 1827 | 9 | 4 | 43 | 7 | 0 | 63 |
| <i>Cryptophagus (Cryptophagus) populi</i> Paykull, 1800 | 0 | 1 | 5 | 1 | 0 | 7 |
| <i>Cryptophagus (Cryptophagus) pseudodentatus</i> Bruce, 1934 | 3 | 3 | 20 | 9 | 3 | 38 |
| <i>Cryptophagus (Cryptophagus) pubescens</i> Sturm, 1845 | 4 | 5 | 12 | 3 | 2 | 26 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| Cryptophagus (Cryptophagus) ruficornis Stephens, 1830 | 5 | 0 | 13 | 1 | 0 | 19 |
| Cryptophagus (Cryptophagus) saginatus Sturm, 1845 | 2 | 1 | 3 | 0 | 0 | 6 |
| Cryptophagus (Cryptophagus) scanicus (Linnaeus, 1758) | 3 | 12 | 23 | 8 | 2 | 48 |
| Cryptophagus (Cryptophagus) schmidtii Leunis in Sturm, 1845 | 0 | 1 | 3 | 0 | 0 | 4 |
| Cryptophagus (Cryptophagus) scutellatus Newman, 1834 | 5 | 3 | 23 | 3 | 0 | 34 |
| Cryptophagus (Cryptophagus) setulosus Sturm, 1845 | 8 | 13 | 15 | 4 | 4 | 44 |
| Cryptophagus (Cryptophagus) subdepressus Gyllenhal, 1827 | 0 | 1 | 0 | 0 | 0 | 1 |
| Micrambe abietis (Paykull, 1798) | 1 | 0 | 0 | 0 | 0 | 1 |
| Micrambe lindbergorum (Bruce, 1934) | 6 | 0 | 0 | 0 | 1 | 7 |
| Micrambe villosa (Heer, 1841) | 13 | 4 | 10 | 0 | 0 | 27 |
| Micrambe vini (Panzer, 1797) | 48 | 18 | 84 | 5 | 15 | 170 |
| Antherophagus nigricornis (Fabricius, 1787) | 4 | 7 | 33 | 4 | 1 | 49 |
| Antherophagus pallens (Linnaeus, 1758) | 2 | 5 | 23 | 8 | 1 | 39 |
| Caenoscelis ferruginea (Sahlberg, C.R., 1820) | 0 | 3 | 4 | 0 | 0 | 7 |
| Caenoscelis sibirica Reitter, 1889 | 0 | 0 | 1 | 0 | 0 | 1 |
| Caenoscelis subdeplanata Brisout de Barneville, 1882 | 2 | 0 | 19 | 1 | 0 | 22 |
| Atomaria (Anchicera) apicalis Erichson, 1846 | 9 | 6 | 30 | 3 | 0 | 48 |
| Atomaria (Anchicera) atra (Herbst, 1793) | 1 | 1 | 0 | 0 | 0 | 2 |
| Atomaria (Anchicera) atricapilla Stephens, 1830 | 28 | 21 | 128 | 16 | 6 | 199 |
| Atomaria (Anchicera) basalis Erichson, 1846 | 8 | 1 | 2 | 1 | 0 | 12 |
| Atomaria (Anchicera) fuscata (Schönherr, 1808) | 56 | 29 | 143 | 34 | 4 | 266 |
| Atomaria (Anchicera) fuscipes (Gyllenhal, 1808) | 2 | 5 | 2 | 2 | 1 | 12 |
| Atomaria (Anchicera) hislopi Wollaston, 1857 | 0 | 1 | 0 | 0 | 0 | 1 |
| Atomaria (Anchicera) lewisi Reitter, 1877 | 67 | 31 | 251 | 41 | 10 | 400 |
| Atomaria (Anchicera) mesomela (Herbst, 1792) | 11 | 1 | 27 | 11 | 3 | 53 |
| Atomaria (Anchicera) morio Kolenati, 1846 | 2 | 0 | 0 | 0 | 0 | 2 |
| Atomaria (Anchicera) munda Erichson, 1846 | 2 | 1 | 7 | 3 | 0 | 13 |
| Atomaria (Anchicera) nigripennis (Kugelann, 1794) | 0 | 0 | 1 | 0 | 0 | 1 |
| Atomaria (Anchicera) nitidula (Marsham, 1802) | 64 | 33 | 267 | 39 | 12 | 415 |
| Atomaria (Anchicera) peltata Kraatz, 1853 | 0 | 1 | 0 | 0 | 0 | 1 |
| Atomaria (Anchicera) pusilla (Paykull, 1798) | 3 | 7 | 7 | 4 | 2 | 23 |
| Atomaria (Anchicera) rhenana Kraatz, 1853 | 11 | 0 | 0 | 0 | 0 | 11 |
| Atomaria (Anchicera) rubella Heer, 1841 | 17 | 5 | 44 | 10 | 1 | 77 |
| Atomaria (Anchicera) scutellaris Motschulsky, 1849 | 0 | 0 | 3 | 0 | 0 | 3 |
| Atomaria (Anchicera) testacea Stephens, 1830 | 74 | 16 | 165 | 34 | 17 | 306 |
| Atomaria (Atomaria) barani Brisout de Barneville, 1863 | 1 | 0 | 0 | 2 | 0 | 3 |
| Atomaria (Atomaria) diluta Erichson, 1846 | 0 | 0 | 2 | 5 | 0 | 7 |
| Atomaria (Atomaria) fimetarii (Fabricius, 1792) | 3 | 1 | 5 | 2 | 0 | 11 |
| Atomaria (Atomaria) impressa Erichson, 1846 | 0 | 1 | 0 | 1 | 1 | 3 |
| Atomaria (Atomaria) linearis Stephens, 1830 | 20 | 16 | 50 | 11 | 7 | 104 |
| Atomaria (Atomaria) lohsei Johnson & Strand, A., 1968 | 0 | 1 | 3 | 0 | 0 | 4 |
| Atomaria (Atomaria) nigrirostris Stephens 1830 | 8 | 7 | 57 | 10 | 2 | 84 |
| Atomaria (Atomaria) nigriventris Stephens, 1830 | 0 | 2 | 3 | 1 | 0 | 6 |
| Atomaria (Atomaria) pulchra Erichson, 1846 | 0 | 5 | 4 | 1 | 0 | 10 |
| Atomaria (Atomaria) puncticollis Thomson, C.G., 1868 | 1 | 0 | 0 | 0 | 0 | 1 |
| Atomaria (Atomaria) punctithorax Reitter, 1887 | 1 | 0 | 0 | 0 | 0 | 1 |
| Atomaria (Atomaria) strandi Johnson, 1967 | 0 | 1 | 3 | 0 | 0 | 4 |
| Atomaria (Atomaria) umbrina (Gyllenhal, 1827) | 0 | 1 | 1 | 0 | 0 | 2 |
| Ootypus globosus (Waltl, 1838) | 5 | 5 | 9 | 3 | 0 | 22 |
| Ephistemus globulus (Paykull, 1798) | 37 | 12 | 125 | 18 | 7 | 199 |

Erotylidae [7/6]

| | | | | | | |
|--|---|----|----|----|---|-----|
| <i>Dacne bipustulata</i> (Thunberg, 1781) | 8 | 14 | 40 | 22 | 2 | 86 |
| <i>Dacne rufifrons</i> (Fabricius, 1775) | 3 | 9 | 7 | 8 | 1 | 28 |
| <i>Triplax aenea</i> (Schaller, 1783) | 2 | 13 | 70 | 15 | 7 | 107 |
| <i>Triplax russica</i> (Linnaeus, 1758) | 2 | 0 | 15 | 0 | 0 | 17 |
| <i>Triplax scutellaris</i> Charpentier, 1825 | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Tritoma bipustulata</i> Fabricius, 1775 | 0 | 2 | 4 | 0 | 0 | 6 |

Byturidae [2/2]

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Byturus ochraceus</i> (Scriba, 1790) | 10 | 9 | 52 | 9 | 1 | 81 |
| <i>Byturus tomentosus</i> (De Geer, 1774) | 29 | 54 | 280 | 27 | 10 | 400 |

Biphyllidae [2/1]

| | | | | | | |
|--|----|---|----|---|---|----|
| <i>Biphyllus lunatus</i> (Fabricius, 1787) | 13 | 8 | 20 | 8 | 1 | 50 |
|--|----|---|----|---|---|----|

Bothrideridae [5/2]

| | | | | | | |
|--|---|---|---|---|---|---|
| <i>Teredus cylindricus</i> (Olivier, 1790) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Anommatus duodecimstriatus</i> (Müller, P.W.J., 1821) | 4 | 1 | 4 | 0 | 0 | 9 |

Cerylonidae [5/3]

| | | | | | | |
|---|---|----|----|----|---|-----|
| <i>Cerylon fagi</i> Brisout de Barneville, 1867 | 1 | 0 | 0 | 1 | 0 | 2 |
| <i>Cerylon ferrugineum</i> Stephens, 1830 | 1 | 17 | 74 | 19 | 4 | 115 |
| <i>Cerylon histeroides</i> (Fabricius, 1792) | 4 | 3 | 26 | 14 | 4 | 51 |

Alexiidae [1/1]

| | | | | | | |
|---|---|---|---|---|---|---|
| <i>Sphaerosoma pilosum</i> (Panzer, 1793) | 0 | 0 | 1 | 0 | 0 | 1 |
|---|---|---|---|---|---|---|

Endomychidae [8/3]

| | | | | | | |
|---|----|----|----|----|---|----|
| <i>Endomychus coccineus</i> (Linnaeus, 1758) | 4 | 12 | 20 | 11 | 0 | 47 |
| <i>Symbiotes latus</i> Redtenbacher, 1849 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Mycetaea subterranea</i> (Fabricius, 1801) | 27 | 5 | 33 | 3 | 1 | 69 |

Coccinellidae [53/41]

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Coccidula rufa</i> (Herbst, 1783) | 96 | 46 | 325 | 46 | 11 | 524 |
| <i>Coccidula scutellata</i> (Herbst, 1783) | 10 | 2 | 33 | 2 | 0 | 47 |
| <i>Rhyzobius litura</i> (Fabricius, 1787) | 72 | 43 | 223 | 26 | 11 | 375 |
| <i>Clitostethus arcuatus</i> (Rossi, 1794) | 0 | 0 | 4 | 0 | 0 | 4 |
| <i>Stethorus punctillum</i> (Weise, 1891) | 1 | 1 | 4 | 0 | 0 | 6 |
| <i>Scymnus</i> (<i>Scymnus</i>) <i>femoralis</i> (Gyllenhal, 1827) | 3 | 2 | 3 | 0 | 0 | 8 |
| <i>Scymnus</i> (<i>Scymnus</i>) <i>frontalis</i> (Fabricius, 1787) | 2 | 1 | 0 | 0 | 0 | 3 |
| <i>Scymnus</i> (<i>Scymnus</i>) <i>nigrinus</i> Kugelann, 1794 | 1 | 2 | 2 | 2 | 0 | 7 |
| <i>Scymnus</i> (<i>Scymnus</i>) <i>schmidti</i> Fürsch, 1958 | 3 | 0 | 0 | 0 | 0 | 3 |
| <i>Scymnus</i> (<i>Pullus</i>) <i>auritus</i> Thunberg, 1795 | 8 | 4 | 18 | 4 | 0 | 34 |
| <i>Scymnus</i> (<i>Pullus</i>) <i>suturalis</i> Thunberg, 1795 | 12 | 19 | 19 | 3 | 2 | 55 |
| <i>Scymnus</i> (<i>Neopullus</i>) <i>haemorrhoidalis</i> Herbst, 1797 | 3 | 3 | 12 | 1 | 0 | 19 |
| <i>Scymnus</i> (<i>Neopullus</i>) <i>limbatus</i> Stephens, 1832 | 2 | 0 | 3 | 1 | 0 | 6 |
| <i>Nephus redtenbacheri</i> (Mulsant, 1846) | 12 | 7 | 16 | 5 | 0 | 40 |
| <i>Hyperaspis pseudopustulata</i> Mulsant, 1853 | 2 | 4 | 5 | 0 | 0 | 11 |
| <i>Chilocorus bipustulatus</i> (Linnaeus, 1758) | 4 | 21 | 31 | 1 | 0 | 57 |
| <i>Chilocorus renipustulatus</i> (Scriba, 1791) | 21 | 11 | 85 | 18 | 2 | 137 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|-----|----|------|----|----|------|
| <i>Exochomus nigromaculatus</i> (Goeze, 1777) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Exochomus quadripustulatus</i> (Linnaeus, 1758) | 16 | 12 | 143 | 12 | 2 | 185 |
| <i>Anisosticta novemdecimpunctata</i> (Linnaeus, 1758) | 34 | 9 | 192 | 34 | 7 | 276 |
| <i>Tytthaspis sedecimpunctata</i> (Linnaeus, 1758) | 0 | 1 | 5 | 2 | 0 | 8 |
| <i>Myzia oblongoguttata</i> (Linnaeus, 1758) | 4 | 14 | 11 | 7 | 2 | 38 |
| <i>Myrrha octodecimguttata</i> (Linnaeus, 1758) | 12 | 13 | 20 | 10 | 0 | 55 |
| <i>Propylea quattuordecimpunctata</i> (Linnaeus, 1758) | 110 | 87 | 1086 | 78 | 31 | 1392 |
| <i>Calvia quattuordecimguttata</i> (Linnaeus, 1758) | 31 | 44 | 313 | 50 | 25 | 463 |
| <i>Halyzia sedecimguttata</i> (Linnaeus, 1758) | 6 | 7 | 63 | 9 | 3 | 88 |
| <i>Psyllobora vigintiduopunctata</i> (Linnaeus, 1758) | 63 | 15 | 398 | 30 | 5 | 511 |
| <i>Anatis ocellata</i> (Linnaeus, 1758) | 6 | 22 | 69 | 14 | 5 | 116 |
| <i>Aphidecta oblitterata</i> (Linnaeus, 1758) | 10 | 37 | 46 | 13 | 17 | 123 |
| <i>Hippodamia</i> (<i>Hippodamia</i>) <i>tredecimpunctata</i> (Linnaeus, 1758) | 1 | 2 | 3 | 0 | 0 | 6 |
| <i>Hippodamia</i> (<i>Adonia</i>) <i>variegata</i> (Goeze, 1777) | 22 | 5 | 95 | 13 | 1 | 136 |
| <i>Coccinella hieroglyphica</i> Linnaeus, 1758 | 6 | 18 | 46 | 4 | 1 | 75 |
| <i>Coccinella magnifica</i> Redtenbacher, 1843 | 0 | 0 | 5 | 0 | 0 | 5 |
| <i>Coccinella quinquepunctata</i> Linnaeus, 1758 | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Coccinella septempunctata</i> Linnaeus, 1758 | 124 | 85 | 1442 | 90 | 48 | 1789 |
| <i>Coccinella undecimpunctata</i> Linnaeus, 1758 | 40 | 8 | 156 | 7 | 3 | 214 |
| <i>Adalia bipunctata</i> (Linnaeus, 1758) | 67 | 45 | 751 | 50 | 24 | 937 |
| <i>Adalia decempunctata</i> (Linnaeus, 1758) | 47 | 44 | 350 | 43 | 14 | 498 |
| <i>Harmonia axyridis</i> (Pallas, 1773) | 12 | 0 | 43 | 1 | 3 | 59 |
| <i>Harmonia quadripunctata</i> (Pontoppidan, 1763) | 2 | 1 | 33 | 2 | 0 | 38 |
| <i>Subcoccinella vigintiquatuorpunctata</i> (Linnaeus, 1758) | 48 | 10 | 28 | 1 | 0 | 87 |

Corylophidae [11/8]

| | | | | | | |
|---|----|---|----|---|---|----|
| <i>Sericoderus lateralis</i> (Gyllenhal, 1827) | 10 | 1 | 24 | 2 | 2 | 39 |
| <i>Corylophus cassidoides</i> (Marsham, 1802) | 2 | 0 | 4 | 0 | 0 | 6 |
| <i>Corylophus sublaevipennis</i> Jacquelin du Val, 1859 | 3 | 0 | 0 | 0 | 0 | 3 |
| <i>Orthoperus aequalis</i> Sharp, 1885 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Orthoperus atomus</i> (Gyllenhal, 1808) | 0 | 0 | 2 | 1 | 0 | 3 |
| <i>Orthoperus brunnipes</i> (Gyllenhal, 1808) | 9 | 2 | 0 | 0 | 0 | 11 |
| <i>Orthoperus corticalis</i> (Redtenbacher, 1849) | 1 | 1 | 4 | 0 | 0 | 6 |
| <i>Orthoperus nigrescens</i> Stephens, 1829 | 1 | 0 | 2 | 0 | 0 | 3 |

Latridiidae [56/38]

| | | | | | | |
|--|-----|----|-----|----|---|-----|
| <i>Lithostygnus serripennis</i> Broun, 1914 | 14 | 0 | 11 | 1 | 0 | 26 |
| <i>Stephostethus angusticollis</i> (Gyllenhal, 1827) | 1 | 0 | 3 | 0 | 0 | 4 |
| <i>Stephostethus lardarius</i> (De Geer, 1775) | 70 | 39 | 158 | 39 | 5 | 311 |
| <i>Cartodere</i> (<i>Cartodere</i>) <i>constricta</i> (Gyllenhal 1827) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Cartodere</i> (<i>Aridius</i>) <i>bifasciata</i> (Reitter, 1877) | 111 | 42 | 350 | 47 | 9 | 559 |
| <i>Cartodere</i> (<i>Aridius</i>) <i>nodifer</i> (Westwood, 1839) | 76 | 38 | 272 | 44 | 9 | 439 |
| <i>Cartodere</i> (<i>Aridius</i>) <i>norvegica</i> (Strand, 1940) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Latridius anthracinus</i> Mannerheim, 1844 | 17 | 1 | 6 | 2 | 1 | 27 |
| <i>Latridius assimilis</i> (Mannerheim, 1844) | 2 | 2 | 16 | 3 | 1 | 24 |
| <i>Latridius consimilis</i> Mannerheim, 1844 | 1 | 0 | 1 | 3 | 0 | 5 |
| <i>Latridius minutus</i> (Linnaeus, 1767) | 1 | 4 | 17 | 1 | 1 | 24 |
| <i>Thes bergrothi</i> (Reitter, 1880) | 0 | 2 | 2 | 2 | 0 | 6 |
| <i>Enicmus brevicornis</i> (Mannerheim, 1844) | 2 | 0 | 3 | 0 | 1 | 6 |
| <i>Enicmus fungicola</i> Thomson, C.G., 1868 | 1 | 0 | 10 | 2 | 0 | 13 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Enicmus histrio</i> Joy & Tomlin, 1910 | 8 | 8 | 8 | 4 | 2 | 30 |
| <i>Enicmus rugosus</i> (Herbst, 1793) | 0 | 4 | 1 | 1 | 0 | 6 |
| <i>Enicmus testaceus</i> (Stephens, 1830) | 2 | 1 | 6 | 2 | 0 | 11 |
| <i>Enicmus transversus</i> (Olivier, 1790) | 68 | 23 | 185 | 33 | 12 | 321 |
| <i>Dienerella elongata</i> (Curtis, 1830) | 0 | 1 | 6 | 1 | 0 | 8 |
| <i>Dienerella filiformis</i> (Gyllenhal, 1827) | 0 | 1 | 2 | 0 | 0 | 3 |
| <i>Dienerella filum</i> (Aubé, 1850) | 1 | 0 | 3 | 0 | 0 | 4 |
| <i>Dienerella ruficollis</i> (Marsham, 1802) | 7 | 1 | 7 | 1 | 0 | 16 |
| <i>Corticaria crenulata</i> (Gyllenhal, 1827) | 18 | 1 | 7 | 0 | 0 | 26 |
| <i>Corticaria elongata</i> (Gyllenhal, 1827) | 10 | 9 | 28 | 5 | 0 | 52 |
| <i>Corticaria fagi</i> Wollaston, 1854 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Corticaria ferruginea</i> Marsham, 1802 | 1 | 1 | 3 | 0 | 0 | 5 |
| <i>Corticaria fulva</i> (Comolli, 1837) | 1 | 1 | 4 | 1 | 0 | 7 |
| <i>Corticaria impressa</i> (Olivier, 1790) | 67 | 10 | 103 | 12 | 0 | 192 |
| <i>Corticaria inconspicua</i> Wollaston, 1860 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Corticaria punctulata</i> Marsham, 1802 | 14 | 2 | 29 | 2 | 0 | 47 |
| <i>Corticaria rubripes</i> Mannerheim, 1844 | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Corticaria serrata</i> (Paykull, 1798) | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Corticaria umbilicata</i> (Beck, 1817) | 1 | 4 | 1 | 1 | 0 | 7 |
| <i>Corticarina fuscata</i> (Gyllenhal, 1827) | 70 | 62 | 206 | 51 | 15 | 404 |
| <i>Corticarina similata</i> (Gyllenhal, 1827) | 0 | 4 | 1 | 1 | 0 | 6 |
| <i>Corticinara gibbosa</i> (Herbst, 1793) | 94 | 74 | 308 | 38 | 7 | 521 |
| <i>Melanophthalma curticolis</i> (Mannerheim, 1844) | 3 | 0 | 0 | 0 | 0 | 3 |
| <i>Melanophthalma suturalis</i> (Mannerheim, 1844) | 1 | 0 | 0 | 0 | 0 | 1 |

Mycetophagidae [13/10]

| | | | | | | |
|---|----|----|----|----|---|-----|
| <i>Pseudotriphyllus suturalis</i> (Fabricius, 1801) | 12 | 23 | 20 | 13 | 4 | 72 |
| <i>Triphyllus bicolor</i> (Fabricius, 1777) | 0 | 13 | 3 | 1 | 0 | 17 |
| <i>Litargus connexus</i> (Geoffroy in Fourcroy, 1785) | 8 | 6 | 20 | 17 | 1 | 52 |
| <i>Mycetophagus atomarius</i> (Fabricius, 1787) | 3 | 3 | 21 | 11 | 3 | 41 |
| <i>Mycetophagus multipunctatus</i> Fabricius, 1792 | 4 | 0 | 12 | 2 | 0 | 18 |
| <i>Mycetophagus piceus</i> (Fabricius, 1777) | 4 | 5 | 14 | 5 | 1 | 29 |
| <i>Mycetophagus populi</i> Fabricius, 1798 | 0 | 0 | 0 | 2 | 0 | 2 |
| <i>Mycetophagus quadriguttatus</i> Müller, P.W.J., 1821 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Mycetophagus quadripustulatus</i> (Linnaeus, 1761) | 8 | 16 | 69 | 23 | 5 | 121 |
| <i>Typhaea stercorea</i> (Linnaeus, 1758) | 17 | 11 | 39 | 11 | 2 | 80 |

Ciidae [22/16]

| | | | | | | |
|--|----|----|-----|----|---|-----|
| <i>Cis bidentatus</i> (Olivier, 1790) | 6 | 10 | 14 | 19 | 1 | 50 |
| <i>Cis bilamellatus</i> Wood, 1884 | 19 | 12 | 70 | 26 | 0 | 127 |
| <i>Cis boleti</i> (Scopoli, 1763) | 9 | 20 | 135 | 35 | 8 | 207 |
| <i>Cis fagi</i> Walth, 1839 | 2 | 1 | 0 | 1 | 0 | 4 |
| <i>Cis festivus</i> (Panzer, 1793) | 0 | 5 | 5 | 2 | 1 | 13 |
| <i>Cis hispidus</i> (Paykull, 1798) | 3 | 0 | 1 | 0 | 0 | 4 |
| <i>Cis lineatocribratus</i> Mellié, 1849 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Cis micans</i> (Fabricius, 1792) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Cis nitidus</i> (Fabricius, 1792) | 4 | 13 | 26 | 7 | 0 | 50 |
| <i>Cis pygmaeus</i> (Marsham, 1802) | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Cis vestitus</i> Mellié, 1849 | 0 | 11 | 2 | 3 | 1 | 17 |
| <i>Cis villosulus</i> (Marsham, 1802) | 1 | 0 | 3 | 1 | 0 | 5 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|---|----|----|----|---|-----|
| <i>Ennearthron cornutum</i> (Gyllenhal, 1827) | 2 | 1 | 6 | 1 | 0 | 10 |
| <i>Orthocis alni</i> (Gyllenhal, 1813) | 1 | 7 | 7 | 3 | 1 | 19 |
| <i>Sulcacis affinis</i> (Gyllenhal, 1827) | 0 | 4 | 4 | 0 | 0 | 8 |
| <i>Octotemnus glabriculus</i> (Gyllenhal, 1827) | 4 | 11 | 75 | 27 | 4 | 121 |

Tetratomidae [4/4]

| | | | | | | |
|--|---|----|----|----|---|-----|
| <i>Hallomenus binotatus</i> (Quensel, 1790) | 0 | 3 | 15 | 8 | 1 | 27 |
| <i>Tetratoma ancora</i> Fabricius, 1791 | 0 | 2 | 0 | 2 | 0 | 4 |
| <i>Tetratoma desmarestii</i> Latreille, 1807 | 1 | 4 | 3 | 0 | 0 | 8 |
| <i>Tetratoma fungorum</i> Fabricius, 1790 | 9 | 14 | 68 | 16 | 3 | 110 |

Melandryidae [17/11]

| | | | | | | |
|--|---|----|----|---|---|----|
| <i>Orchesia micans</i> (Panzer, 1793) | 4 | 7 | 8 | 6 | 0 | 25 |
| <i>Orchesia minor</i> Walker, 1836 | 0 | 10 | 6 | 0 | 4 | 20 |
| <i>Orchesia undulata</i> Kraatz, 1853 | 1 | 16 | 48 | 7 | 4 | 76 |
| <i>Anisoxya fuscata</i> (Illiger, 1798) | 0 | 2 | 2 | 0 | 0 | 4 |
| <i>Abdera flexuosa</i> (Paykull, 1799) | 0 | 9 | 1 | 1 | 1 | 12 |
| <i>Abdera quadrifasciata</i> (Curtis, 1829) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Phloiotrya vaudoueri</i> Mulsant, 1856 | 1 | 0 | 4 | 0 | 0 | 5 |
| <i>Hypulus quercinus</i> (Quensel, 1790) | 0 | 3 | 2 | 0 | 0 | 5 |
| <i>Melandrya caraboides</i> (Linnaeus, 1761) | 1 | 10 | 9 | 5 | 1 | 26 |
| <i>Conopalpus testaceus</i> (Olivier, 1790) | 0 | 3 | 6 | 1 | 0 | 10 |
| <i>Osphya bipunctata</i> (Fabricius, 1775) | 0 | 1 | 0 | 0 | 0 | 1 |

Mordellidae [17/4]

| | | | | | | |
|--|---|---|----|---|---|----|
| <i>Mordellistena (Mordellistena) neuwaldeggiana</i> (Panzer, 1796) | 0 | 1 | 2 | 1 | 0 | 4 |
| <i>Mordellistena (Mordellistena) pumila</i> (Gyllenhal, 1810) | 2 | 0 | 25 | 1 | 0 | 28 |
| <i>Mordellistena (Mordellistena) variegata</i> (Fabricius, 1798) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Mordellochroa abdominalis</i> (Fabricius, 1775) | 0 | 5 | 2 | 0 | 0 | 7 |

Ripiphoridae [1/1]

| | | | | | | |
|--|---|---|---|---|---|----|
| <i>Metoecus paradoxus</i> (Linnaeus, 1761) | 3 | 9 | 9 | 6 | 2 | 29 |
|--|---|---|---|---|---|----|

Colydiidae [12/5]

| | | | | | | |
|--|---|---|----|---|---|----|
| <i>Orthocerus clavicornis</i> (Linnaeus, 1758) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Synchita humeralis</i> (Fabricius, 1792) | 1 | 0 | 6 | 1 | 0 | 8 |
| <i>Synchita separanda</i> (Reitter, 1882) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Bitoma crenata</i> (Fabricius, 1775) | 6 | 5 | 27 | 6 | 0 | 44 |
| <i>Colydium elongatum</i> (Fabricius, 1787) | 0 | 0 | 1 | 0 | 0 | 1 |

Tenebrionidae [47/31]

| | | | | | | |
|--|----|----|-----|----|----|-----|
| <i>Lagria hirta</i> (Linnaeus, 1758) | 55 | 23 | 123 | 28 | 35 | 264 |
| <i>Eledona agricola</i> (Herbst, 1783) | 5 | 13 | 14 | 2 | 0 | 34 |
| <i>Tenebrio molitor</i> Linnaeus, 1758 | 8 | 6 | 12 | 4 | 1 | 31 |
| <i>Tenebrio obscurus</i> Fabricius, 1792 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Alphitobius diaperinus</i> (Panzer, 1796) | 1 | 1 | 6 | 1 | 1 | 10 |
| <i>Alphitobius laevigatus</i> (Fabricius, 1781) | 1 | 1 | 2 | 0 | 0 | 4 |
| <i>Tribolium castaneum</i> (Herbst, 1797) | 3 | 4 | 8 | 3 | 1 | 19 |
| <i>Tribolium confusum</i> Jacquelin du Val, 1863 | 0 | 1 | 4 | 0 | 0 | 5 |
| <i>Palorus ratzeburgii</i> (Wissmann, 1848) | 1 | 1 | 0 | 1 | 0 | 3 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|----|---|---|----|
| <i>Phylan gibbus</i> (Fabricius, 1775) | 4 | 1 | 0 | 0 | 0 | 5 |
| <i>Melanimon tibialis</i> (Fabricius, 1781) | 0 | 1 | 6 | 1 | 0 | 8 |
| <i>Nalassus laevioctostriatus</i> (Goeze, 1777) | 0 | 3 | 10 | 0 | 0 | 13 |
| <i>Xanthomus pallidus</i> (Curtis, 1830) | 3 | 0 | 0 | 0 | 0 | 3 |
| <i>Blaps mortisaga</i> (Linnaeus, 1758) | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Blaps mucronata</i> Latreille, 1804 | 3 | 11 | 8 | 1 | 1 | 24 |
| <i>Prionychus melanarius</i> (Germar, 1813) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Gonodera luperus</i> (Herbst, 1783) | 0 | 0 | 2 | 2 | 0 | 4 |
| <i>Pseudocistela ceramboides</i> (Linnaeus, 1758) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Isomira murina</i> (Linnaeus, 1758) | 9 | 7 | 12 | 4 | 2 | 34 |
| <i>Mycetochara humeralis</i> (Fabricius, 1787) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Crypticus quisquilius</i> (Linnaeus, 1761) | 14 | 0 | 0 | 0 | 0 | 14 |
| <i>Phaleria cadaverina</i> (Fabricius, 1792) | 5 | 0 | 0 | 0 | 0 | 5 |
| <i>Myrmechixenus subterraneus</i> Chevrolat, 1835 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Myrmechixenus vaporariorum</i> Guérin-Méneville, 1843 | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Corticeus bicolor</i> (Olivier, 1790) | 0 | 5 | 9 | 1 | 0 | 15 |
| <i>Corticeus fraxini</i> (Kugelann, 1794) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Corticeus linearis</i> (Fabricius, 1790) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Corticeus unicolor</i> Piller & Mitterpacher, 1783 | 6 | 0 | 15 | 1 | 1 | 23 |
| <i>Scaphidema metallicum</i> (Fabricius, 1792) | 25 | 11 | 25 | 3 | 0 | 64 |
| <i>Alphitophagus bifasciatus</i> (Say, 1823) | 3 | 1 | 4 | 0 | 0 | 8 |
| <i>Gnatocerus cornutus</i> (Fabricius, 1798) | 2 | 4 | 2 | 0 | 0 | 8 |

Oedemeridae [10/8]

| | | | | | | |
|---|----|----|-----|----|---|-----|
| <i>Nacerdes melanura</i> (Linnaeus, 1758) | 18 | 4 | 7 | 6 | 0 | 35 |
| <i>Ischnomera caerulea</i> (Linnaeus, 1758) | 0 | 4 | 0 | 1 | 0 | 5 |
| <i>Ischnomera cinerascens</i> (Pandellé, 1867) | 0 | 3 | 0 | 0 | 0 | 3 |
| <i>Ischnomera cyanea</i> (Fabricius, 1792) | 0 | 8 | 0 | 1 | 0 | 9 |
| <i>Ischnomera sanguinicollis</i> (Fabricius, 1787) | 0 | 6 | 0 | 0 | 2 | 8 |
| <i>Oedemera (Oedemera) lurida</i> (Marsham, 1802) | 24 | 3 | 140 | 11 | 7 | 185 |
| <i>Oedemera (Oedemera) nobilis</i> (Scopoli, 1763) | 0 | 4 | 6 | 1 | 0 | 11 |
| <i>Oedemera (Oedemera) virescens</i> (Linnaeus, 1767) | 0 | 28 | 0 | 0 | 0 | 28 |

Meloidae [10/2]

| | | | | | | |
|---|---|---|----|---|---|----|
| <i>Meloe proscarabaeus</i> Linnaeus, 1758 | 1 | 8 | 6 | 5 | 2 | 22 |
| <i>Meloe violaceus</i> Marsham, 1802 | 2 | 6 | 20 | 4 | 3 | 35 |

Pyrochroidae [3/2]

| | | | | | | |
|--|----|----|----|----|----|-----|
| <i>Pyrochroa coccinea</i> (Linnaeus, 1761) | 1 | 10 | 8 | 0 | 0 | 19 |
| <i>Pyrochroa serraticornis</i> (Scopoli, 1763) | 38 | 18 | 43 | 26 | 11 | 136 |

Salpingidae [11/9]

| | | | | | | |
|---|----|----|----|----|---|----|
| <i>Aglenus brunneus</i> (Gyllenhal, 1813) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Lissodema cursor</i> (Gyllenhal, 1813) | 0 | 0 | 0 | 2 | 0 | 2 |
| <i>Lissodema denticolle</i> (Gyllenhal, 1813) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Rabocerus foveolatus</i> (Ljungh, 1824) | 0 | 1 | 2 | 0 | 0 | 3 |
| <i>Rabocerus gabrieli</i> Gerhardt, 1901 | 2 | 3 | 9 | 6 | 0 | 20 |
| <i>Sphaeriestes castaneus</i> (Panzer, 1796) | 1 | 8 | 6 | 0 | 1 | 16 |
| <i>Sphaeriestes reyi</i> (Abeille, 1874) | 5 | 4 | 3 | 1 | 1 | 14 |
| <i>Vincenzellus ruficollis</i> (Panzer, 1794) | 10 | 22 | 41 | 20 | 4 | 97 |

Yorkshire Coleoptera checklist 2011

Salpingus planirostris (Fabricius, 1787) 19 49 89 19 5 181

Anthicidae [13/7]

Notoxus monoceros (Linnaeus, 1761) 32 3 0 3 0 38
 Anthicus antherinus (Linnaeus, 1761) 2 0 2 0 0 4
 Cyclodinus constrictus (Curtis, 1838) 1 1 0 0 0 2
 Omonadus floralis (Linnaeus, 1758) 39 15 56 18 3 131
 Omonadus formicarius (Goeze, 1777) 13 8 26 7 5 59
 Cordicomus instabilis (Schmidt, 1842) 0 1 0 0 0 1
 Stricticomus tobias (Marseul, 1879) 1 0 5 0 0 6

Aderidae [3/3]

Aderus populneus (Creutzer in Panzer, 1796) 2 0 8 0 0 10
 Eugenes oculus (Paykull, 1798) 1 10 2 6 0 19
 Vanonus brevicornis (Perris, 1869) 0 0 1 0 0 1

Scraptiidae [14/10]

Anaspis (Anaspis) frontalis (Linnaeus, 1758) 10 36 121 24 10 201
 Anaspis (Anaspis) garneysi Fowler, 1889 3 8 7 5 0 23
 Anaspis (Anaspis) humeralis (Fabricius, 1775) 17 25 64 21 4 131
 Anaspis (Anaspis) lurida Stephens, 1832 0 1 2 3 1 7
 Anaspis (Anaspis) maculata (Geoffrey in Fourcroy, 1785) 71 74 352 56 20 573
 Anaspis (Anaspis) pulicaria Costa, A., 1854 0 4 0 2 0 6
 Anaspis (Anaspis) regimbarti Schilsky, 1895 24 24 112 12 4 176
 Anaspis (Anaspis) thoracica (Linnaeus, 1758) 1 2 6 1 0 10
 Anaspis (Nassipa) costai Emery, 1876 2 3 11 0 0 16
 Anaspis (Nassipa) rufilabris (Gyllenhal, 1827) 13 36 121 16 10 196

Cerambycidae [65/47]

Rhagium (Rhagium) inquisitor (Linnaeus, 1758) 2 4 2 0 0 8
 Rhagium (Hagrium) bifasciatum Fabricius, 1775 2 22 49 8 5 86
 Rhagium (Megarhagium) mordax (De Geer, 1775) 1 22 34 16 4 77
 Stenocorus meridianus (Linnaeus, 1758) 5 13 16 12 3 49
 Grammoptera abdominalis (Stephens, 1831) 1 0 3 1 0 5
 Grammoptera ruficornis (Fabricius, 1781) 38 45 263 45 13 404
 Leptura quadrifasciata Linnaeus, 1758 11 19 31 4 0 65
 Anastrangalia sanguinolenta (Linnaeus, 1761) 0 0 1 0 0 1
 Stictoleptura rubra (Linnaeus, 1758) 1 0 2 0 0 3
 Stictoleptura scutellata (Fabricius, 1781) 0 0 2 0 0 2
 Anoplodera sexguttata (Fabricius, 1775) 0 9 0 0 0 9
 Pachytodes cerambyciformis (Schränk, 1781) 0 0 4 0 0 4
 Alosterna tabacicolor (De Geer, 1775) 0 27 6 2 0 35
 Rutpela maculata (Poda, 1761) 20 14 111 11 0 156
 Stenurella melanura (Linnaeus, 1758) 0 6 2 0 0 8
 Asemum striatum (Linnaeus, 1758) 1 3 2 2 0 8
 Tetropium gabrieli Weise, 1905 0 4 4 2 0 10
 Arhopalus fesus (Mulsant, 1839) 1 0 0 0 0 1
 Arhopalus rusticus (Linnaeus, 1758) 2 0 1 0 0 3
 Cerambyx cerdo Linnaeus, 1758 1 0 0 0 0 1
 Cerambyx scopolii Fuessly, 1775 0 1 2 0 0 3

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|----|----|---|-----|
| Gracilia minuta (Fabricius, 1781) | 2 | 0 | 4 | 2 | 0 | 8 |
| Nathrius brevipennis (Mulsant, 1839) | 0 | 0 | 3 | 1 | 0 | 4 |
| Molorchus minor (Linnaeus, 1758) | 1 | 1 | 0 | 3 | 0 | 5 |
| Glaphyra umbellatarum (von Schreber, 1759) | 0 | 0 | 0 | 2 | 0 | 2 |
| Aromia moschata (Linnaeus, 1758) | 1 | 2 | 2 | 3 | 0 | 8 |
| Hylotrupes bajulus (Linnaeus, 1758) | 0 | 0 | 2 | 0 | 0 | 2 |
| Callidium violaceum (Linnaeus, 1758) | 1 | 0 | 1 | 0 | 0 | 2 |
| Pyrrhidium sanguineum (Linnaeus, 1758) | 0 | 0 | 0 | 1 | 0 | 1 |
| Phymatodes testaceus (Linnaeus, 1758) | 2 | 10 | 1 | 2 | 0 | 15 |
| Poecilium alni (Linnaeus, 1767) | 0 | 0 | 2 | 0 | 0 | 2 |
| Clytus arietis (Linnaeus, 1758) | 5 | 27 | 96 | 11 | 1 | 140 |
| Plagionotus arcuatus (Linnaeus, 1758) | 0 | 0 | 1 | 1 | 0 | 2 |
| Anaglyptus mysticus (Linnaeus, 1758) | 6 | 2 | 13 | 3 | 0 | 24 |
| Mesosa nebulosa (Fabricius, 1781) | 0 | 0 | 2 | 0 | 0 | 2 |
| Agapanthia villosviridescens (De Geer, 1775) | 19 | 0 | 36 | 0 | 1 | 56 |
| Pogonocherus fasciculatus (De Geer, 1775) | 0 | 1 | 0 | 1 | 0 | 2 |
| Pogonocherus hispidulus (Piller & Mitterpacher, 1783) | 4 | 18 | 5 | 3 | 3 | 33 |
| Pogonocherus hispidus (Linnaeus, 1758) | 1 | 10 | 1 | 3 | 1 | 16 |
| Acanthocinus aedilis (Linnaeus, 1758) | 2 | 1 | 6 | 1 | 0 | 10 |
| Leiopus nebulosus (Linnaeus, 1758) | 5 | 22 | 32 | 19 | 0 | 78 |
| Saperda carcharias (Linnaeus, 1758) | 0 | 3 | 7 | 0 | 0 | 10 |
| Saperda populnea (Linnaeus, 1758) | 5 | 6 | 15 | 11 | 0 | 37 |
| Saperda scalaris (Linnaeus, 1758) | 0 | 8 | 5 | 0 | 0 | 13 |
| Stenostola dubia (Laicharting, 1784) | 0 | 6 | 7 | 9 | 3 | 25 |
| Phytoecia cylindrica (Linnaeus, 1758) | 2 | 0 | 0 | 1 | 0 | 3 |
| Tetrops praeustus (Linnaeus, 1758) | 2 | 6 | 22 | 6 | 0 | 36 |

Megalopodidae [3/1]

| | | | | | | |
|---|---|---|----|---|---|----|
| Zeugophora subspinosa (Fabricius, 1781) | 4 | 3 | 16 | 1 | 0 | 24 |
|---|---|---|----|---|---|----|

Orsodacnidae [2/1]

| | | | | | | |
|-----------------------------------|---|----|---|---|---|----|
| Orsodacne cerasi (Linnaeus, 1758) | 0 | 31 | 7 | 8 | 0 | 46 |
|-----------------------------------|---|----|---|---|---|----|

Chrysomelidae [270/211]

| | | | | | | |
|---|----|----|-----|----|---|-----|
| Bruchus loti Paykull, 1800 | 37 | 24 | 134 | 15 | 0 | 210 |
| Bruchus pisorum (Linnaeus, 1758) | 2 | 1 | 2 | 0 | 0 | 5 |
| Bruchus rufimanus Boheman, 1833 | 15 | 8 | 30 | 1 | 0 | 54 |
| Bruchus rufipes Herbst, 1783 | 0 | 0 | 4 | 0 | 0 | 4 |
| Bruchidius cisti (Fabricius, 1775) | 5 | 11 | 9 | 3 | 0 | 28 |
| Bruchidius varius (Olivier, 1795) | 3 | 0 | 2 | 0 | 0 | 5 |
| Bruchidius villosus (Fabricius, 1792) | 4 | 2 | 63 | 9 | 0 | 78 |
| Callosobruchus chinensis (Linnaeus, 1758) | 0 | 1 | 0 | 0 | 0 | 1 |
| Macrolea appendiculata (Panzer, 1794) | 4 | 1 | 0 | 9 | 0 | 14 |
| Macrolea mutica (Fabricius, 1792) | 4 | 0 | 4 | 0 | 0 | 8 |
| Donacia aquatica (Linnaeus, 1758) | 0 | 1 | 1 | 1 | 0 | 3 |
| Donacia bicolora Zschach, 1788 | 0 | 2 | 0 | 0 | 0 | 2 |
| Donacia cinerea Herbst, 1784 | 7 | 3 | 6 | 1 | 0 | 17 |
| Donacia clavipes Fabricius, 1792 | 12 | 2 | 3 | 3 | 0 | 20 |
| Donacia crassipes Fabricius, 1775 | 0 | 0 | 1 | 0 | 0 | 1 |
| Donacia dentata Hoppe, 1795 | 1 | 0 | 0 | 0 | 0 | 1 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|---|-----|
| <i>Donacia impressa</i> Paykull, 1799 | 5 | 1 | 1 | 0 | 0 | 7 |
| <i>Donacia marginata</i> Hoppe, 1795 | 1 | 1 | 1 | 5 | 0 | 8 |
| <i>Donacia semicuprea</i> Panzer, 1796 | 12 | 2 | 21 | 1 | 0 | 36 |
| <i>Donacia simplex</i> Fabricius, 1775 | 18 | 11 | 36 | 13 | 2 | 80 |
| <i>Donacia sparganii</i> Ahrens, 1810 | 7 | 0 | 0 | 2 | 0 | 9 |
| <i>Donacia thalassina</i> Germar, 1811 | 4 | 6 | 4 | 0 | 0 | 14 |
| <i>Donacia versicolore</i> a (Brahm, 1791) | 14 | 5 | 15 | 11 | 0 | 45 |
| <i>Donacia vulgaris</i> Zschach, 1788 | 14 | 9 | 20 | 6 | 0 | 49 |
| <i>Plateumaris affinis</i> (Kunze, 1818) | 0 | 3 | 2 | 2 | 4 | 11 |
| <i>Plateumaris braccata</i> (Scopoli, 1772) | 4 | 0 | 0 | 0 | 0 | 4 |
| <i>Plateumaris discolor</i> (Panzer, 1795) | 0 | 21 | 21 | 14 | 5 | 61 |
| <i>Plateumaris sericea</i> (Linnaeus, 1758) | 16 | 12 | 17 | 8 | 1 | 54 |
| <i>Lema cyanella</i> (Linnaeus, 1758) | 4 | 6 | 2 | 2 | 0 | 14 |
| <i>Oulema melanopus</i> (Linnaeus, 1758) | 6 | 0 | 5 | 4 | 1 | 16 |
| <i>Oulema obscura</i> (Stephens, 1831) | 13 | 30 | 11 | 17 | 1 | 72 |
| <i>Oulema rufocyanea</i> (Suffrian, 1847) | 37 | 11 | 48 | 14 | 6 | 116 |
| <i>Crioceris asparagi</i> (Linnaeus, 1758) | 0 | 0 | 20 | 0 | 0 | 20 |
| <i>Lilioceris lili</i> (Scopoli, 1763) | 7 | 0 | 9 | 3 | 0 | 19 |
| <i>Labidostomis tridentata</i> (Linnaeus, 1758) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Clytra quadripunctata</i> (Linnaeus, 1758) | 0 | 12 | 16 | 0 | 0 | 28 |
| <i>Cryptocephalus aureolus</i> Suffrian, 1847 | 2 | 11 | 0 | 0 | 0 | 13 |
| <i>Cryptocephalus biguttatus</i> (Scopoli, 1763) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Cryptocephalus bipunctatus</i> (Linnaeus, 1758) | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Cryptocephalus fulvus</i> (Goeze, 1777) | 2 | 2 | 20 | 2 | 0 | 26 |
| <i>Cryptocephalus hypochaeridis</i> (Linnaeus, 1758) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Cryptocephalus labiatus</i> (Linnaeus, 1761) | 4 | 13 | 85 | 12 | 0 | 114 |
| <i>Cryptocephalus moraei</i> (Linnaeus, 1758) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Cryptocephalus parvulus</i> Müller, O.F., 1776 | 0 | 0 | 8 | 0 | 0 | 8 |
| <i>Cryptocephalus punctiger</i> Paykull, 1799 | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Cryptocephalus pusillus</i> Fabricius, 1777 | 13 | 10 | 34 | 6 | 0 | 63 |
| <i>Oomorplus concolor</i> (Sturm, 1807) | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Timarcha goettingensis</i> (Linnaeus, 1758) | 2 | 1 | 27 | 8 | 1 | 39 |
| <i>Timarcha tenebricosa</i> (Fabricius, 1775) | 0 | 0 | 9 | 3 | 0 | 12 |
| <i>Leptinotarsa decemlineata</i> (Say, 1824) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Chrysolina americana</i> (Linnaeus, 1758) | 2 | 0 | 0 | 0 | 0 | 2 |
| <i>Chrysolina banksi</i> (Fabricius, 1775) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Chrysolina brunsvicensis</i> (Gravenhorst, 1807) | 0 | 5 | 0 | 3 | 2 | 10 |
| <i>Chrysolina fastuosa</i> (Scopoli, 1763) | 0 | 0 | 28 | 0 | 0 | 28 |
| <i>Chrysolina graminis</i> (Linnaeus, 1758) | 5 | 12 | 1 | 15 | 0 | 33 |
| <i>Chrysolina herbacea</i> (Duftschmid, 1825) | 0 | 0 | 1 | 1 | 0 | 2 |
| <i>Chrysolina hyperici</i> (Forster, 1771) | 2 | 6 | 23 | 2 | 0 | 33 |
| <i>Chrysolina marginata</i> (Linnaeus, 1758) | 1 | 1 | 3 | 0 | 2 | 7 |
| <i>Chrysolina oricalcia</i> (Müller, O.F., 1776) | 3 | 1 | 10 | 5 | 5 | 24 |
| <i>Chrysolina polita</i> (Linnaeus, 1758) | 29 | 27 | 96 | 26 | 5 | 183 |
| <i>Chrysolina sanguinolenta</i> (Linnaeus, 1758) | 1 | 0 | 7 | 0 | 0 | 8 |
| <i>Chrysolina staphylaea</i> (Linnaeus, 1758) | 10 | 13 | 29 | 11 | 4 | 67 |
| <i>Chrysolina varians</i> (Schaller, 1783) | 2 | 13 | 3 | 4 | 2 | 24 |
| <i>Chrysolina violacea</i> (Müller, O.F., 1776) | 0 | 1 | 3 | 2 | 0 | 6 |
| <i>Gastrophysa polygoni</i> (Linnaeus, 1758) | 20 | 10 | 50 | 31 | 2 | 113 |
| <i>Gastrophysa viridula</i> (De Geer, 1775) | 28 | 20 | 143 | 32 | 5 | 228 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| <i>Phaedon armoraciae</i> (Linnaeus, 1758) | 21 | 22 | 56 | 16 | 3 | 118 |
| <i>Phaedon cochleariae</i> (Fabricius, 1792) | 27 | 12 | 43 | 18 | 10 | 110 |
| <i>Phaedon concinnus</i> Stephens, 1831 | 10 | 0 | 0 | 0 | 0 | 10 |
| <i>Phaedon tumidulus</i> (Germar, 1824) | 53 | 32 | 177 | 26 | 4 | 292 |
| <i>Hydrothassa glabra</i> (Herbst, 1783) | 3 | 5 | 3 | 9 | 1 | 21 |
| <i>Hydrothassa hannoveriana</i> (Fabricius, 1775) | 0 | 0 | 0 | 9 | 2 | 11 |
| <i>Hydrothassa marginella</i> (Linnaeus, 1758) | 15 | 35 | 42 | 29 | 5 | 126 |
| <i>Prasocuris junci</i> (Brahm, 1790) | 11 | 9 | 6 | 5 | 1 | 32 |
| <i>Prasocuris phellandrii</i> (Linnaeus, 1758) | 5 | 2 | 2 | 3 | 1 | 13 |
| <i>Plagioderia versicolora</i> (Laicharting, 1781) | 2 | 3 | 4 | 0 | 0 | 9 |
| <i>Chrysomela aenea</i> Linnaeus, 1758 | 1 | 39 | 9 | 9 | 5 | 63 |
| <i>Chrysomela populi</i> Linnaeus, 1758 | 10 | 10 | 7 | 1 | 0 | 28 |
| <i>Gonioctena decemnotata</i> (Marsham, 1802) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Gonioctena olivacea</i> (Forster, 1771) | 0 | 4 | 7 | 0 | 1 | 12 |
| <i>Gonioctena pallida</i> (Linnaeus, 1758) | 0 | 33 | 2 | 9 | 4 | 48 |
| <i>Gonioctena viminalis</i> (Linnaeus, 1758) | 0 | 3 | 0 | 0 | 0 | 3 |
| <i>Phratora laticollis</i> Suffrian, 1851 | 12 | 6 | 11 | 3 | 0 | 32 |
| <i>Phratora vitellinae</i> (Linnaeus, 1758) | 14 | 5 | 31 | 29 | 12 | 91 |
| <i>Phratora vulgatissima</i> (Linnaeus, 1758) | 51 | 13 | 110 | 23 | 18 | 215 |
| <i>Galerucella calmariensis</i> (Linnaeus, 1767) | 9 | 3 | 20 | 5 | 0 | 37 |
| <i>Galerucella lineola</i> (Fabricius, 1781) | 9 | 12 | 73 | 15 | 7 | 116 |
| <i>Galerucella nymphaeae</i> (Linnaeus, 1758) | 7 | 6 | 70 | 8 | 0 | 91 |
| <i>Galerucella pusilla</i> (Duftschmid, 1825) | 2 | 0 | 9 | 6 | 0 | 17 |
| <i>Galerucella sagittariae</i> (Gyllenhal, 1813) | 27 | 6 | 17 | 8 | 0 | 58 |
| <i>Galerucella tenella</i> (Linnaeus, 1761) | 14 | 17 | 16 | 29 | 2 | 78 |
| <i>Pyrrhalta viburni</i> (Paykull, 1799) | 1 | 6 | 23 | 4 | 0 | 34 |
| <i>Xanthogaleruca luteola</i> (Müller, O.F., 1766) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Galeruca tanacetii</i> (Linnaeus, 1758) | 8 | 3 | 16 | 3 | 0 | 30 |
| <i>Lochmaea caprea</i> (Linnaeus, 1758) | 7 | 9 | 100 | 14 | 1 | 131 |
| <i>Lochmaea crataegi</i> (Forster, 1771) | 6 | 19 | 79 | 14 | 5 | 123 |
| <i>Lochmaea suturalis</i> (Thomson, C.G., 1866) | 9 | 27 | 129 | 7 | 6 | 178 |
| <i>Phyllotreta quadrimaculata</i> (Linnaeus, 1758) | 2 | 7 | 12 | 12 | 0 | 33 |
| <i>Luperus flavipes</i> (Linnaeus, 1767) | 1 | 6 | 15 | 0 | 0 | 22 |
| <i>Luperus longicornis</i> (Fabricius, 1781) | 8 | 21 | 13 | 16 | 5 | 63 |
| <i>Calomicrus circumfusus</i> (Marsham, 1802) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Agelastica alni</i> (Linnaeus, 1758) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Sermylassa halensis</i> (Linnaeus, 1767) | 20 | 12 | 29 | 2 | 2 | 65 |
| <i>Phyllotreta atra</i> (Fabricius, 1775) | 3 | 9 | 6 | 0 | 0 | 18 |
| <i>Phyllotreta consobrina</i> (Curtis, 1837) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Phyllotreta cruciferae</i> (Goeze, 1777) | 0 | 3 | 0 | 0 | 0 | 3 |
| <i>Phyllotreta diademata</i> Foudras, 1860 | 0 | 0 | 5 | 0 | 0 | 5 |
| <i>Phyllotreta exclamationis</i> (Thunberg, 1784) | 9 | 3 | 7 | 4 | 1 | 24 |
| <i>Phyllotreta flexuosa</i> (Illiger, 1794) | 2 | 3 | 2 | 10 | 2 | 19 |
| <i>Phyllotreta nemorum</i> (Linnaeus, 1758) | 4 | 2 | 10 | 14 | 7 | 37 |
| <i>Phyllotreta nigripes</i> (Fabricius, 1775) | 8 | 6 | 42 | 6 | 0 | 62 |
| <i>Phyllotreta nodicornis</i> (Marsham, 1802) | 4 | 1 | 19 | 1 | 1 | 26 |
| <i>Phyllotreta ochripes</i> (Curtis, 1837) | 8 | 3 | 3 | 0 | 0 | 14 |
| <i>Phyllotreta punctulata</i> (Marsham, 1802) | 1 | 1 | 1 | 0 | 0 | 3 |
| <i>Phyllotreta tetrastigma</i> (Comolli, 1837) | 12 | 6 | 26 | 15 | 1 | 60 |
| <i>Phyllotreta undulata</i> Kutschera, 1860 | 29 | 39 | 78 | 26 | 4 | 176 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Phyllotreta vittula</i> (Redtenbacher, 1849) | 1 | 3 | 7 | 3 | 0 | 14 |
| <i>Aphthona ?atratula</i> Allard, 1859 | 1 | 1 | 9 | 1 | 0 | 12 |
| <i>Aphthona atrocaerulea</i> (Stephens, 1829) | 5 | 3 | 15 | 1 | 0 | 24 |
| <i>Aphthona euphorbiae</i> (Schrank, 1781) | 71 | 43 | 264 | 25 | 4 | 407 |
| <i>Aphthona herbigrada</i> (Curtis, 1837) | 3 | 7 | 0 | 5 | 1 | 16 |
| <i>Aphthona lutescens</i> (Gyllenhal, 1808) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Aphthona melancholica</i> Weise, 1888 | 1 | 1 | 8 | 2 | 0 | 12 |
| <i>Aphthona nigriceps</i> (Redtenbacher, 1842) | 1 | 2 | 1 | 10 | 0 | 14 |
| <i>Aphthona nonstriata</i> (Goeze, 1777) | 9 | 7 | 21 | 32 | 1 | 70 |
| <i>Longitarsus aeneicollis</i> (Faldermann, 1837) | 2 | 2 | 3 | 1 | 0 | 8 |
| <i>Longitarsus anchusae</i> (Paykull, 1799) | 0 | 2 | 5 | 0 | 0 | 7 |
| <i>Longitarsus atricillus</i> (Linnaeus, 1761) | 3 | 5 | 5 | 0 | 1 | 14 |
| <i>Longitarsus brunneus</i> (Duftschmid, 1825) | 0 | 2 | 2 | 3 | 0 | 7 |
| <i>Longitarsus dorsalis</i> (Fabricius, 1781) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Longitarsus exoletus</i> (Linnaeus, 1758) | 3 | 1 | 11 | 1 | 0 | 16 |
| <i>Longitarsus flavicornis</i> (Stephens, 1831) | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Longitarsus ganglbaueri</i> Heikertinger, 1912 | 1 | 3 | 2 | 0 | 2 | 8 |
| <i>Longitarsus gracilis</i> Kutschera, 1864 | 21 | 9 | 10 | 5 | 2 | 47 |
| <i>Longitarsus holsaticus</i> (Linnaeus, 1758) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Longitarsus jacobaeae</i> (Waterhouse, G.R., 1858) | 26 | 8 | 63 | 10 | 6 | 113 |
| <i>Longitarsus kutscherae</i> (Rye, 1872) | 4 | 2 | 16 | 0 | 0 | 22 |
| <i>Longitarsus luridus</i> (Scopoli, 1763) | 59 | 34 | 199 | 43 | 7 | 342 |
| <i>Longitarsus melanocephalus</i> (De Geer, 1775) | 11 | 5 | 44 | 8 | 0 | 68 |
| <i>Longitarsus membranaceus</i> (Foudras, 1860) | 2 | 5 | 0 | 0 | 4 | 11 |
| <i>Longitarsus nasturtii</i> (Fabricius, 1792) | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Longitarsus nigerrimus</i> (Gyllenhal, 1827) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Longitarsus nigrofasciatus</i> (Goeze, 1777) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Longitarsus obliteratus</i> (Rosenhauer, 1847) | 2 | 0 | 1 | 0 | 0 | 3 |
| <i>Longitarsus ochroleucus</i> (Marsham, 1802) | 3 | 2 | 7 | 2 | 0 | 14 |
| <i>Longitarsus parvulus</i> (Paykull, 1799) | 22 | 18 | 49 | 5 | 0 | 94 |
| <i>Longitarsus pellucidus</i> (Foudras, 1860) | 2 | 1 | 7 | 0 | 0 | 10 |
| <i>Longitarsus pratensis</i> (Panzer, 1794) | 20 | 5 | 9 | 4 | 0 | 38 |
| <i>Longitarsus reichei</i> (Allard, 1860) | 3 | 0 | 11 | 0 | 0 | 14 |
| <i>Longitarsus rubiginosus</i> (Foudras, 1860) | 6 | 0 | 17 | 0 | 0 | 23 |
| <i>Longitarsus succineus</i> (Foudras, 1860) | 15 | 7 | 41 | 5 | 3 | 71 |
| <i>Longitarsus suturellus</i> (Duftschmid, 1825) | 25 | 14 | 89 | 22 | 2 | 152 |
| <i>Longitarsus tabidus</i> (Fabricius, 1775) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Altica brevicollis</i> Foudras, 1860 | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Altica ericeti</i> (Allard, 1859) | 1 | 10 | 13 | 2 | 3 | 29 |
| <i>Altica helianthemi</i> (Allard, 1859) | 3 | 5 | 10 | 4 | 0 | 22 |
| <i>Altica lythri</i> Aubé, 1843 | 58 | 44 | 262 | 34 | 16 | 414 |
| <i>Altica oleracea</i> (Linnaeus, 1758) | 12 | 10 | 45 | 7 | 0 | 74 |
| <i>Altica palustris</i> Weise, 1888 | 16 | 11 | 69 | 13 | 2 | 111 |
| <i>Hermæophaga mercurialis</i> (Fabricius, 1792) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Batophila aerata</i> (Marsham, 1802) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Batophila rubi</i> (Paykull, 1799) | 3 | 23 | 43 | 11 | 3 | 83 |
| <i>Lythriaria salicariae</i> (Paykull, 1800) | 0 | 1 | 2 | 0 | 0 | 3 |
| <i>Ochrosis ventralis</i> (Illiger, 1807) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Neocrepidodera ferruginea</i> (Scopoli, 1763) | 63 | 29 | 190 | 36 | 10 | 328 |
| <i>Neocrepidodera impressa</i> (Fabricius, 1801) | 1 | 0 | 0 | 0 | 0 | 1 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Neocrepidodera transversa</i> (Marsham, 1802) | 15 | 23 | 121 | 21 | 5 | 185 |
| <i>Derocrepis rufipes</i> (Linnaeus, 1758) | 9 | 17 | 23 | 15 | 6 | 70 |
| <i>Hippuriphila modeeri</i> (Linnaeus, 1761) | 13 | 12 | 14 | 5 | 4 | 48 |
| <i>Crepidodera aurata</i> (Marsham, 1802) | 14 | 4 | 11 | 4 | 0 | 33 |
| <i>Crepidodera aurea</i> (Fourcroy, 1785) | 0 | 2 | 12 | 1 | 1 | 16 |
| <i>Crepidodera fulvicornis</i> (Fabricius, 1792) | 86 | 40 | 320 | 52 | 23 | 521 |
| <i>Crepidodera plutus</i> (Latreille, 1804) | 2 | 0 | 9 | 0 | 0 | 11 |
| <i>Epitrix atropae</i> Foudras, 1860 | 2 | 0 | 1 | 0 | 0 | 3 |
| <i>Epitrix pubescens</i> (Koch, J.D.W., 1803) | 3 | 0 | 13 | 2 | 0 | 18 |
| <i>Podagrica fuscicornis</i> (Linnaeus, 1767) | 0 | 3 | 0 | 0 | 0 | 3 |
| <i>Mantura matthewsii</i> (Curtis, 1833) | 4 | 5 | 0 | 8 | 0 | 17 |
| <i>Mantura obtusata</i> (Gyllenhal, 1813) | 0 | 4 | 1 | 7 | 0 | 12 |
| <i>Mantura rustica</i> (Linnaeus, 1767) | 5 | 6 | 13 | 8 | 0 | 32 |
| <i>Chaetocnema arida</i> Foudras, 1860 | 1 | 2 | 0 | 0 | 0 | 3 |
| <i>Chaetocnema concinna</i> (Marsham, 1802) | 49 | 29 | 216 | 46 | 8 | 348 |
| <i>Chaetocnema confusa</i> (Boheman, 1851) | 3 | 0 | 1 | 0 | 0 | 4 |
| <i>Chaetocnema hortensis</i> (Fourcroy, 1785) | 15 | 11 | 45 | 19 | 8 | 98 |
| <i>Chaetocnema sahlbergii</i> (Gyllenhal, 1827) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Chaetocnema subcoerulea</i> (Kutschera, 1864) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Sphaeroderma rubidum</i> (Graëlls, 1858) | 16 | 12 | 27 | 2 | 1 | 58 |
| <i>Sphaeroderma testaceum</i> (Fabricius, 1775) | 28 | 16 | 79 | 19 | 5 | 147 |
| <i>Apteropeda globosa</i> (Illiger, 1794) | 0 | 3 | 1 | 2 | 0 | 6 |
| <i>Apteropeda orbiculata</i> (Marsham, 1802) | 2 | 7 | 7 | 1 | 1 | 18 |
| <i>Mniophila muscorum</i> (Koch, J.D.W., 1803) | 0 | 3 | 0 | 3 | 1 | 7 |
| <i>Psylliodes affinis</i> (Paykull, 1799) | 32 | 4 | 111 | 23 | 1 | 171 |
| <i>Psylliodes attenuata</i> (Koch, J.D.W., 1803) | 0 | 0 | 3 | 1 | 0 | 4 |
| <i>Psylliodes chalcomera</i> (Illiger, 1807) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Psylliodes chrysocephala</i> (Linnaeus, 1758) | 20 | 15 | 149 | 7 | 8 | 199 |
| <i>Psylliodes cuprea</i> (Koch, J.D.W., 1803) | 3 | 10 | 12 | 1 | 1 | 27 |
| <i>Psylliodes dulcamarae</i> (Koch, J.D.W., 1803) | 1 | 0 | 5 | 0 | 0 | 6 |
| <i>Psylliodes hyoscyami</i> (Linnaeus, 1758) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Psylliodes laticollis</i> Kutschera, 1864 | 11 | 4 | 15 | 0 | 0 | 30 |
| <i>Psylliodes marcida</i> (Illiger, 1807) | 7 | 3 | 0 | 0 | 0 | 10 |
| <i>Psylliodes napi</i> (Fabricius, 1792) | 8 | 10 | 26 | 13 | 6 | 63 |
| <i>Psylliodes picina</i> (Marsham, 1802) | 15 | 6 | 29 | 9 | 0 | 59 |
| <i>Cassida flaveola</i> Thunberg, 1794 | 4 | 12 | 27 | 14 | 1 | 58 |
| <i>Cassida hemisphaerica</i> Herbst, 1799 | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Cassida nobilis</i> Linnaeus, 1758 | 0 | 1 | 2 | 0 | 0 | 3 |
| <i>Cassida prasina</i> Illiger, 1798 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Cassida rubiginosa</i> Müller, O.F., 1776 | 44 | 22 | 155 | 31 | 13 | 265 |
| <i>Cassida vibex</i> Linnaeus, 1767 | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Cassida viridis</i> Linnaeus, 1758 | 5 | 4 | 12 | 3 | 0 | 24 |
| <u>Nemonychidae [1/1]</u> | | | | | | |
| <i>Cimberis attelaboides</i> (Fabricius, 1787) | 0 | 9 | 0 | 1 | 0 | 10 |

Anthribidae [9/7]

| | | | | | | |
|---|---|----|----|----|---|----|
| <i>Anthribus fasciatus</i> (Forster, 1771) | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Anthribus nebulosus</i> (Forster, 1771) | 4 | 14 | 7 | 10 | 2 | 37 |
| <i>Platyrhinus resinosus</i> (Scopoli, 1763) | 5 | 1 | 22 | 2 | 0 | 30 |
| <i>Platystomos albinus</i> (Linnaeus, 1758) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Choragus sheppardi</i> Kirby, 1819 | 3 | 1 | 1 | 1 | 0 | 6 |
| <i>Araecerus fasciculatus</i> (De Geer, 1775) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Bruchela rufipes</i> (Olivier, 1790) | 2 | 0 | 15 | 0 | 0 | 17 |

Rhynchitidae [18/14]

| | | | | | | |
|--|----|----|----|----|---|-----|
| <i>Involvlus cupreus</i> (Linnaeus, 1758) | 0 | 21 | 1 | 0 | 0 | 22 |
| <i>Lasiorrhynchites</i> (<i>Lasiorrhynchites</i>) <i>cavifrons</i> (Gyllenhal, 1833) | 0 | 2 | 4 | 2 | 0 | 8 |
| <i>Lasiorrhynchites</i> (<i>Lasiorrhynchites</i>) <i>olivaceus</i> (Gyllenhal, 1833) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Neocoenorrhinus aeneovirens</i> (Marsham, 1802) | 0 | 12 | 3 | 0 | 0 | 15 |
| <i>Neocoenorrhinus aequatus</i> (Linnaeus, 1767) | 5 | 5 | 37 | 1 | 0 | 48 |
| <i>Neocoenorrhinus germanicus</i> (Herbst, 1797) | 11 | 8 | 21 | 3 | 0 | 43 |
| <i>Neocoenorrhinus interpunctatus</i> (Stephens, 1831) | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Neocoenorrhinus pauxillus</i> (Germar, 1824) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Temnocerus longiceps</i> (Thomson, C.G., 1888) | 2 | 4 | 13 | 6 | 0 | 25 |
| <i>Temnocerus nanus</i> (Paykull, 1792) | 13 | 10 | 49 | 12 | 0 | 84 |
| <i>Temnocerus tomentosus</i> (Gyllenhal, 1839) | 1 | 1 | 5 | 1 | 0 | 8 |
| <i>Byctiscus betulae</i> (Linnaeus, 1758) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Deporaus</i> (<i>Deporaus</i>) <i>betulae</i> (Linnaeus, 1758) | 9 | 20 | 67 | 10 | 0 | 106 |
| <i>Deporaus</i> (<i>Caenorhinus</i>) <i>mannerheimii</i> (Hummel, 1823) | 8 | 15 | 10 | 2 | 0 | 35 |

Attelabidae [2/2]

| | | | | | | |
|---|---|----|----|---|---|----|
| <i>Attelabus nitens</i> (Scopoli, 1763) | 9 | 14 | 50 | 6 | 0 | 79 |
| <i>Apoderus coryli</i> (Linnaeus, 1758) | 0 | 3 | 1 | 1 | 0 | 5 |

Apionidae [87/68]

| | | | | | | |
|--|----|----|-----|----|----|-----|
| <i>Apion cruentatum</i> Walton, 1844 | 7 | 5 | 10 | 3 | 1 | 26 |
| <i>Apion frumentarium</i> (Linnaeus, 1758) | 41 | 10 | 88 | 15 | 6 | 160 |
| <i>Apion haematodes</i> Kirby, 1808 | 14 | 6 | 33 | 6 | 3 | 62 |
| <i>Apion rubens</i> Stephens, 1839 | 2 | 3 | 1 | 2 | 1 | 9 |
| <i>Apion rubiginosum</i> Grill, 1893 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Perapion</i> (<i>Perapion</i>) <i>affine</i> (Kirby, 1808) | 2 | 3 | 3 | 1 | 0 | 9 |
| <i>Perapion</i> (<i>Perapion</i>) <i>curtirostre</i> (Germar, 1817) | 28 | 22 | 103 | 20 | 6 | 179 |
| <i>Perapion</i> (<i>Perapion</i>) <i>hydrolapathi</i> (Marsham, 1802) | 19 | 14 | 33 | 9 | 0 | 75 |
| <i>Perapion</i> (<i>Perapion</i>) <i>marchicum</i> (Herbst, 1797) | 10 | 5 | 25 | 2 | 0 | 42 |
| <i>Perapion</i> (<i>Perapion</i>) <i>violaceum</i> (Kirby, 1808) | 30 | 18 | 115 | 28 | 18 | 209 |
| <i>Aspidapion</i> (<i>Aspidapion</i>) <i>radiolus</i> (Marsham, 1802) | 16 | 9 | 20 | 2 | 2 | 49 |
| <i>Aspidapion</i> (<i>Koestlinia</i>) <i>aeneum</i> (Fabricius, 1775) | 7 | 6 | 18 | 0 | 1 | 32 |
| <i>Ceratapion</i> (<i>Ceratapion</i>) <i>carduorum</i> (Kirby, 1808) | 9 | 7 | 26 | 6 | 0 | 48 |
| <i>Ceratapion</i> (<i>Ceratapion</i>) <i>gibbirostre</i> (Gyllenhal, 1813) | 31 | 1 | 47 | 9 | 5 | 93 |
| <i>Ceratapion</i> (<i>Acanephodus</i>) <i>onopordi</i> (Kirby, 1808) | 19 | 3 | 8 | 1 | 2 | 33 |
| <i>Diplapion confluens</i> (Kirby, 1808) | 7 | 0 | 1 | 0 | 0 | 8 |
| <i>Diplapion stolidum</i> (Germar, 1817) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Omphalapion hookerorum</i> (Kirby, 1808) | 13 | 2 | 13 | 3 | 1 | 32 |
| <i>Omphalapion laevigatum</i> (Paykull, 1792) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Exapion</i> (<i>Exapion</i>) <i>fuscirostre</i> (Fabricius, 1775) | 1 | 0 | 8 | 0 | 0 | 9 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| Exapion (Ulapion) ulicis (Forster, 1771) | 54 | 14 | 77 | 9 | 9 | 163 |
| Kalcapion pallipes (Kirby, 1808) | 5 | 15 | 5 | 10 | 0 | 35 |
| Melanapion minimum (Herbst, 1797) | 0 | 4 | 1 | 0 | 0 | 5 |
| Squamapion atomarium (Kirby, W., 1808) | 3 | 1 | 0 | 0 | 0 | 4 |
| Squamapion vicinum (Kirby, W., 1808) | 0 | 0 | 1 | 1 | 0 | 2 |
| Malvapion malvae (Fabricius, 1775) | 15 | 0 | 31 | 2 | 0 | 48 |
| Pseudapion rufirostre (Fabricius, 1775) | 4 | 0 | 9 | 0 | 1 | 14 |
| Rhopalapion longirostre (Olivier, 1807) | 0 | 0 | 0 | 1 | 0 | 1 |
| Cyanapion (Cyanapion) spencii (Kirby, 1808) | 1 | 3 | 10 | 5 | 1 | 20 |
| Cyanapion (Bothryorrhynchapion) afer (Gyllenhal, 1833) | 3 | 2 | 8 | 3 | 1 | 17 |
| Cyanapion (Bothryorrhynchapion) gyllenhalii (Kirby, 1808) | 1 | 2 | 1 | 1 | 0 | 5 |
| Eutrichapion (Eutrichapion) ervi (Kirby, 1808) | 30 | 19 | 76 | 18 | 5 | 148 |
| Eutrichapion (Eutrichapion) viciae (Paykull, 1800) | 12 | 7 | 61 | 14 | 2 | 96 |
| Eutrichapion (Cnemapion) vorax (Herbst, 1797) | 9 | 5 | 19 | 5 | 0 | 38 |
| Eutrichapion (Psilocalymma) punctigerum (Paykull, 1792) | 1 | 6 | 4 | 1 | 2 | 14 |
| Hemitrichapion (Tinocyba) reflexum (Gyllenhal, 1833) | 0 | 1 | 1 | 0 | 0 | 2 |
| Holotrichapion (Holotrichapion) ononis (Kirby, 1808) | 20 | 4 | 2 | 3 | 1 | 30 |
| Holotrichapion (Apiops) pisi (Fabricius, 1801) | 10 | 6 | 15 | 6 | 2 | 39 |
| Holotrichapion (Legaricapion) aethiops (Herbst, 1797) | 5 | 3 | 4 | 2 | 2 | 16 |
| Oxystoma cerdo (Gerstaecker, 1854) | 6 | 0 | 15 | 3 | 0 | 24 |
| Oxystoma craccae (Linnaeus, 1767) | 1 | 0 | 1 | 1 | 0 | 3 |
| Oxystoma pomonae (Fabricius, 1798) | 0 | 0 | 1 | 0 | 0 | 1 |
| Oxystoma subulatum (Kirby, 1808) | 21 | 31 | 58 | 16 | 2 | 128 |
| Pirapion immune (Kirby, W., 1808) | 2 | 3 | 6 | 0 | 0 | 11 |
| Catapion curtisi (Stephens, 1831) | 1 | 0 | 0 | 0 | 0 | 1 |
| Catapion pubescens (Kirby, W., 1811) | 2 | 1 | 5 | 1 | 1 | 10 |
| Catapion seniculus (Kirby, W., 1808) | 15 | 6 | 3 | 5 | 4 | 33 |
| Ischnopterapion (Ischnopterapion) loti (Kirby, 1808) | 39 | 29 | 88 | 32 | 4 | 192 |
| Ischnopterapion (Ischnopterapion) modestum (Germar, 1817) | 0 | 0 | 3 | 1 | 0 | 4 |
| Ischnopterapion (Chlorapion) virens (Herbst, 1797) | 25 | 12 | 19 | 8 | 2 | 66 |
| Protopirapion atratum (Germar, 1817) | 2 | 5 | 6 | 0 | 3 | 16 |
| Stenopterapion (Stenopterapion) meliloti (Kirby, 1808) | 10 | 3 | 7 | 0 | 0 | 20 |
| Stenopterapion (Stenopterapion) tenue (Kirby, 1808) | 13 | 2 | 10 | 4 | 0 | 29 |
| Stenopterapion (Cobosiotherium) scutellare (Kirby, 1811) | 0 | 1 | 4 | 0 | 0 | 5 |
| Synapion ebeninum (Kirby, 1808) | 0 | 0 | 0 | 1 | 0 | 1 |
| Betulapion simile (Kirby, 1811) | 21 | 4 | 66 | 10 | 0 | 101 |
| Protapion apricans (Herbst, 1797) | 39 | 21 | 154 | 28 | 15 | 257 |
| Protapion assimile (Kirby, 1808) | 18 | 36 | 43 | 5 | 1 | 103 |
| Protapion difforme (Germar, 1818) | 0 | 0 | 1 | 0 | 0 | 1 |
| Protapion dissimile (Germar, 1817) | 0 | 0 | 2 | 0 | 0 | 2 |
| Protapion filirostre (Kirby, 1808) | 0 | 0 | 3 | 0 | 0 | 3 |
| Protapion fulvipes (Geoffroy in Fourcroy, 1785) | 67 | 51 | 163 | 35 | 4 | 320 |
| Protapion nigrirtarse (Kirby, 1808) | 21 | 13 | 61 | 13 | 4 | 112 |
| Protapion ononidis (Gyllenhal, 1827) | 6 | 2 | 6 | 4 | 0 | 18 |
| Protapion schoenherri (Boheman, 1839) | 0 | 1 | 0 | 1 | 0 | 2 |
| Protapion trifolii (Linnaeus, 1768) | 13 | 3 | 14 | 1 | 2 | 33 |
| Protapion varipes (Germar, 1817) | 0 | 0 | 2 | 0 | 0 | 2 |
| Pseudoprotapion astragali (Paykull, 1800) | 0 | 0 | 0 | 1 | 0 | 1 |

Nanophyidae [2/1]

| | | | | | | |
|------------------------------------|----|---|----|---|---|----|
| Nanophyes marmoratus (Goeze, 1777) | 11 | 4 | 33 | 7 | 0 | 55 |
|------------------------------------|----|---|----|---|---|----|

Dryophthoridae [4/4]

| | | | | | | |
|---|---|---|---|---|---|----|
| Dryophthorus corticalis (Paykull, 1792) | 0 | 0 | 2 | 0 | 0 | 2 |
| Sitophilus granarius (Linnaeus, 1758) | 3 | 4 | 5 | 1 | 0 | 13 |
| Sitophilus oryzae (Linnaeus, 1763) | 2 | 5 | 4 | 2 | 0 | 13 |
| Sitophilus zeamais Motschulsky, 1855 | 0 | 0 | 1 | 0 | 0 | 1 |

Erihniidae [12/9]

| | | | | | | |
|---|----|----|----|----|---|-----|
| Grypus equiseti (Fabricius, 1775) | 11 | 22 | 23 | 16 | 4 | 76 |
| Notaris acridulus (Linnaeus, 1758) | 31 | 11 | 76 | 26 | 5 | 149 |
| Notaris aethiops (Fabricius, 1792) | 0 | 1 | 0 | 1 | 0 | 2 |
| Notaris scirpi (Fabricius, 1792) | 3 | 0 | 25 | 5 | 0 | 33 |
| Thryogenes festucae (Herbst, 1795) | 4 | 0 | 0 | 0 | 0 | 4 |
| Thryogenes nereis (Paykull, 1800) | 0 | 1 | 0 | 1 | 0 | 2 |
| Thryogenes scirrhosus (Gyllenhal, 1836) | 3 | 0 | 0 | 0 | 0 | 3 |
| Tournotaris bimaculatus (Fabricius, 1787) | 16 | 2 | 10 | 0 | 0 | 28 |
| Stenopelmus rufinasus Gyllenhal, 1836 | 4 | 0 | 6 | 1 | 0 | 11 |

Curculionidae [475/312]

| | | | | | | |
|--|----|----|----|----|----|-----|
| Archarius pyrrhoceras (Marsham, 1802) | 15 | 22 | 90 | 12 | 3 | 142 |
| Archarius salicivorus (Paykull, 1792) | 23 | 13 | 83 | 21 | 11 | 151 |
| Archarius villosus (Fabricius, 1781) | 1 | 2 | 16 | 1 | 0 | 20 |
| Curculio betulae (Stephens, 1831) | 3 | 1 | 24 | 0 | 1 | 29 |
| Curculio glandium Marsham, 1802 | 11 | 1 | 40 | 0 | 0 | 52 |
| Curculio nucum Linnaeus, 1758 | 0 | 0 | 7 | 0 | 0 | 7 |
| Curculio rubidus (Gyllenhal, 1836) | 0 | 1 | 6 | 0 | 0 | 7 |
| Curculio venosus (Gravenhorst, 1807) | 1 | 1 | 33 | 2 | 0 | 37 |
| Anthonomus (Anthonomus) bituberculatus Thomson, C.G., 1868 | 4 | 3 | 10 | 4 | 3 | 24 |
| Anthonomus (Anthonomus) conspersus Desbrochers, 1868 | 0 | 5 | 1 | 0 | 0 | 6 |
| Anthonomus (Anthonomus) pedicularius (Linnaeus, 1758) | 12 | 21 | 54 | 11 | 2 | 100 |
| Anthonomus (Anthonomus) piri Kollar, 1837 | 0 | 1 | 1 | 0 | 0 | 2 |
| Anthonomus (Anthonomus) pomorum (Linnaeus, 1758) | 4 | 5 | 4 | 2 | 0 | 15 |
| Anthonomus (Anthonomus) rubi (Herbst, 1795) | 16 | 11 | 44 | 8 | 4 | 83 |
| Anthonomus (Anthonomus) ulmi (De Geer, 1775) | 0 | 3 | 4 | 1 | 4 | 12 |
| Anthonomus (Furcibus) rectirostris (Linnaeus, 1758) | 0 | 4 | 0 | 0 | 0 | 4 |
| Cionus alauda (Herbst, 1784) | 8 | 9 | 23 | 7 | 1 | 48 |
| Cionus hortulanus (Fourcroy, 1785) | 0 | 0 | 1 | 0 | 0 | 1 |
| Cionus scrophulariae (Linnaeus, 1758) | 16 | 15 | 90 | 20 | 8 | 149 |
| Cleopus pulchellus (Herbst, 1795) | 1 | 7 | 7 | 6 | 0 | 21 |
| Elleucus bipunctatus (Linnaeus, 1758) | 0 | 3 | 0 | 0 | 0 | 3 |
| Dorytomus dejeani Faust, 1882 | 5 | 1 | 28 | 4 | 0 | 38 |
| Dorytomus filirostris (Gyllenhal, 1836) | 7 | 0 | 5 | 2 | 0 | 14 |
| Dorytomus hirtipennis Bedel, 1884 | 2 | 0 | 1 | 0 | 0 | 3 |
| Dorytomus longimanus (Forster, 1771) | 9 | 5 | 18 | 3 | 0 | 35 |
| Dorytomus melanophthalmus (Paykull, 1792) | 7 | 7 | 16 | 7 | 0 | 37 |
| Dorytomus rufatus Bedel, 1888 | 11 | 11 | 32 | 4 | 0 | 58 |
| Dorytomus salicinus (Gyllenhal, 1827) | 1 | 0 | 2 | 7 | 0 | 10 |
| Dorytomus salicis Walton, 1851 | 3 | 1 | 2 | 0 | 0 | 6 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|-----|----|-----|----|----|-----|
| <i>Dorytomus taeniatus</i> (Fabricius, 1781) | 11 | 13 | 75 | 15 | 10 | 124 |
| <i>Dorytomus tortrix</i> (Linnaeus, 1761) | 6 | 2 | 24 | 3 | 1 | 36 |
| <i>Dorytomus tremulae</i> (Fabricius, 1787) | 1 | 0 | 9 | 1 | 0 | 11 |
| <i>Cleopomiarus graminis</i> (Gyllenhal, 1813) | 1 | 0 | 3 | 1 | 0 | 5 |
| <i>Cleopomiarus plantarum</i> (Germar, 1824) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Gymnetron beccabungae</i> (Linnaeus, 1761) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Gymnetron melanarium</i> (Germar, 1821) | 3 | 0 | 0 | 0 | 0 | 3 |
| <i>Gymnetron veronicae</i> (Germar, 1821) | 1 | 2 | 0 | 1 | 0 | 4 |
| <i>Gymnetron villosulum</i> Gyllenhal, 1838 | 11 | 0 | 2 | 1 | 0 | 14 |
| <i>Mecinus labilis</i> (Herbst, 1795) | 1 | 5 | 1 | 1 | 0 | 8 |
| <i>Mecinus pascuorum</i> (Gyllenhal, 1813) | 31 | 25 | 71 | 7 | 2 | 136 |
| <i>Mecinus pyraster</i> (Herbst, 1795) | 24 | 8 | 41 | 7 | 1 | 81 |
| <i>Miarus campanulae</i> (Linnaeus, 1767) | 2 | 2 | 10 | 0 | 0 | 14 |
| <i>Rhinusa antirrhini</i> (Paykull, 1800) | 6 | 0 | 32 | 7 | 0 | 45 |
| <i>Rhinusa linariae</i> (Panzer, 1795) | 0 | 3 | 1 | 0 | 0 | 4 |
| <i>Isochnus foliorum</i> (Müller, O.F., 1764) | 0 | 2 | 2 | 1 | 1 | 6 |
| <i>Orchestes</i> (<i>Orchestes</i>) <i>alni</i> (Linnaeus, 1758) | 8 | 0 | 2 | 3 | 0 | 13 |
| <i>Orchestes</i> (<i>Orchestes</i>) <i>pilosus</i> (Fabricius, 1781) | 1 | 7 | 2 | 0 | 0 | 10 |
| <i>Orchestes</i> (<i>Orchestes</i>) <i>quercus</i> (Linnaeus, 1758) | 24 | 24 | 33 | 12 | 0 | 93 |
| <i>Orchestes</i> (<i>Orchestes</i>) <i>rusci</i> (Herbst, 1795) | 17 | 17 | 63 | 8 | 2 | 107 |
| <i>Orchestes</i> (<i>Orchestes</i>) <i>signifer</i> (Creutzer, 1799) | 8 | 7 | 11 | 5 | 0 | 31 |
| <i>Orchestes</i> (<i>Orchestes</i>) <i>testaceus</i> (Müller, O.F., 1776) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Orchestes</i> (<i>Salius</i>) <i>fagi</i> (Linnaeus, 1758) | 14 | 31 | 109 | 34 | 2 | 190 |
| <i>Pseudorchestes pratensis</i> (Germar, 1821) | 0 | 0 | 1 | 1 | 0 | 2 |
| <i>Rhamphus oxyacanthae</i> (Marsham, 1802) | 6 | 4 | 31 | 3 | 1 | 45 |
| <i>Rhamphus pulicarius</i> (Herbst, 1795) | 10 | 16 | 44 | 3 | 2 | 75 |
| <i>Tachyerges salicis</i> (Linnaeus, 1758) | 4 | 15 | 12 | 6 | 1 | 38 |
| <i>Tachyerges stigma</i> (Germar, 1821) | 0 | 1 | 1 | 1 | 0 | 3 |
| <i>Smicronyx reichi</i> (Gyllenhal, 1836) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Orthochaetes setiger</i> (Beck, 1817) | 4 | 5 | 3 | 2 | 0 | 14 |
| <i>Sibinia primita</i> (Herbst, 1795) | 0 | 0 | 3 | 0 | 0 | 3 |
| <i>Tychius lineatulus</i> Stephens, 1831 | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Tychius meliloti</i> Stephens, 1831 | 7 | 2 | 17 | 1 | 0 | 27 |
| <i>Tychius picirostris</i> (Fabricius, 1787) | 38 | 23 | 169 | 22 | 11 | 263 |
| <i>Tychius schneideri</i> (Herbst, 1795) | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Bagous</i> (<i>Bagous</i>) <i>limosus</i> (Gyllenhal, 1827) | 0 | 1 | 4 | 1 | 0 | 6 |
| <i>Bagous</i> (<i>Bagous</i>) <i>lutulosus</i> (Gyllenhal, 1827) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Bagous</i> (<i>Bagous</i>) <i>subcarinatus</i> Gyllenhal, 1836 | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Bagous</i> (<i>Bagous</i>) <i>tempestivus</i> (Herbst, 1795) | 2 | 1 | 1 | 1 | 0 | 5 |
| <i>Bagous</i> (<i>Ephimeropus</i>) <i>petro</i> (Herbst, 1795) | 0 | 0 | 0 | 3 | 0 | 3 |
| <i>Bagous</i> (<i>Hydronomus</i>) <i>alismatis</i> (Marsham, 1802) | 3 | 1 | 13 | 2 | 0 | 19 |
| <i>Aulacobaris lepidii</i> (Germar, 1824) | 0 | 3 | 0 | 1 | 0 | 4 |
| <i>Aulacobaris picicornis</i> (Marsham, 1802) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Melanobaris laticollis</i> (Marsham, 1802) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Limnobaris dolorosa</i> (Goeze, 1777) | 10 | 14 | 16 | 8 | 0 | 48 |
| <i>Limnobaris t-album</i> (Linnaeus, 1758) | 0 | 6 | 2 | 3 | 1 | 12 |
| <i>Amalorrhynchus melanarius</i> (Stephens, 1831) | 0 | 0 | 9 | 0 | 0 | 9 |
| <i>Calosirus terminatus</i> (Herbst, 1795) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Ceutorhynchus alliariae</i> Brisout, 1860 | 0 | 2 | 3 | 2 | 0 | 7 |
| <i>Ceutorhynchus assimilis</i> (Paykull, 1792) | 133 | 65 | 201 | 47 | 11 | 457 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| <i>Ceutorhynchus atomus</i> Boheman, 1845 | 1 | 0 | 0 | 1 | 0 | 2 |
| <i>Ceutorhynchus chalybaeus</i> Germar, 1824 | 0 | 0 | 1 | 1 | 0 | 2 |
| <i>Ceutorhynchus cochleariae</i> (Gyllenhal, 1813) | 3 | 2 | 2 | 2 | 0 | 9 |
| <i>Ceutorhynchus constrictus</i> (Marsham, 1802) | 1 | 4 | 3 | 1 | 0 | 9 |
| <i>Ceutorhynchus contractus</i> (Marsham, 1802) | 20 | 15 | 19 | 13 | 2 | 69 |
| <i>Ceutorhynchus erysimi</i> (Fabricius, 1787) | 22 | 26 | 32 | 28 | 5 | 113 |
| <i>Ceutorhynchus hirtulus</i> Germar, 1824 | 0 | 1 | 0 | 1 | 0 | 2 |
| <i>Ceutorhynchus insularis</i> Dieckmann, 1971 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Ceutorhynchus pallidactylus</i> (Marsham, 1802) | 68 | 24 | 132 | 31 | 19 | 274 |
| <i>Ceutorhynchus parvulus</i> Brisout, Ch., 1869 | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Ceutorhynchus picitarsis</i> Gyllenhal, 1837 | 8 | 0 | 10 | 0 | 0 | 18 |
| <i>Ceutorhynchus pumilio</i> (Gyllenhal, 1827) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Ceutorhynchus pyrrhorhynchus</i> (Marsham, 1802) | 5 | 2 | 9 | 1 | 0 | 17 |
| <i>Ceutorhynchus resedae</i> (Marsham, 1802) | 2 | 0 | 2 | 0 | 0 | 4 |
| <i>Ceutorhynchus sulcicollis</i> (Paykull, 1800) | 1 | 0 | 0 | 1 | 0 | 2 |
| <i>Ceutorhynchus turbatus</i> Schultze, 1903 | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Ceutorhynchus typhae</i> (Herbst, 1795) | 59 | 41 | 47 | 24 | 11 | 182 |
| <i>Coeliodes rana</i> (Fabricius, 1787) | 6 | 18 | 20 | 6 | 0 | 50 |
| <i>Coeliodes ruber</i> (Marsham, 1802) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Coeliodes transversealbofasciatus</i> (Goeze, 1777) | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Coeliodinus rubicundus</i> (Herbst, 1795) | 8 | 18 | 30 | 7 | 0 | 63 |
| <i>Datonychus angulosus</i> (Boheman, 1845) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Datonychus melanostictus</i> (Marsham, 1802) | 0 | 0 | 5 | 3 | 0 | 8 |
| <i>Datonychus urticae</i> (Boheman, 1845) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Drupenatus nasturtii</i> (Germar, 1824) | 6 | 0 | 10 | 0 | 0 | 16 |
| <i>Glocianus distinctus</i> (Brisout, Ch., 1870) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Glocianus punctiger</i> (Gyllenhal, 1837) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Hadroplontus litura</i> (Fabricius, 1775) | 16 | 10 | 44 | 11 | 5 | 86 |
| <i>Hadroplontus trimaculatus</i> (Fabricius, 1775) | 0 | 3 | 0 | 0 | 0 | 3 |
| <i>Micrelus ericae</i> (Gyllenhal, 1813) | 4 | 18 | 42 | 4 | 6 | 74 |
| <i>Microplontus campestris</i> (Gyllenhal, 1837) | 1 | 0 | 10 | 1 | 0 | 12 |
| <i>Microplontus rugulosus</i> (Herbst, 1795) | 17 | 1 | 12 | 3 | 0 | 33 |
| <i>Mogulones asperifoliarum</i> (Gyllenhal, 1813) | 12 | 5 | 7 | 2 | 2 | 28 |
| <i>Nedus quadrimaculatus</i> (Linnaeus, 1758) | 62 | 33 | 213 | 65 | 15 | 388 |
| <i>Parethelcus pollinarius</i> (Forster, 1771) | 36 | 19 | 91 | 15 | 0 | 161 |
| <i>Poophagus sisymbrii</i> (Fabricius, 1777) | 9 | 7 | 12 | 12 | 1 | 41 |
| <i>Sirocalodes depressicollis</i> (Gyllenhal, 1813) | 2 | 1 | 0 | 3 | 0 | 6 |
| <i>Sirocalodes mixtus</i> (Mulsant & Rey, 1858) | 2 | 0 | 1 | 0 | 0 | 3 |
| <i>Sirocalodes quercicola</i> (Paykull, 1792) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Stenocarus ruficornis</i> (Stephens, 1831) | 2 | 7 | 0 | 2 | 0 | 11 |
| <i>Thamicolus viduatus</i> (Gyllenhal, 1813) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Trichosirocalus dawsoni</i> (Brisout, Ch., 1869) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Trichosirocalus horridus</i> (Panzer, 1801) | 0 | 2 | 1 | 0 | 0 | 3 |
| <i>Trichosirocalus thalhammeri</i> (Schultze, 1906) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Trichosirocalus troglodytes</i> (Fabricius, 1787) | 53 | 37 | 100 | 14 | 0 | 204 |
| <i>Zacladus exiguus</i> (Olivier, 1807) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Zacladus geranii</i> (Paykull, 1800) | 13 | 9 | 14 | 20 | 11 | 67 |
| <i>Amalus scortillum</i> (Herbst, 1795) | 7 | 1 | 7 | 5 | 0 | 20 |
| <i>Eubrychius velutus</i> (Beck, 1817) | 0 | 0 | 6 | 2 | 0 | 8 |
| <i>Neophytobius muricatus</i> (Brisout, Ch., 1867) | 0 | 1 | 0 | 0 | 0 | 1 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|-----|----|----|-----|
| <i>Neophytobius quadrinodosus</i> (Gyllenhal, 1813) | 1 | 2 | 0 | 0 | 0 | 3 |
| <i>Pelenomus canaliculatus</i> (Fåhraeus, 1843) | 0 | 1 | 1 | 1 | 0 | 3 |
| <i>Pelenomus comari</i> (Herbst, 1795) | 2 | 3 | 2 | 8 | 0 | 15 |
| <i>Pelenomus quadricorniger</i> (Colonnelli, 1986) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Pelenomus quadrituberculatus</i> (Fabricius, 1787) | 9 | 8 | 13 | 10 | 0 | 40 |
| <i>Phytobius leucogaster</i> (Marsham, 1802) | 1 | 2 | 3 | 0 | 0 | 6 |
| <i>Rhinoncus bruchoides</i> (Herbst, 1784) | 2 | 1 | 0 | 0 | 0 | 3 |
| <i>Rhinoncus castor</i> (Fabricius, 1792) | 16 | 8 | 27 | 6 | 2 | 59 |
| <i>Rhinoncus inconspectus</i> (Herbst, 1795) | 17 | 1 | 10 | 2 | 0 | 30 |
| <i>Rhinoncus pericarpus</i> (Linnaeus, 1758) | 49 | 26 | 108 | 29 | 9 | 221 |
| <i>Rhinoncus perpendicularis</i> (Reich, 1797) | 25 | 6 | 31 | 10 | 2 | 74 |
| <i>Rhopalomesites tardyi</i> (Curtis, 1825) | 0 | 7 | 3 | 1 | 0 | 11 |
| <i>Pselactus spadix</i> (Herbst, 1795) | 17 | 0 | 0 | 0 | 0 | 17 |
| <i>Pseudophloeophagus aeneopiceus</i> (Boheman, 1845) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Euophryum confine</i> (Broun, 1881) | 17 | 7 | 59 | 7 | 0 | 90 |
| <i>Euophryum rufum</i> (Broun, 1880) | 0 | 0 | 1 | 1 | 0 | 2 |
| <i>Pentarthrum huttoni</i> Wollaston, 1854 | 2 | 0 | 8 | 2 | 0 | 12 |
| <i>Rhyncolus ater</i> (Linnaeus, 1758) | 0 | 2 | 2 | 0 | 0 | 4 |
| <i>Phloeophagus lignarius</i> (Marsham, 1802) | 2 | 0 | 8 | 1 | 0 | 11 |
| <i>Cryptorhynchus lapathi</i> (Linnaeus, 1758) | 1 | 3 | 18 | 3 | 2 | 27 |
| <i>Acalles misellus</i> Boheman, 1844 | 5 | 2 | 3 | 0 | 0 | 10 |
| <i>Acalles ptinoides</i> (Marsham, 1802) | 0 | 2 | 14 | 2 | 4 | 22 |
| <i>Kykliaocalles roboris</i> (Curtis, 1835) | 0 | 7 | 9 | 4 | 0 | 20 |
| <i>Gronops lunatus</i> (Fabricius, 1775) | 1 | 0 | 4 | 0 | 0 | 5 |
| <i>Graptus triguttatus</i> (Fabricius, 1775) | 10 | 7 | 1 | 9 | 3 | 30 |
| <i>Neliocarus faber</i> (Herbst, 1784) | 1 | 3 | 3 | 1 | 0 | 8 |
| <i>Neliocarus nebulosus</i> (Stephens, 1831) | 2 | 20 | 12 | 4 | 0 | 38 |
| <i>Neliocarus sus</i> (Stephens, 1831) | 11 | 27 | 16 | 4 | 3 | 61 |
| <i>Strophosoma capitatum</i> (De Geer, 1775) | 1 | 2 | 5 | 3 | 0 | 11 |
| <i>Strophosoma melanogrammum</i> (Forster, 1771) | 26 | 42 | 146 | 17 | 6 | 237 |
| <i>Attactagenus plumbeus</i> (Marsham, 1802) | 6 | 9 | 7 | 1 | 1 | 24 |
| <i>Philopeton plagiatum</i> (Schaller, 1783) | 37 | 5 | 5 | 0 | 0 | 47 |
| <i>Barynotus moerens</i> (Fabricius, 1792) | 2 | 12 | 27 | 11 | 4 | 56 |
| <i>Barynotus obscurus</i> (Fabricius, 1775) | 8 | 11 | 30 | 8 | 2 | 59 |
| <i>Barynotus squamosus</i> Germar, 1824 | 0 | 4 | 10 | 8 | 5 | 27 |
| <i>Omiamima mollina</i> (Boheman, 1834) | 10 | 5 | 15 | 2 | 0 | 32 |
| <i>Otiorhynchus</i> (<i>Otiorhynchus</i>) <i>clavipes</i> (Bonsdorff, 1785) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Otiorhynchus</i> (<i>Otiorhynchus</i>) <i>ligneus</i> (Olivier, 1807) | 1 | 1 | 2 | 3 | 0 | 7 |
| <i>Otiorhynchus</i> (<i>Otiorhynchus</i>) <i>nodosus</i> (Müller, O. F. 1764) | 0 | 4 | 3 | 3 | 2 | 12 |
| <i>Otiorhynchus</i> (<i>Otiorhynchus</i>) <i>porcatus</i> (Herbst, 1795) | 0 | 2 | 2 | 3 | 0 | 7 |
| <i>Otiorhynchus</i> (<i>Otiorhynchus</i>) <i>raucus</i> (Fabricius, 1777) | 0 | 1 | 1 | 2 | 0 | 4 |
| <i>Otiorhynchus</i> (<i>Otiorhynchus</i>) <i>rugosostriatus</i> (Goeze, 1777) | 2 | 1 | 10 | 1 | 0 | 14 |
| <i>Otiorhynchus</i> (<i>Arammichnus</i>) <i>atroapterus</i> (De Geer, 1775) | 0 | 3 | 1 | 0 | 0 | 4 |
| <i>Otiorhynchus</i> (<i>Arammichnus</i>) <i>ligustici</i> (Linnaeus, 1758) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Otiorhynchus</i> (<i>Dorymerus</i>) <i>rugifrons</i> (Gyllenhal, 1813) | 1 | 3 | 4 | 1 | 3 | 12 |
| <i>Otiorhynchus</i> (<i>Dorymerus</i>) <i>singularis</i> (Linnaeus, 1767) | 31 | 52 | 198 | 31 | 12 | 324 |
| <i>Otiorhynchus</i> (<i>Dorymerus</i>) <i>sulcatus</i> (Fabricius, 1775) | 5 | 13 | 70 | 11 | 6 | 105 |
| <i>Otiorhynchus</i> (<i>Tournieria</i>) <i>desertus</i> Rosenhauer, 1847 | 0 | 0 | 2 | 0 | 1 | 3 |
| <i>Otiorhynchus</i> (<i>Tournieria</i>) <i>ovatus</i> (Linnaeus, 1758) | 39 | 6 | 19 | 9 | 1 | 74 |
| <i>Caenopsis fissirostris</i> (Walton, 1847) | 0 | 0 | 0 | 1 | 0 | 1 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|-----|----|-----|-----|----|-----|
| Phyllobius (Phyllobius) pyri (Linnaeus, 1758) | 57 | 59 | 267 | 38 | 19 | 440 |
| Phyllobius (Phyllobius) vespertinus (Fabricius, 1792) | 0 | 3 | 1 | 0 | 0 | 4 |
| Phyllobius (Dieletus) argentatus (Linnaeus, 1758) | 16 | 44 | 192 | 24 | 17 | 293 |
| Phyllobius (Metaphyllobius) glaucus (Scopoli, 1763) | 6 | 41 | 109 | 18 | 9 | 183 |
| Phyllobius (Metaphyllobius) pomaceus Gyllenhal, 1834 | 17 | 30 | 117 | 28 | 5 | 197 |
| Phyllobius (Nemoicus) oblongus (Linnaeus, 1758) | 7 | 20 | 135 | 27 | 12 | 201 |
| Phyllobius (Parnemoicus) roboretanus Gredler, 1882 | 63 | 49 | 202 | 39 | 13 | 366 |
| Phyllobius (Parnemoicus) viridicollis (Fabricius, 1801) | 5 | 7 | 2 | 21 | 1 | 36 |
| Phyllobius (Pterygorrhynchus) maculicornis Germar, 1824 | 8 | 16 | 76 | 13 | 0 | 113 |
| Phyllobius (Subphyllobius) virideaeris (Laicharting, 1781) | 2 | 0 | 3 | 0 | 0 | 5 |
| Liophloeus tessulatus (Müller, O.F., 1776) | 22 | 11 | 32 | 7 | 2 | 74 |
| Polydrusus (Polydrusus) tereticollis (De Geer, 1775) | 0 | 29 | 58 | 5 | 0 | 92 |
| Polydrusus (Chrysophis) formosus (Mayer, 1779) | 0 | 0 | 2 | 0 | 0 | 2 |
| Polydrusus (Eudipnus) mollis (Ström, 1768) | 0 | 27 | 9 | 4 | 4 | 44 |
| Polydrusus (Eurodrusus) confluens Stephens, 1831 | 1 | 3 | 2 | 0 | 0 | 6 |
| Polydrusus (Eustolus) flavipes (De Geer, 1775) | 1 | 0 | 0 | 0 | 0 | 1 |
| Polydrusus (Eustolus) pterygomalis Boheman, 1840 | 4 | 43 | 19 | 15 | 4 | 85 |
| Polydrusus (Neoeustolus) cervinus (Linnaeus, 1758) | 15 | 27 | 80 | 10 | 4 | 136 |
| Polydrusus (Neoeustolus) pilosus Gredler, 1866 | 1 | 9 | 3 | 2 | 2 | 17 |
| Polydrusus (Neoeustolus) pulchellus Stephens, 1831 | 3 | 0 | 0 | 0 | 0 | 3 |
| Barypeithes (Barypeithes) sulcifrons (Boheman, 1843) | 2 | 3 | 7 | 0 | 0 | 12 |
| Barypeithes (Exomias) araneiformis (Schrank, 1781) | 8 | 22 | 41 | 18 | 5 | 94 |
| Barypeithes (Exomias) pellucidus (Boheman, 1834) | 26 | 43 | 205 | 41 | 11 | 326 |
| Barypeithes (Exomias) pyrenaeus Seidlitz, 1868 | 0 | 0 | 3 | 0 | 0 | 3 |
| Brachysomus echinatus (Bonsdorff, 1785) | 5 | 11 | 8 | 10 | 1 | 35 |
| Sciaphilus asperatus (Bonsdorff, 1785) | 4 | 10 | 42 | 15 | 2 | 73 |
| Sitona (Sitona) ambiguus Gyllenhal, 1834 | 0 | 3 | 1 | 1 | 0 | 5 |
| Sitona (Sitona) cambricus Stephens, 1831 | 0 | 0 | 7 | 1 | 0 | 8 |
| Sitona (Sitona) cylindricollis (Fåhraeus, 1840) | 8 | 2 | 7 | 0 | 1 | 18 |
| Sitona (Sitona) hispidulus (Fabricius, 1777) | 21 | 22 | 66 | 19 | 8 | 136 |
| Sitona (Sitona) humeralis Stephens, 1831 | 10 | 3 | 16 | 8 | 1 | 38 |
| Sitona (Sitona) lepidus Gyllenhal, 1834 | 24 | 34 | 93 | 30 | 12 | 193 |
| Sitona (Sitona) lineatus (Linnaeus, 1758) | 163 | 95 | 395 | 101 | 26 | 780 |
| Sitona (Sitona) lineellus (Bonsdorff, 1785) | 5 | 3 | 3 | 2 | 0 | 13 |
| Sitona (Sitona) macularius (Marsham, 1802) | 2 | 2 | 0 | 0 | 0 | 4 |
| Sitona (Sitona) ononidis Sharp, 1866 | 0 | 1 | 2 | 0 | 0 | 3 |
| Sitona (Sitona) puncticollis Stephens, 1831 | 9 | 10 | 8 | 3 | 0 | 30 |
| Sitona (Sitona) regensteinensis (Herbst, 1797) | 13 | 9 | 89 | 7 | 3 | 121 |
| Sitona (Sitona) striatellus Gyllenhal, 1834 | 10 | 3 | 13 | 6 | 1 | 33 |
| Sitona (Sitona) sulcifrons (Thunberg, 1798) | 6 | 7 | 9 | 9 | 2 | 33 |
| Sitona (Sitona) suturalis Stephens, 1831 | 20 | 23 | 54 | 15 | 5 | 117 |
| Sitona (Sitona) waterhousei Walton, 1846 | 0 | 1 | 1 | 0 | 0 | 2 |
| Sitona (Charagmus) griseus (Fabricius, 1775) | 3 | 2 | 5 | 0 | 0 | 10 |
| Tanymecus palliatus (Fabricius, 1787) | 2 | 5 | 1 | 1 | 0 | 9 |
| Trachyploeus alternans Gyllenhal, 1834 | 0 | 2 | 0 | 0 | 0 | 2 |
| Trachyploeus angustisetulus Hansen, 1915 | 0 | 0 | 1 | 0 | 0 | 1 |
| Trachyploeus aristatus (Gyllenhal, 1827) | 1 | 1 | 4 | 0 | 0 | 6 |
| Trachyploeus asperatus Boheman, 1843 | 0 | 2 | 1 | 0 | 0 | 3 |
| Trachyploeus bifoveolatus (Beck, 1817) | 1 | 2 | 3 | 0 | 0 | 6 |
| Trachyploeus scabriculus (Linnaeus, 1771) | 2 | 0 | 1 | 0 | 0 | 3 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|--|----|----|-----|----|----|-----|
| <i>Cathormiocerus myrmecophilus</i> (Seidlitz, 1868) | 1 | 0 | 0 | 0 | 0 | 1 |
| <i>Tropiphorus elevatus</i> (Herbst, 1795) | 0 | 1 | 2 | 1 | 0 | 4 |
| <i>Tropiphorus obtusus</i> (Bonsdorff, 1785) | 0 | 1 | 1 | 3 | 0 | 5 |
| <i>Tropiphorus terricola</i> (Newman, 1838) | 2 | 7 | 8 | 2 | 5 | 24 |
| <i>Hypera</i> (<i>Hypera</i>) <i>arator</i> (Linnaeus, 1758) | 10 | 2 | 5 | 3 | 0 | 20 |
| <i>Hypera</i> (<i>Hypera</i>) <i>fuscocinerea</i> (Marsham, 1802) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Hypera</i> (<i>Hypera</i>) <i>nigrirostris</i> (Fabricius, 1775) | 27 | 10 | 38 | 15 | 0 | 90 |
| <i>Hypera</i> (<i>Hypera</i>) <i>plantaginis</i> (De Geer, 1775) | 2 | 8 | 11 | 9 | 0 | 30 |
| <i>Hypera</i> (<i>Hypera</i>) <i>postica</i> (Gyllenhal, 1813) | 17 | 9 | 12 | 6 | 0 | 44 |
| <i>Hypera</i> (<i>Hypera</i>) <i>suspiciosa</i> (Herbst, 1795) | 1 | 3 | 5 | 3 | 3 | 15 |
| <i>Hypera</i> (<i>Hypera</i>) <i>venusta</i> (Fabricius, 1781) | 2 | 1 | 5 | 0 | 0 | 8 |
| <i>Hypera</i> (<i>Antidonus</i>) <i>dauci</i> (Olivier, 1807) | 3 | 1 | 0 | 0 | 0 | 4 |
| <i>Hypera</i> (<i>Antidonus</i>) <i>zoilus</i> (Scopoli, 1763) | 7 | 8 | 5 | 11 | 0 | 31 |
| <i>Hypera</i> (<i>Boreohypera</i>) <i>diversipunctata</i> (Schrank, 1798) | 3 | 0 | 1 | 2 | 0 | 6 |
| <i>Hypera</i> (<i>Dapalinus</i>) <i>meles</i> (Fabricius, 1792) | 0 | 0 | 1 | 1 | 0 | 2 |
| <i>Hypera</i> (<i>Eirinomorphus</i>) <i>pollux</i> (Fabricius, 1801) | 10 | 2 | 3 | 3 | 0 | 18 |
| <i>Hypera</i> (<i>Eirinomorphus</i>) <i>rumicis</i> (Linnaeus, 1758) | 4 | 3 | 7 | 4 | 1 | 19 |
| <i>Limobius borealis</i> (Paykull, 1792) | 2 | 3 | 0 | 5 | 2 | 12 |
| <i>Lixus</i> (<i>Lixus</i>) <i>paraplecticus</i> (Linnaeus, 1758) | 1 | 0 | 1 | 0 | 0 | 2 |
| <i>Lixus</i> (<i>Dilixellus</i>) <i>angustatus</i> (Fabricius, 1775) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Cleonis pigra</i> (Scopoli, 1763) | 15 | 2 | 0 | 0 | 0 | 17 |
| <i>Magdalis</i> (<i>Magdalis</i>) <i>duplicata</i> Germar, 1818 | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Magdalis</i> (<i>Magdalis</i>) <i>phlegmatica</i> (Herbst, 1797) | 0 | 0 | 0 | 2 | 0 | 2 |
| <i>Magdalis</i> (<i>Edo</i>) <i>ruficornis</i> (Linnaeus, 1758) | 3 | 5 | 6 | 6 | 1 | 21 |
| <i>Magdalis</i> (<i>Odontomagdalis</i>) <i>armigera</i> (Fourcroy, 1785) | 3 | 7 | 12 | 7 | 2 | 31 |
| <i>Magdalis</i> (<i>Odontomagdalis</i>) <i>carbonaria</i> (Linnaeus, 1758) | 2 | 3 | 6 | 4 | 1 | 16 |
| <i>Magdalis</i> (<i>Panus</i>) <i>barbicornis</i> (Latreille, 1804) | 1 | 0 | 3 | 0 | 0 | 4 |
| <i>Magdalis</i> (<i>Porrothus</i>) <i>cerasi</i> (Linnaeus, 1758) | 3 | 1 | 11 | 5 | 0 | 20 |
| <i>Liparus coronatus</i> (Goeze, 1777) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Leiosoma deflexum</i> (Panzer, 1795) | 11 | 25 | 132 | 21 | 12 | 201 |
| <i>Leiosoma oblongulum</i> Boheman, 1842 | 0 | 0 | 2 | 2 | 0 | 4 |
| <i>Anoplus plantaris</i> (Naezen, 1794) | 4 | 16 | 26 | 2 | 0 | 48 |
| <i>Anoplus roboris</i> Suffrian, 1840 | 0 | 1 | 2 | 1 | 0 | 4 |
| <i>Hylobius</i> (<i>Callirus</i>) <i>abietis</i> (Linnaeus, 1758) | 6 | 5 | 11 | 6 | 1 | 29 |
| <i>Pissodes castaneus</i> (De Geer, 1775) | 3 | 2 | 1 | 1 | 0 | 7 |
| <i>Pissodes pini</i> (Linnaeus, 1758) | 4 | 6 | 9 | 3 | 0 | 22 |
| <i>Trachodes hispidus</i> (Linnaeus, 1758) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Orobitis cyaneus</i> (Linnaeus, 1758) | 1 | 7 | 1 | 2 | 0 | 11 |
| <i>Scolytus intricatus</i> (Ratzeburg, 1837) | 1 | 9 | 10 | 7 | 0 | 27 |
| <i>Scolytus mali</i> (Bechstein & Scharfenberg, 1805) | 0 | 3 | 3 | 4 | 0 | 10 |
| <i>Scolytus multistriatus</i> (Marsham, 1802) | 1 | 0 | 17 | 0 | 1 | 19 |
| <i>Scolytus ratzeburgi</i> Janson, 1856 | 0 | 1 | 1 | 0 | 0 | 2 |
| <i>Scolytus rugulosus</i> (Müller, P.W.J., 1818) | 1 | 0 | 3 | 2 | 0 | 6 |
| <i>Scolytus scolytus</i> (Fabricius, 1775) | 1 | 9 | 32 | 7 | 0 | 49 |
| <i>Pityophthorus lichtensteinii</i> (Ratzeburg, 1837) | 1 | 0 | 2 | 0 | 0 | 3 |
| <i>Pityophthorus pubescens</i> (Marsham, 1802) | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Cryphalus asperatus</i> (Gyllenhal, 1813) | 2 | 5 | 0 | 3 | 0 | 10 |
| <i>Ernoporicus caucasicus</i> (Lindemann, 1876) | 0 | 2 | 0 | 1 | 0 | 3 |
| <i>Ernoporicus fagi</i> (Fabricius, 1798) | 0 | 0 | 0 | 2 | 0 | 2 |
| <i>Ernoporus tiliae</i> (Panzer, 1793) | 0 | 0 | 1 | 1 | 0 | 2 |

Yorkshire Coleoptera checklist 2011

| | | | | | | |
|---|----|----|----|----|---|----|
| <i>Dryocoetes alni</i> (Georg, 1856) | 0 | 0 | 5 | 5 | 2 | 12 |
| <i>Dryocoetes autographus</i> (Ratzeburg, 1837) | 2 | 7 | 0 | 4 | 0 | 13 |
| <i>Dryocoetes villosus</i> (Fabricius, 1792) | 1 | 8 | 23 | 6 | 0 | 38 |
| <i>Taphrorhynchus bicolor</i> (Herbst, 1793) | 0 | 0 | 0 | 1 | 0 | 1 |
| <i>Ips acuminatus</i> (Gyllenhal, 1827) | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Ips sexdentatus</i> (Boerner, 1776) | 0 | 0 | 1 | 0 | 0 | 1 |
| <i>Ips typographus</i> (Linnaeus, 1758) | 1 | 1 | 0 | 0 | 0 | 2 |
| <i>Orthotomicus laricis</i> (Fabricius, 1792) | 2 | 1 | 0 | 1 | 0 | 4 |
| <i>Pityogenes bidentatus</i> (Herbst, 1783) | 5 | 7 | 9 | 3 | 1 | 25 |
| <i>Pityogenes chalcographus</i> (Linnaeus, 1761) | 1 | 4 | 1 | 0 | 0 | 6 |
| <i>Pityogenes quadridens</i> (Hartig, 1834) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Xyleborinus saxesenii</i> (Ratzeburg, 1837) | 2 | 4 | 4 | 1 | 0 | 11 |
| <i>Xyleborus dispar</i> (Fabricius, 1792) | 0 | 1 | 6 | 1 | 0 | 8 |
| <i>Xyleborus dryographus</i> (Ratzeburg, 1837) | 0 | 0 | 2 | 0 | 0 | 2 |
| <i>Trypodendron domesticum</i> (Linnaeus, 1758) | 2 | 10 | 26 | 11 | 1 | 50 |
| <i>Trypodendron lineatum</i> (Olivier, 1795) | 0 | 1 | 4 | 1 | 0 | 6 |
| <i>Trypodendron signatum</i> (Fabricius, 1792) | 0 | 3 | 6 | 0 | 0 | 9 |
| <i>Hylesinus crenatus</i> (Fabricius, 1787) | 1 | 6 | 12 | 5 | 3 | 27 |
| <i>Hylesinus toranio</i> (Danthoine in Bernard, 1788) | 3 | 2 | 5 | 2 | 0 | 12 |
| <i>Hylesinus varius</i> (Fabricius, 1775) | 5 | 9 | 25 | 6 | 4 | 49 |
| <i>Hylastinus obscurus</i> (Marsham, 1802) | 1 | 7 | 7 | 0 | 1 | 16 |
| <i>Kissophagus hederæ</i> (Schmitt, 1843) | 0 | 2 | 0 | 0 | 0 | 2 |
| <i>Pteleobius vittatus</i> (Fabricius, 1787) | 1 | 0 | 4 | 1 | 0 | 6 |
| <i>Hylastes ater</i> (Fabricius, 1792) | 2 | 5 | 7 | 2 | 1 | 17 |
| <i>Hylastes brunneus</i> Erichson, 1836 | 0 | 1 | 1 | 0 | 1 | 3 |
| <i>Hylastes cunicularius</i> Erichson, 1836 | 0 | 2 | 0 | 1 | 0 | 3 |
| <i>Hylastes opacus</i> Erichson, 1836 | 2 | 2 | 2 | 0 | 0 | 6 |
| <i>Hylurgops palliatus</i> (Gyllenhal, 1813) | 7 | 9 | 12 | 3 | 0 | 31 |
| <i>Phloeotribus rhododactylus</i> (Marsham, 1802) | 7 | 4 | 6 | 0 | 0 | 17 |
| <i>Polygraphus poligraphus</i> (Linnaeus, 1758) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Tomicus piniperda</i> (Linnaeus, 1758) | 2 | 8 | 10 | 3 | 0 | 23 |
| <i>Xylechinus pilosus</i> (Ratzeburg, 1837) | 0 | 1 | 0 | 0 | 0 | 1 |
| <i>Tanysphyrus lemnae</i> (Paykull, 1792) | 12 | 7 | 24 | 13 | 0 | 56 |

Platypodidae [1/1]

| | | | | | | |
|---|---|---|---|---|---|---|
| <i>Platypus cylindrus</i> (Fabricius, 1792) | 1 | 1 | 2 | 0 | 0 | 4 |
|---|---|---|---|---|---|---|

It has not been possible to include a column containing species status in this version of the checklist owing to software problems.

Please pass comments on the content of this checklist either to R J Marsh or M L Denton.